


Felix Klein Famous Problems of Elementary Geometry




A diagram illustrating the quadrature of the circle. It features a large blue rectangle on the left and a yellow circle on the right. A black line segment connects the top-right corner of the rectangle to the center of the circle, and another black line segment connects the bottom-right corner of the rectangle to the center of the circle. The text "QUADRATURE OF THE CIRCLE" is printed in white capital letters across the bottom of the blue rectangle.

QUADRATURE OF THE CIRCLE



A diagram illustrating the trisection of an angle. It features a large green rectangle on the left and a yellow sector of a circle on the right. The sector is divided into three equal parts by two black lines that meet at the center of the circle. The text "TRISECTION OF AN ANGLE" is printed in white capital letters across the bottom of the green rectangle.

TRISECTION OF AN ANGLE



A diagram illustrating the duplication of the cube. It features three 3D cubes. The first cube is yellow, the second is green, and the third is blue. The text "513" and "672 F" is printed in black on a white rectangular background to the left of the cubes. The text "DUPLICATION OF THE CUBE" is printed in white capital letters across the bottom of the cubes.

513
672 F

DUPLICATION OF THE CUBE

DOVER BOOKS ON ELEMENTARY AND INTERMEDIATE MATHEMATICS

- Teach Yourself Algebra*, P. Abbott. Clothbound \$2.00
Teach Yourself Calculus, P. Abbott. Clothbound \$2.00
Teach Yourself Geometry, P. Abbott. Clothbound \$2.00
Teach Yourself Trigonometry, P. Abbott. Clothbound \$2.00
Handbook of Mathematical Functions: with Formulas, Graphs and Mathematical Tables, Milton Abramowitz and Irene A. Stegun. \$4.00
A Short Account of the History of Mathematics, W. W. Rouse Ball. \$2.25
Recreations in the Theory of Numbers: The Queen of Mathematics Entertains, Albert Beiler. \$2.00
Probabilities and Life, Emile Borel. \$1.00
Arithmetical Excursions: An Enrichment of Elementary Mathematics, Henry Bowers and Joan E. Bowers. \$1.65
Mathematical Tables and Formulas, Robert D. Carmichael and Edwin R. Smith. \$1.25
Textbook of Algebra, G. Chrystal. Two-volume set \$4.70
The Mathematics of Great Amateurs, Julian Lowell Coolidge. \$1.50
Elementary Statistics with Applications in Medicine and the Biological Sciences, Frederick E. Croxton. \$2.00
Statistical Adjustment of Data, W. Edwards Deming. \$2.00
Contemporary Geometry, André Delachet. \$1.00
100 Great Problems of Elementary Mathematics: Their History and Solution, Heinrich Dörrie. \$2.00
Mathematical Tables of Elementary and Some Higher Mathematical Functions, Herbert Bristol Dwight. \$2.00
Coordinate Geometry, Luther P. Eisenhart. \$1.65
The Theory and Operation of the Slide Rule, John P. Ellis. \$1.50
College Algebra, Henry Burchard Fine. \$2.75
Mathematical Fun, Games and Puzzles, Jack Frolichstein. \$2.00
A Treatise on Plane and Advanced Trigonometry, Ernest W. Hobson. \$2.25
Arithmetic Refresher for Practical Men, A. Albert Klaf. \$2.00
Calculus Refresher for Technical Men, A. Albert Klaf. \$2.00
Trigonometry Refresher for Technical Men, A. Albert Klaf. \$2.00
Famous Problems of Elementary Geometry, Felix Klein. \$1.00
An Introduction to Symbolic Logic, Susanne L. Langer. \$2.00

(continued on back flap)

DOVER BOOKS ON ELEMENTARY AND INTERMEDIATE MATHEMATICS

- Teach Yourself Algebra*, P. Abbott. Clothbound \$2.00
Teach Yourself Calculus, P. Abbott. Clothbound \$2.00
Teach Yourself Geometry, P. Abbott. Clothbound \$2.00
Teach Yourself Trigonometry, P. Abbott. Clothbound \$2.00
Handbook of Mathematical Functions: with Formulas, Graphs and Mathematical Tables, Milton Abramowitz and Irene A. Stegun. \$4.00
A Short Account of the History of Mathematics, W. W. Rouse Ball. \$2.25
Recreations in the Theory of Numbers: The Queen of Mathematics Entertains, Albert Beiler. \$2.00
Probabilities and Life, Emile Borel. \$1.00
Arithmetical Excursions: An Enrichment of Elementary Mathematics, Henry Bowers and Joan E. Bowers. \$1.65
Mathematical Tables and Formulas, Robert D. Carmichael and Edwin R. Smith. \$1.25
Textbook of Algebra, G. Chrystal. Two-volume set \$4.70
The Mathematics of Great Amateurs, Julian Lowell Coolidge. \$1.50
Elementary Statistics with Applications in Medicine and the Biological Sciences, Frederick E. Croxton. \$2.00
Statistical Adjustment of Data, W. Edwards Deming. \$2.00
Contemporary Geometry, André Delachet. \$1.00
100 Great Problems of Elementary Mathematics: Their History and Solution, Heinrich Dörrie. \$2.00
Mathematical Tables of Elementary and Some Higher Mathematical Functions, Herbert Bristol Dwight. \$2.00
Coordinate Geometry, Luther P. Eisenhart. \$1.65
The Theory and Operation of the Slide Rule, John P. Ellis. \$1.50
College Algebra, Henry Burchard Fine. \$2.75
Mathematical Fun, Games and Puzzles, Jack Frolichstein. \$2.00
A Treatise on Plane and Advanced Trigonometry, Ernest W. Hobson. \$2.25
Arithmetic Refresher for Practical Men, A. Albert Klaf. \$2.00
Calculus Refresher for Technical Men, A. Albert Klaf. \$2.00
Trigonometry Refresher for Technical Men, A. Albert Klaf. \$2.00
Famous Problems of Elementary Geometry, Felix Klein. \$1.00
An Introduction to Symbolic Logic, Susanne L. Langer. \$2.00

(continued on back flap)

Famous Problems of Elementary Geometry

THE DUPLICATION OF THE CUBE
THE TRISECTION OF AN ANGLE
THE QUADRATURE OF THE CIRCLE

by

Felix Klein



Translated by

Wooster Woodruff Beman

and

David Eugene Smith

Second Edition Revised and Enlarged with Notes by
RAYMOND CLARE ARCHIBALD

Dover Publications, Inc., New York



Library

IAS, Shimla

513 K 672 F




00039358

Published in Canada by General Publishing Company, Ltd., 30 Lesmill Road, Don Mills, Toronto, Ontario.

Published in the United Kingdom by Constable and Company, Ltd., 10 Orange Street, London WC 2.

This Dover edition, first published in 1956, is an unabridged and unaltered republication of the second revised edition of the authorized translation of *Vortrage über ausgewählte Fragen der Elementargeometrie*, as published by G. E. Stechert and Company in 1930.



513
K672 F

Manufactured in the United States of America
Dover Publications, Inc.
180 Varick Street
New York, N. Y. 10014

PREFACE.

THE more precise definitions and more rigorous methods of demonstration developed by modern mathematics are looked upon by the mass of gymnasium professors as abstruse and excessively abstract, and accordingly as of importance only for the small circle of specialists. With a view to counteracting this tendency it gave me pleasure to set forth last summer in a brief course of lectures before a larger audience than usual what modern science has to say regarding the possibility of elementary geometric constructions. Some time before, I had had occasion to present a sketch of these lectures in an Easter vacation course at Göttingen. The audience seemed to take great interest in them, and this impression has been confirmed by the experience of the summer semester. I venture therefore to present a short exposition of my lectures to the Association for the Advancement of the Teaching of Mathematics and the Natural Sciences, for the meeting to be held at Göttingen. This exposition has been prepared by Oberlehrer Tägert, of Ems, who attended the vacation course just mentioned. He also had at his disposal the lecture notes written out under my supervision by several of my summer semester students. I hope that this unpretending little book may contribute to promote the useful work of the association.

GÖTTINGEN, Easter, 1895.

F. KLEIN.

TRANSLATORS' PREFACE.

At the Göttingen meeting of the German Association for the Advancement of the Teaching of Mathematics and the Natural Sciences, Professor Felix Klein presented a discussion of the three famous geometric problems of antiquity, — the duplication of the cube, the trisection of an angle, and the quadrature of the circle, as viewed in the light of modern research.

This was done with the avowed purpose of bringing the study of mathematics in the university into closer touch with the work of the gymnasium. That Professor Klein is likely to succeed in this effort is shown by the favorable reception accorded his lectures by the association, the uniform commendation of the educational journals, and the fact that translations into French and Italian have already appeared.

The treatment of the subject is elementary, not even a knowledge of the differential and integral calculus being required. Among the questions answered are such as these: Under what circumstances is a geometric construction possible? By what means can it be effected? What are transcendental numbers? How can we prove that e and π are transcendental?

With the belief that an English presentation of so important a work would appeal to many unable to read the original,

Professor Klein's consent to a translation was sought and readily secured.

In its preparation the authors have also made free use of the French translation by Professor J. Griess, of Algiers, following its modifications where it seemed advisable.

They desire further to thank Professor Ziwet for assistance in improving the translation and in reading the proof-sheets.

W. W. BEMAN.

D. E. SMITH.

August, 1897.

EDITOR'S PREFACE.



Within three years of its publication thirty-five years ago Klein's little work was translated into English, French, Italian, and Russian¹. In the United States it filled a decided need for many years, and not a few teachers regretted that the work was allowed to go out of print. No other work supplied in such compact form just the information here found. Hence it seemed desirable to have a new edition with at least some of the slips of the first edition rectified, and with added notes illuminating the text.

The corrections and notes of the present edition are little more than revised extracts from my article in *The American Mathematical Monthly*², 1914. I am indebted to the Editors for courteously allowing the reproduction of this material.

R. C. A.

February, 1930.

¹ French translation by Griess, Paris, Nony, 1896; Italian by Giudice, Turin, Rosenberg e Sallier, 1896; Russian by Parfentiev and Sintsov, Kazan, 1898. This last translation seems to have been unknown to the editors of Klein's *Abhandlungen* (see v. 3, 1923, p. 28).

² Remarks on Klein's "Famous Problems of Elementary Geometry", v. 21, p. 247—259.

CONTENTS.



INTRODUCTION.

PRACTICAL AND THEORETICAL CONSTRUCTIONS	PAGE
STATEMENT OF THE PROBLEM IN ALGEBRAIC FORM	2
	3

PART I.

The Possibility of the Construction of Algebraic Expressions.

CHAPTER I. ALGEBRAIC EQUATIONS SOLVABLE BY SQUARE ROOTS.

1-4. Structure of the expression x to be constructed	5
5, 6. Normal form of x	6
7, 8. Conjugate values	7
9. The corresponding equation $F(x) = 0$	8
10. Other rational equations $f(x) = 0$	8
11, 12. The irreducible equation $\phi(x) = 0$	10
13, 14. The degree of the irreducible equation a power of 2	11

CHAPTER II. THE DELIAN PROBLEM AND THE TRISECTION OF THE ANGLE.

1. The impossibility of solving the Delian problem with straight edge and compasses	13
2. The general equation $x^3 = \lambda$	13
3. The impossibility of trisecting an angle with straight edge and compasses	14

CHAPTER III. THE DIVISION OF THE CIRCLE INTO EQUAL PARTS.

1. History of the problem	16
2-4. Gauss's prime numbers	17
5. The cyclotomic equation	19
6. Gauss's Lemma	19
7, 8. The irreducibility of the cyclotomic equation	21

CHAPTER IV. THE CONSTRUCTION OF THE REGULAR POLYGON OF 17 SIDES.

	PAGE
1. Algebraic statement of the problem	24
2-4. The periods formed from the roots	25
5, 6. The quadratic equations satisfied by the periods	27
7. Historical account of constructions with straight edge and compasses	32
8, 9. Von Staudt's construction of the regular polygon of 17 sides	34

CHAPTER V. GENERAL CONSIDERATIONS ON ALGEBRAIC CONSTRUCTIONS.

1. Paper folding	42
2. The conic sections	42
3. The Cissoid of Diocles	44
4. The Conchoid of Nicomedes	45
5. Mechanical devices	47

PART II.

Transcendental Numbers and the Quadrature of the Circle.

CHAPTER I. CANTOR'S DEMONSTRATION OF THE EXISTENCE OF TRANSCENDENTAL NUMBERS.

1. Definition of algebraic and of transcendental numbers	49
2. Arrangement of algebraic numbers according to <i>height</i>	50
3. Demonstration of the existence of transcendental numbers	53

CHAPTER II. HISTORICAL SURVEY OF THE ATTEMPTS AT THE COMPUTATION AND CONSTRUCTION OF π .

1. The empirical stage	56
2. The Greek mathematicians	56
3. Modern analysis from 1670 to 1770	58
4, 5. Revival of critical rigor since 1770	59

CHAPTER III. THE TRANSCENDENCE OF THE NUMBER e .

1. Outline of the demonstration	61
2. The symbol h_r and the function $\phi(x)$	62
3. Hermite's Theorem	65

CHAPTER IV. THE TRANSCENDENCE OF THE NUMBER π .

	PAGE
1. Outline of the demonstration	68
2. The function $\psi(x)$	70
3. Lindemann's Theorem	73
4. Lindemann's Corollary	74
5. The transcendence of π	76
6. The transcendence of $y = e^x$	77
7. The transcendence of $y = \sin^{-1}x$	77

CHAPTER V. THE INTEGRAPH AND THE GEOMETRIC CONSTRUCTION
OF π .

1. The impossibility of the quadrature of the circle with straight edge and compasses	78
2. Principle of the integraph	78
3. Geometric construction of π	79

NOTES	81
-----------------	----

INTRODUCTION.

THIS course of lectures is due to the desire on my part to bring the study of mathematics in the university into closer touch with the needs of the secondary schools. Still it is not intended for beginners, since the matters under discussion are treated from a higher standpoint than that of the schools. On the other hand, it presupposes but little preliminary work, only the elements of analysis being required, as, for example, in the development of the exponential function into a series.

We propose to treat of geometrical constructions, and our object will not be so much to find the solution suited to each case as to determine the *possibility* or *impossibility* of a solution.

Three problems, the object of much research in ancient times, will prove to be of special interest. They are

1. *The problem of the duplication of the cube* (also called the *Delian problem*).

2. *The trisection of an arbitrary angle.*

3. *The quadrature of the circle, i.e., the construction of π .*

In all these problems the ancients sought in vain for a solution with straight edge and compasses, and the celebrity of these problems is due chiefly to the fact that their solution seemed to demand the use of appliances of a higher order. In fact, we propose to show that a solution by the use of straight edge and compasses is impossible.

The impossibility of the solution of the third problem was demonstrated only very recently. That of the first and second is implicitly involved in the Galois theory as presented to-day in treatises on higher algebra. On the other hand, we find no explicit demonstration in elementary form unless it be in Petersen's text-books, works which are also noteworthy in other respects.

At the outset we must insist upon the difference between *practical* and *theoretical* constructions. For example, if we need a divided circle as a measuring instrument, we construct it simply on trial. Theoretically, in earlier times, it was possible (*i.e.*, by the use of straight edge and compasses) only to divide the circle into a number of parts represented by 2^n , 3, and 5, and their products. Gauss added other cases by showing the possibility of the division into parts where p is a prime number of the form $p = 2^m + 1$, and the impossibility for all other numbers. No practical advantage is derived from these results; *the significance of Gauss's developments is purely theoretical*. The same is true of all the discussions of the present course.

Our fundamental problem may be stated: *What geometrical constructions are, and what are not, theoretically possible?* To define sharply the meaning of the word "construction," we must designate the instruments which we propose to use in each case. We shall consider

1. Straight edge and compasses,
2. Compasses alone,
3. Straight edge alone,
4. Other instruments used in connection with straight edge and compasses.

The singular thing is that elementary geometry furnishes no answer to the question. We must fall back upon algebra and the higher analysis. The question then arises: How

shall we use the language of these sciences to express the employment of straight edge and compasses? This new method of attack is rendered necessary because elementary geometry possesses no general method, no *algorithm*, as do the last two sciences.

In analysis we have first *rational* operations: addition, subtraction, multiplication, and division. These operations can be directly effected geometrically upon two given segments by the aid of proportions, if, in the case of multiplication and division, we introduce an auxiliary unit-segment.

Further, there are *irrational* operations, subdivided into *algebraic* and *transcendental*. The simplest algebraic operations are the extraction of square and higher roots, and the solution of algebraic equations not solvable by radicals, such as those of the fifth and higher degrees. As we know how to construct \sqrt{ab} , rational operations in general, and irrational operations involving only square roots, can be constructed. On the other hand, every *individual* geometrical construction which can be reduced to the intersection of two straight lines, a straight line and a circle, or two circles, is equivalent to a rational operation or the extraction of a square root. In the higher irrational operations the construction is therefore impossible, *unless we can find a way of effecting it by the aid of square roots*. In all these constructions it is obvious that the number of operations must be limited.

We may therefore state the following fundamental theorem: *The necessary and sufficient condition that an analytic expression can be constructed with straight edge and compasses is that it can be derived from the known quantities by a finite number of rational operations and square roots.*

Accordingly, if we wish to show that a quantity cannot be constructed with straight edge and compasses, we must prove that the corresponding equation is not solvable by a finite number of square roots.

A fortiori the solution is impossible when the problem has *no* corresponding algebraic equation. An expression which satisfies no algebraic equation is called a transcendental number. This case occurs, as we shall show, with the number π .

PART I

THE POSSIBILITY OF THE CONSTRUCTION OF ALGEBRAIC EXPRESSIONS



CHAPTER I.

Algebraic Equations Solvable by Square Roots.

The following propositions taken from the theory of algebraic equations are probably known to the reader, yet to secure greater clearness of view we shall give brief demonstrations.

If x , the quantity to be constructed, depends only upon rational expressions and square roots, it is a root of an irreducible equation $\phi(x) = 0$, whose degree is always a power of 2.

1. To get a clear idea of the structure of the quantity x , suppose it, *e.g.*, of the form

$$x = \frac{\sqrt{a + \sqrt{c + ef}} + \sqrt{d + \sqrt{b}}}{\sqrt{a} + \sqrt{b}} + \frac{p + \sqrt{q}}{\sqrt{r}},$$

where $a, b, c, d, e, f, p, q, r$ are rational expressions.

2. The number of radicals one over another occurring in any term of x is called the *order of the term*; the preceding expression contains terms of orders 0, 1, 2.

3. Let μ designate the *maximum order*, so that no term can have more than μ radicals one over another.

PART I

THE POSSIBILITY OF THE CONSTRUCTION OF ALGEBRAIC EXPRESSIONS

4. In the example $x = \sqrt{2} + \sqrt{3} + \sqrt{6}$, we have three expressions of the first order, but as it may be written

$$x = \sqrt{2} + \sqrt{3} + \sqrt{2} \cdot \sqrt{3},$$

it really depends on only two distinct expressions.

We shall suppose that this reduction has been made in all the terms of x , so that among the n terms of order μ none can be expressed rationally as a function of any other terms of order μ or of lower order.

We shall make the same supposition regarding terms of the order $\mu - 1$ or of lower order, whether these occur explicitly or implicitly. This hypothesis is obviously a very natural one and of great importance in later discussions.

5. NORMAL FORM OF x .

If the expression x is a sum of terms with different denominators we may reduce them to the same denominator and thus obtain x as the quotient of two integral functions.

Suppose \sqrt{Q} one of the terms of x of order μ ; it can occur in x only explicitly, since μ is the maximum order. Since, further, the powers of \sqrt{Q} may be expressed as functions of \sqrt{Q} and Q , which is a term of lower order, we may put

$$x = \frac{a + b\sqrt{Q}}{c + d\sqrt{Q}},$$

where a, b, c, d contain no more than $n - 1$ terms of order μ , besides terms of lower order.

Multiplying both terms of the fraction by $c - d\sqrt{Q}$, \sqrt{Q} disappears from the denominator, and we may write

$$x = \frac{(ac - bdQ) + (bc - ad)\sqrt{Q}}{c^2 - d^2Q} = \alpha + \beta\sqrt{Q},$$

where α and β contain no more than $n - 1$ terms of order μ .

For a second term of order μ we have, in a similar manner, $x = \alpha_1 + \beta_1\sqrt{Q_1}$, etc.

The x may, therefore, be transformed so as to contain a term of given order μ only in its numerator and there only linearly.

We observe, however, that products of terms of order μ may occur, for α and β still depend upon $n - 1$ terms of order μ . We may, then, put

$$\alpha = \alpha_{11} + \alpha_{12} \sqrt{Q_1}, \quad \beta = \beta_{11} + \beta_{12} \sqrt{Q_1},$$

and hence

$$x = (\alpha_{11} + \alpha_{12} \sqrt{Q_1}) + (\beta_{11} + \beta_{12} \sqrt{Q_1}) \sqrt{Q}.$$

6. We proceed in a similar way with the different terms of order $\mu - 1$, which occur explicitly and in Q, Q_1 , etc., so that each of these quantities becomes an integral linear function of the term of order $\mu - 1$ under consideration. We then pass on to terms of lower order and finally obtain x , or rather its terms of different orders, under the form of rational integral linear functions of the individual radical expressions which occur explicitly. We then say that x is reduced to the *normal form*.

7. Let m be the total number of independent (4) square roots occurring in this normal form. Giving the double sign to these square roots and combining them in all possible ways, we obtain a system of 2^m values.

$$x_1, x_2, \dots, x_{2^m},$$

which we shall call *conjugate values*.

We must now investigate the equation admitting these conjugate values as roots.

8. These values are not necessarily all distinct; thus, if

$$\text{we have} \quad x = \sqrt{a + \sqrt{b}} + \sqrt{a - \sqrt{b}},$$

this expression is not changed when we change the sign of \sqrt{b} .

9. If x is an arbitrary quantity and we form the polynomial

$$F(x) = (x - x_1)(x - x_2) \dots (x - x_{2m}),$$

$F(x) = 0$ is clearly an equation having as roots these conjugate values. It is of degree $2m$, but may have equal roots (8).

The coefficients of the polynomial $F(x)$ arranged with respect to x are rational.

For let us change the sign of one of the square roots; this will permute two roots, say x_λ and $x_{\lambda'}$, since the roots of $F(x) = 0$ are precisely all the conjugate values. As these roots enter $F(x)$ only under the form of the product

$$(x - x_\lambda)(x - x_{\lambda'}),$$

we merely change the order of the factors of $F(x)$. Hence the polynomial is not changed.

$F(x)$ remains, then, invariable when we change the sign of any one of the square roots; it therefore contains only their squares; and hence $F(x)$ has only rational coefficients.

10. *When any one of the conjugate values satisfies a given equation with rational coefficients, $f(x) = 0$, the same is true of all the others.*

$f(x)$ is not necessarily equal to $F(x)$, and may admit other roots besides the x_i 's.

Let $x_1 = \alpha + \beta\sqrt{Q}$ be one of the conjugate values; \sqrt{Q} , a term of order μ ; α and β now depend only upon other terms of order μ and terms of lower order. There must, then, be a conjugate value

$$x_1' = \alpha - \beta\sqrt{Q}.$$

Let us now form the equation $f(x_1) = 0$. $f(x_1)$ may be put into the normal form with respect to \sqrt{Q} ,

$$f(x_1) = A + B\sqrt{Q};$$

this expression can equal zero only when A and B are simultaneously zero. Otherwise we should have

$$\sqrt{Q} = -\frac{A}{B};$$

i.e., \sqrt{Q} could be expressed rationally as a function of terms of order μ and of terms of lower order contained in A and B, which is contrary to the hypothesis of the independence of all the square roots (4).

But we evidently have

$$f(x_1') = A - B \sqrt{Q};$$

hence if $f(x_1) = 0$, so also $f(x_1') = 0$. Whence the following proposition :

If x_1 satisfies the equation $f(x) = 0$, the same is true of all the conjugate values derived from x_1 by changing the signs of the roots of order μ .

The proof for the other conjugate values is obtained in an analogous manner. Suppose, for example, as may be done without affecting the generality of the reasoning, that the expression x_1 depends on only two terms of order μ , \sqrt{Q} and $\sqrt{Q'}$. $f(x_1)$ may be reduced to the following normal form :

$$(a) \quad f(x_1) = p + q \sqrt{Q} + r \sqrt{Q'} + s \sqrt{Q} \cdot \sqrt{Q'} = 0.$$

If x_1 depended on more than two terms of order μ , we should only have to add to the preceding expression a greater number of terms of analogous structure.

Equation (a) is possible only when we have separately

$$(b) \quad p = 0, \quad q = 0, \quad r = 0, \quad s = 0.$$

Otherwise \sqrt{Q} and $\sqrt{Q'}$ would be connected by a rational relation, contrary to our hypothesis.

Let now \sqrt{R} , $\sqrt{R'}$, ... be the terms of order $\mu - 1$ on which x_1 depends; they occur in p, q, r, s; then can the quantities p, q, r, s, in which they occur, be reduced to the

normal form with respect to \sqrt{R} and $\sqrt{R'}$; and if, for the sake of simplicity, we take only two quantities, \sqrt{R} and $\sqrt{R'}$, we have

$$(c) \quad p = \kappa_1 + \lambda_1 \sqrt{R} + \mu_1 \sqrt{R'} + \nu_1 \sqrt{R} \cdot \sqrt{R'} = 0,$$

and three analogous equations for q, r, s .

The hypothesis, already used several times, of the independence of the roots, furnishes the equations

$$(d) \quad \kappa = 0, \quad \lambda = 0, \quad \mu = 0, \quad \nu = 0.$$

Hence equations (c) and consequently $f(x) = 0$ are satisfied when for x_1 we substitute the conjugate values deduced by changing the signs of \sqrt{R} and $\sqrt{R'}$.

Therefore *the equation $f(x) = 0$ is also satisfied by all the conjugate values deduced from x_1 by changing the signs of the roots of order $\mu - 1$.*

The same reasoning is applicable to the terms of order $\mu - 2, \mu - 3, \dots$ and our theorem is completely proved.

11. We have so far considered two equations

$$F(x) = 0 \quad \text{and} \quad f(x) = 0.$$

Both have rational coefficients and contain the x_1 's as roots. $F(x)$ is of degree 2^m and may have multiple roots; $f(x)$ may have other roots besides the x_1 's. We now introduce a third equation, $\phi(x) = 0$, defined as the equation of lowest degree, with rational coefficients, admitting the root x_1 and consequently all the x_1 's (10).

12. PROPERTIES OF THE EQUATION $\phi(x) = 0$.

I. $\phi(x) = 0$ is an irreducible equation, i.e., $\phi(x)$ cannot be resolved into two rational polynomial factors. This irreducibility is due to the hypothesis that $\phi(x) = 0$ is the rational equation of lowest degree satisfied by the x_1 's.

For if we had

$$\phi(x) = \psi(x) \chi(x),$$

then $\phi(x_1) = 0$ would require either $\psi(x_1) = 0$, or $\chi(x_1) = 0$, or both. But since these equations are satisfied by all the conjugate values (10), $\phi(x) = 0$ would not then be the equation of lowest degree satisfied by the x_1 's.

II. $\phi(x) = 0$ has no multiple roots. Otherwise $\phi(x)$ could be decomposed into rational factors by the well-known methods of Algebra, and $\phi(x) = 0$ would not be irreducible.

III. $\phi(x) = 0$ has no other roots than the x_1 's. Otherwise $F(x)$ and $\phi(x)$ would admit a highest common divisor, which could be determined rationally. We could then decompose $\phi(x)$ into rational factors, and $\phi(x)$ would not be irreducible.

IV. Let M be the number of x_1 's which have distinct values, and let

$$x_1, x_2, \dots, x_M$$

be these quantities. We shall then have

$$\phi(x) = C(x - x_1)(x - x_2) \dots (x - x_M).$$

For $\phi(x) = 0$ is satisfied by the quantities x_1 and it has no multiple roots. The polynomial $\phi(x)$ is then determined save for a constant factor whose value has no effect upon $\phi(x) = 0$

V. $\phi(x) = 0$ is the only irreducible equation with rational coefficients satisfied by the x_1 's. For if $f(x) = 0$ were another rational irreducible equation satisfied by x_1 and consequently by the x_1 's, $f(x)$ would be divisible by $\phi(x)$ and therefore would not be irreducible.

By reason of the five properties of $\phi(x) = 0$ thus established, we may designate this equation, in short, as *the* irreducible equation satisfied by the x_1 's.

13. Let us now compare $F(x)$ and $\phi(x)$. These two polynomials have the x_1 's as their only roots, and $\phi(x)$ has no multiple roots. $F(x)$ is, then, divisible by $\phi(x)$; that is,

$$F(x) = F_1(x) \phi(x).$$

$F_1(x)$ necessarily has rational coefficients, since it is the quotient obtained by dividing $F(x)$ by $\phi(x)$. If $F_1(x)$ is not a constant it admits roots belonging to $F(x)$; and admitting one it admits all the x_i 's (10). Hence $F_1(x)$ is also divisible by $\phi(x)$, and

$$F_1(x) = F_2(x) \phi(x).$$

If $F_2(x)$ is not a constant the same reasoning still holds, the degree of the quotient being lowered by each operation. Hence at the end of a limited number of divisions we reach an equation of the form

$$F_{\nu-1}(x) = C_1 \cdot \phi(x),$$

and for $F(x)$,

$$F(x) = C_1 \cdot [\phi(x)]^\nu.$$

The polynomial $F(x)$ is then a power of the polynomial of minimum degree $\phi(x)$, except for a constant factor.

14. We can now determine the degree M of $\phi(x)$. $F(x)$ is of degree 2^m ; further, it is the ν th power of $\phi(x)$. Hence

$$2^m = \nu \cdot M.$$

Therefore M is also a power of 2 and we obtain the following theorem :

The degree of the irreducible equation satisfied by an expression composed of square roots only is always a power of 2

15. Since, on the other hand, there is only one irreducible equation satisfied by all the x_i 's (12, V.), we have the converse theorem :

If an irreducible equation is not of degree 2^h , it cannot be solved by square roots.

CHAPTER II.

The Delian Problem and the Trisection of the Angle.

1. Let us now apply the general theorem of the preceding chapter to the *Delian problem*, i.e., to the problem of the *duplication of the cube*. The equation of the problem is manifestly

$$x^3 = 2.$$

This is irreducible, since otherwise $\sqrt[3]{2}$ would have a rational value. For an equation of the third degree which is reducible must have a rational linear factor. Further, the degree of the equation is not of the form 2^h ; hence it cannot be solved by means of square roots, and the geometric construction with straight edge and compasses is impossible.

2. Next let us consider the more general equation

$$x^3 = \lambda,$$

λ designating a parameter which may be a complex quantity of the form $a + ib$. This equation furnishes us the analytical expressions for the geometrical problems of the multiplication of the cube and the trisection of an arbitrary angle. The question arises whether this equation is reducible, i.e., whether one of its roots can be expressed as a rational function of λ . It should be remarked that the irreducibility of an expression always depends upon the values of the quantities supposed to be known. In the case $x^3 = 2$, we were dealing with numerical quantities, and the question was whether $\sqrt[3]{2}$ could have a rational numerical value. In the equation $x^3 = \lambda$ we ask whether a root can be represented by a rational function of λ . In the first case, the so-called

domain of rationality comprehends the totality of rational numbers; in the second, it is made up of the rational functions of a parameter. If no limitation is placed upon this parameter we see at once that no expression of the form $\frac{\phi(\lambda)}{\psi(\lambda)}$, in which $\phi(\lambda)$ and $\psi(\lambda)$ are polynomials, can satisfy our equation. Under our hypothesis the equation is therefore irreducible, and since its degree is not of the form 2^h , it cannot be solved by square roots.

8. Let us now restrict the variability of λ . Assume




FIG. 1.

$$\lambda = r(\cos \phi + i \sin \phi);$$

whence $\sqrt[3]{\lambda} = \sqrt[3]{r} \sqrt[3]{\cos \phi + i \sin \phi}.$

Our problem resolves itself into two, to extract the cube root of a real number and also that of a complex number of the form $\cos \phi + i \sin \phi$, both numbers being regarded as arbitrary. We shall treat these separately.

I. The roots of the equation $x^3 = r$ are

$$\sqrt[3]{r}, \epsilon \sqrt[3]{r}, \epsilon^2 \sqrt[3]{r},$$

representing by ϵ and ϵ^2 the complex cube roots of unity

$$\epsilon = \frac{-1 + i\sqrt{3}}{2}, \quad \epsilon^2 = \frac{-1 - i\sqrt{3}}{2}.$$

Taking for the domain of rationality the totality of rational functions of r , we know by the previous reasoning that the equation $x^3 = r$ is irreducible. Hence the problem of the multiplication of the cube does not admit, in general, of a construction by means of straight edge and compasses.

II. The roots of the equation

$$x^3 = \cos \phi + i \sin \phi$$

are, by De Moivre's formula,

$$x_1 = \cos \frac{\phi}{3} + i \sin \frac{\phi}{3},$$

$$x_2 = \cos \frac{\phi + 2\pi}{3} + i \sin \frac{\phi + 2\pi}{3},$$

$$x_3 = \cos \frac{\phi + 4\pi}{3} + i \sin \frac{\phi + 4\pi}{3}.$$

These roots are represented geometrically by the vertices of an equilateral triangle inscribed in the circle with radius unity and center at the origin. The figure shows that to the root x_1 corresponds the argument $\frac{\phi}{3}$. Hence the equation

$$x^3 = \cos \phi + i \sin \phi$$

is the analytic expression of the problem of the trisection of the angle.




FIG. 2.

If this equation were reducible, one, at least, of its roots could be represented as a rational function of $\cos \phi$ and $\sin \phi$, its value remaining unchanged on substituting $\phi + 2\pi$ for ϕ . But if we effect this change by a continuous variation of the angle ϕ , we see that the roots x_1, x_2, x_3 undergo a cyclic permutation. Hence no root can be represented as a rational function of $\cos \phi$ and $\sin \phi$. The equation under consideration is irreducible and therefore cannot be solved by the aid of a finite number of square roots. Hence the trisection of the angle cannot be effected with straight edge and compasses.

This demonstration and the general theorem evidently hold good only when ϕ is an arbitrary angle; but for certain special values of ϕ the construction may prove to be possible, e.g., when $\phi = \frac{\pi}{2}$.

CHAPTER III.

The Division of the Circle into Equal Parts.

1. The problem of dividing a given circle into n equal parts has come down from antiquity ; for a long time we have known the possibility of solving it when $n = 2^h, 3, 5$, or the product of any two or three of these numbers. In his *Disquisitiones Arithmeticae*, Gauss extended this series of numbers by showing that the division is possible for every prime number of the form $p = 2^{2^h} + 1$ but impossible for all other prime numbers and their powers. If in $p = 2^{2^h} + 1$ we make $\mu = 0$ and 1, we get $p = 3$ and 5, cases already known to the ancients. For $\mu = 2$ we get $p = 2^{2^2} + 1 = 17$, a case completely discussed by Gauss.

For $\mu = 3$ we get $p = 2^{2^3} + 1 = 257$, likewise a prime number. The regular polygon of 257 sides can be constructed. Similarly for $\mu = 4$, since $2^{2^4} + 1 = 65537$ is a prime number. $\mu = 5, \mu = 6, \mu = 7, \mu = 8, \mu = 9, \mu = 11, \mu = 12, \mu = 15, \mu = 18, \mu = 23, \mu = 36, \mu = 38, \mu = 73$ give no prime numbers. The proof that the large numbers corresponding to $\mu = 5, 6, \dots, 73$ are not prime has required a large expenditure of labor and ingenuity. It is, therefore, quite possible that $\mu = 4$ is the last number for which a solution can be effected.

Upon the regular polygon of 257 sides Richelot published an extended investigation in Crelle's *Journal*, IX, 1832, pp. 1-26, 146-161, 209-230, 337-356. The title of the memoir is : *De resolutione algebraica aequationis $x^{257} = 1$, sive de divisione circuli per bisectionem anguli septies repetitam in partes 257 inter se aequales commentatio coronata.*

To the regular polygon of 65537 sides Professor Hermes of Lingen devoted ten years of his life, examining with care all the roots furnished by Gauss's method. His MSS. are preserved in the collection of the mathematical seminary in Göttingen. (Compare a communication of Professor Hermes in No. 3 of the *Göttinger Nachrichten* for 1894.)

2. We may restrict the problem of the division of the circle into n equal parts to the cases where n is a prime number p or a power p^a of such a number. For if n is a composite number and if μ and ν are factors of n , prime to each other, we can always find integers a and b , positive or negative, such that

$$1 = a\mu + b\nu;$$

whence

$$\frac{1}{\mu\nu} = \frac{a}{\nu} + \frac{b}{\mu}.$$

To divide the circle into $\mu\nu = n$ equal parts it is sufficient to know how to divide it into μ and ν equal parts respectively. Thus, for $n = 15$, we have

$$\frac{1}{15} = \frac{2}{3} - \frac{3}{5}.$$

3. As will appear, the division into p equal parts (p being a prime number) is possible only when p is of the form $p = 2^h + 1$. We shall next show that a prime number can be of this form only when $h = 2^k$. For this we shall make use of Fermat's Theorem :

If p is a prime number and a an integer not divisible by p , these numbers satisfy the congruence

$$a^{p-1} \equiv +1 \pmod{p}.$$

$p - 1$ is not necessarily the lowest exponent which, for a given value of a , satisfies the congruence. If s is the lowest exponent it may be shown that s is a divisor of $p - 1$. In particular, if $s = p - 1$ we say that a is a *primitive root* of p ,

and notice that for every prime number p there is a primitive root. We shall make use of this notion further on.

Suppose, then, p a prime number such that

$$(1) \quad p = 2^h + 1,$$

and s the least integer satisfying

$$(2) \quad 2^s \equiv +1 \pmod{p}.$$

From (1) $2^h < p$; from (2) $2^s > p$.

$$\therefore s > h.$$

(1) shows that h is the least integer satisfying the congruence

$$(3) \quad 2^h \equiv -1 \pmod{p}.$$

From (2) and (3), by division,

$$2^{s-h} \equiv -1 \pmod{p}.$$

$$\therefore (4) \quad s - h \not\leq h, \quad s \not\leq 2h.$$

From (3), by squaring,

$$2^{2h} \equiv 1 \pmod{p}.$$

Comparing with (2) and observing that s is the least exponent satisfying congruences of the form

$$2^x \equiv 1 \pmod{p},$$

we have

$$(5) \quad s \not\geq 2h.$$

$$\therefore s = 2h.$$

We have observed that s is a divisor of $p - 1 = 2^h$; the same is true of h , which is, therefore, a power of 2. Hence prime numbers of the form $2^h + 1$ are necessarily of the form $2^{2^r} + 1$.

4. This conclusion may be established otherwise. Suppose that h is divisible by an odd number, so that

$$h = h'(2r + 1);$$

then, by reason of the formula

$$x^{2n+1} + 1 = (x + 1)(x^{2n} - x^{2n-1} + \dots - x + 1),$$

$p = 2^{h(2n+1)} + 1$ is divisible by $2^h + 1$, and hence is not a prime number.

5. We now reach our fundamental proposition :

p being a prime number, the division of the circle into p equal parts by the straight edge and compasses is impossible unless p is of the form

$$p = 2^h + 1 = 2^{2^u} + 1.$$

Let us trace in the z -plane ($z = x + iy$) a circle of radius 1. To divide this circle into n equal parts, beginning at $z = 1$, is the same as to solve the equation

$$z^n - 1 = 0.$$

This equation admits the root $z = 1$; let us suppress this root by dividing by $z - 1$, which is the same geometrically as to disregard the initial point of the division. We thus obtain the equation

$$z^{n-1} + z^{n-2} + \dots + z + 1 = 0,$$

which may be called the *cyclotomic equation*. As noticed above, we may confine our attention to the cases where n is a prime number or a power of a prime number. We shall first investigate the case when $n = p$. The essential point of the proof is to show *that the above equation is irreducible*. For since, as we have seen, irreducible equations can only be solved by means of square roots in finite number when their degree is a power of 2, a division into p parts is always impossible when $p - 1$ is not equal to a power of 2, *i.e.* when

$$p \neq 2^h + 1 \neq 2^{2^u} + 1.$$

Thus we see why Gauss's prime numbers occupy such an exceptional position.

6. At this point we introduce a lemma known as *Gauss's Lemma*. If

$$F(z) = z^m + Az^{m-1} + Bz^{m-2} + \dots + Lz + M,$$

where A, B, \dots are integers, and $F(z)$ can be resolved into two rational factors $f(z)$ and $\phi(z)$, so that

$$F(z) = f(z) \cdot \phi(z) = (z^{m'} + \alpha_1 z^{m'-1} + \alpha_2 z^{m'-2} + \dots) \\ (z^{m''} + \beta_1 z^{m''-1} + \beta_2 z^{m''-2} + \dots),$$

then must the α 's and β 's also be integers. In other words :

If an integral expression can be resolved into rational factors these factors must be integral expressions.

Let us suppose the α 's and β 's to be fractional. In each factor reduce all the coefficients to the least common denominator. Let a_0 and b_0 be these common denominators. Finally multiply both members of our equation by $a_0 b_0$. It takes the form

$$a_0 b_0 F(z) = f_1(z) \phi_1(z) = (a_0 z^{m'} + a_1 z^{m'-1} + \dots) \\ (b_0 z^{m''} + b_1 z^{m''-1} + \dots).$$

The a 's are integral and prime to one another, as also the b 's, since a_0 and b_0 are the least common denominators.

Suppose a_0 and b_0 different from unity and let q be a prime divisor of $a_0 b_0$. Further, let a_1 be the first coefficient of $f_1(z)$ and b_k the first coefficient of $\phi_1(z)$ not divisible by q . Let us develop the product $f_1(z) \phi_1(z)$ and consider the coefficient of $z^{m'+m''-1-k}$. It will be

$$a_1 b_k + a_{1-1} b_{k+1} + a_{1-2} b_{k+2} + \dots + a_{1+1} b_{k-1} + a_{1+2} b_{k-2} + \dots$$

According to our hypotheses, all the terms after the first are divisible by q , but the first is not. Hence this coefficient is not divisible by q . Now the coefficient of $z^{m'+m''-1-k}$ in the first member is divisible by $a_0 b_0$, i.e., by q . Hence if the identity is true it is impossible for a coefficient not divisible by q to occur in each polynomial. The coefficients of one at least of the polynomials are then all divisible by q . Here is another absurdity, since we have seen that all the coefficients are

prime to one another. Hence we cannot suppose a_0 and b_0 different from 1, and consequently the α 's and β 's are integral.

7. In order to show that the cyclotomic equation is irreducible, it is sufficient to show by Gauss's Lemma that the first member cannot be resolved into factors with integral coefficients. To this end we shall employ the simple method due to Eisenstein, in Crelle's *Journal*, XXXIX, p. 167, which depends upon the substitution

$$z = x + 1.$$

We obtain

$$f(z) = \frac{z^p - 1}{z - 1} = \frac{(x + 1)^p - 1}{x} = x^{p-1} + px^{p-2} + \frac{p(p-1)}{1 \cdot 2} x^{p-3} \\ + \dots + \frac{p(p-1)}{1 \cdot 2} x + p = 0.$$

All the coefficients of the expanded member except the first are divisible by p ; the last coefficient is always p itself, by hypothesis a prime number. An expression of this class is always irreducible.

For if this were not the case we should have

$$f(x + 1) = (x^m + a_1 x^{m-1} + \dots + a_{m-1} x + a_m) \\ (x^{m'} + b_1 x^{m'-1} + \dots + b_{m'-1} x + b_{m'}),$$


where the a 's and b 's are integers.

Since the term of zero degree in the above expression of $f(z)$ is p , we have $a_m b_{m'} = p$. p being prime, one of the factors of $a_m b_{m'}$ must be unity. Suppose, then,

$$a_m = \pm p, \quad b_m = \pm 1.$$

Equating the coefficients of the terms in x , we have

$$\frac{p(p-1)}{2} = a_{m-1} b_{m'} + a_m b_{m'-1}.$$



The first member and the second term of the second being divisible by p , $a_{m-1}b_m$ must be so also. Since $b_m = \pm 1$, a_{m-1} is divisible by p . Equating the coefficients of the terms in x^2 we may show that a_{m-2} is divisible by p . Similarly we show that all of the remaining coefficients of the factor $x^m + a_1x^{m-1} + \dots + a_{m-1}x + a_m$ are divisible by p . But this cannot be true of the coefficient of x^m , which is 1. The assumed equality is impossible and hence the cyclotomic equation is irreducible when p is a prime.

8. We now consider the case where n is a power of a prime number, say $n = p^a$. We propose to show that when $p > 2$ the division of the circle into p^2 equal parts is impossible. The general problem will then be solved, since the division into p^a equal parts evidently includes the division into p^2 equal parts.

The cyclotomic equation is now

$$\frac{z^{p^2} - 1}{z - 1} = 0.$$

It admits as roots extraneous to the problem those which come from the division into p equal parts, i.e., the roots of the equation

$$\frac{z^p - 1}{z - 1} = 0.$$

Suppressing these roots by division we obtain

$$f(z) = \frac{z^{p^2} - 1}{z^p - 1} = 0$$

as the cyclotomic equation. This may be written

$$z^{p(p-1)} + z^{p(p-2)} + \dots + z^p + 1 = 0.$$

Transforming by the substitution

$$z = x + 1,$$

we have

$$(x + 1)^{p(p-1)} + (x + 1)^{p(p-2)} + \dots + (x + 1)^p + 1 = 0.$$

The number of terms being p , the term independent of x after development will be equal to p , and the sum will take the form

$$x^{p(p-1)} + p \cdot \chi(x),$$

where $\chi(x)$ is a polynomial with integral coefficients whose constant term is 1. We have just shown that such an expression is always irreducible. Consequently *the new cyclotomic equation is also irreducible.*

The degree of this equation is $p(p-1)$. On the other hand an irreducible equation is solvable by square roots only when its degree is a power of 2. Hence a circle is divisible into p^2 equal parts only when $p=2$, p being assumed to be a prime.

The same is true, as already noted, for the division into p^a equal parts when $a > 2$.

CHAPTER IV.

The Construction of the Regular Polygon of 17 Sides.

1. We have just seen that the division of the circle into equal parts by the straight edge and compasses is possible only for the prime numbers studied by Gauss. It will now be of interest to learn how the construction can actually be effected.

The purpose of this chapter, then, will be to show in an elementary way how to inscribe in the circle the regular polygon of 17 sides.

Since we possess as yet no method of construction based upon considerations purely geometrical, we must follow the path indicated by our general discussions. We consider, first of all, the roots of the cyclotomic equation

$$x^{16} + x^{15} + \dots + x^2 + x + 1 = 0,$$

and construct geometrically the expression, formed of square roots, deduced from it.

We know that the roots can be put into the transcendental form

$$\epsilon_{\kappa} = \cos \frac{2\kappa\pi}{17} + i \sin \frac{2\kappa\pi}{17} \quad (\kappa = 1, 2, \dots, 16);$$

and if

$$\epsilon_1 = \cos \frac{2\pi}{17} + i \sin \frac{2\pi}{17},$$

that

$$\epsilon_{\kappa} = \epsilon_1^{\kappa}.$$

Geometrically, these roots are represented in the complex plane by the vertices, different from 1, of the regular polygon of 17 sides inscribed in a circle of radius 1, having the origin

as center. The selection of ϵ_1 is arbitrary, but for the construction it is essential to indicate some ϵ as the point of departure. Having fixed upon ϵ_1 , the angle corresponding to ϵ_κ is κ times the angle corresponding to ϵ_1 , which completely determines ϵ_κ .

2. The fundamental idea of the solution is the following : *Forming a primitive root to the modulus 17 we may arrange the 16 roots of the equation in a cycle in a determinate order.*

As already stated, a number a is said to be a primitive root to the modulus 17 when the congruence

$$a^s \equiv +1 \pmod{17}$$

has for least solution $s = 17 - 1 = 16$. The number 3 possesses this property; for we have

$$\left. \begin{array}{llll} 3^1 \equiv 3 & 3^5 \equiv 5 & 3^9 \equiv 14 & 3^{13} \equiv 12 \\ 3^2 \equiv 9 & 3^6 \equiv 15 & 3^{10} \equiv 8 & 3^{14} \equiv 2 \\ 3^3 \equiv 10 & 3^7 \equiv 11 & 3^{11} \equiv 7 & 3^{15} \equiv 6 \\ 3^4 \equiv 13 & 3^8 \equiv 16 & 3^{12} \equiv 4 & 3^{16} \equiv 1 \end{array} \right\} \pmod{17}.$$

Let us then arrange the roots ϵ_κ so that their subscripts are the preceding remainders in order

$$\epsilon_3, \epsilon_9, \epsilon_{10}, \epsilon_{13}, \epsilon_5, \epsilon_{15}, \epsilon_{11}, \epsilon_{16}, \epsilon_{14}, \epsilon_8, \epsilon_7, \epsilon_4, \epsilon_{12}, \epsilon_2, \epsilon_6, \epsilon_1.$$

Notice that if r is the remainder of $3^\kappa \pmod{17}$, we have

$$3^\kappa = 17q + r,$$

whence

$$\epsilon_r = \epsilon_1^r = \epsilon_1^{3^\kappa}$$

If r' is the next remainder, we have similarly

$$\epsilon_{r'} = \epsilon_1^{3^{\kappa+1}} = (\epsilon_1^{3^\kappa})^3 = (\epsilon_r)^3.$$

Hence in this series of roots each root is the cube of the preceding.

Gauss's method consists in decomposing this cycle into sums containing 8, 4, 2, 1 roots respectively, corresponding to the divisors of 16. Each of these sums is called a period.

The periods thus obtained may be calculated successively as roots of certain quadratic equations.

The process just outlined is only a particular case of that employed in the general case of the division into p equal parts. The $p - 1$ roots of the cyclotomic equation are cyclically arranged by means of a primitive root of p , and the periods may be calculated as roots of certain auxiliary equations. The degree of these last depends upon the prime factors of $p - 1$. They are not necessarily equations of the second degree.

The general case has, of course, been treated in detail by Gauss in his *Disquisitiones*, and also by Bachmann in his work, *Die Lehre von der Kreisteilung* (Leipzig, 1872).

3. In our case of the 16 roots the periods may be formed in the following manner: Form two periods of 8 roots by taking in the cycle, first, the roots of even order, then those of odd order. Designate these periods by x_1 and x_2 , and replace each root by its index. We may then write symbolically

$$\begin{aligned}x_1 &= 9 + 13 + 15 + 16 + 8 + 4 + 2 + 1, \\x_2 &= 3 + 10 + 5 + 11 + 14 + 7 + 12 + 6.\end{aligned}$$

Operating upon x_1 and x_2 in the same way, we form 4 periods of 4 terms:

$$\begin{aligned}y_1 &= 13 + 16 + 4 + 1, \\y_2 &= 9 + 15 + 8 + 2, \\y_3 &= 10 + 11 + 7 + 6, \\y_4 &= 3 + 5 + 14 + 12.\end{aligned}$$

Operating in the same way upon the y 's, we obtain 8 periods of 2 terms:

$$\begin{aligned}z_1 &= 16 + 1, & z_5 &= 11 + 6, \\z_2 &= 13 + 4, & z_6 &= 10 + 7, \\z_3 &= 15 + 2, & z_7 &= 5 + 12, \\z_4 &= 9 + 8, & z_8 &= 3 + 14.\end{aligned}$$

It now remains to show that *these periods can be calculated successively by the aid of square roots.*

4. It is readily seen that the sum of the remainders corresponding to the roots forming a period z is always equal to 17. These roots are then ϵ_r and ϵ_{17-r} ;

$$\begin{aligned}\epsilon_r &= \cos r \frac{2\pi}{17} + i \sin r \frac{2\pi}{17}, \\ \epsilon_1 = \epsilon_{17-r} &= \cos (17-r) \frac{2\pi}{17} + i \sin (17-r) \frac{2\pi}{17}, \\ &= \cos r \frac{2\pi}{17} - i \sin r \frac{2\pi}{17}.\end{aligned}$$

Hence

$$\epsilon_r + \epsilon_r = 2 \cos r \frac{2\pi}{17}.$$

Therefore all the periods z are real, and we readily obtain

$$\begin{aligned}z_1 &= 2 \cos \frac{2\pi}{17}, & z_5 &= 2 \cos 6 \frac{2\pi}{17}, \\ z_2 &= 2 \cos 4 \frac{2\pi}{17}, & z_6 &= 2 \cos 7 \frac{2\pi}{17}, \\ z_3 &= 2 \cos 2 \frac{2\pi}{17}, & z_7 &= 2 \cos 5 \frac{2\pi}{17}, \\ z_4 &= 2 \cos 8 \frac{2\pi}{17}, & z_8 &= 2 \cos 3 \frac{2\pi}{17}.\end{aligned}$$

Moreover, by definition,

$$\begin{aligned}x_1 &= z_1 + z_2 + z_3 + z_4, & x_2 &= z_5 + z_6 + z_7 + z_8, \\ y_1 &= z_1 + z_2, & y_2 &= z_3 + z_4, & y_3 &= z_5 + z_6, & y_4 &= z_7 + z_8.\end{aligned}$$

5. It will be necessary to determine the relative magnitude of the different periods. For this purpose we shall employ the following artifice: We divide the semicircle of unit radius into 17 equal parts and denote by S_1, S_2, \dots, S_{17} the distances

of the consecutive points of division A_1, A_2, \dots, A_{17} from the initial point of the semicircle, S_{17} being equal to the diameter, i.e., equal to 2. The angle $A_\kappa A_{17} O$ has the same measure as the half of the arc $A_\kappa O$, which equals $\frac{2\kappa\pi}{34}$. Hence




FIG. 3.

$$S_\kappa = 2 \sin \frac{\kappa\pi}{34} = 2 \cos \frac{(17 - \kappa)\pi}{34}.$$

That this may be identical with $2 \cos h \frac{2\pi}{17}$, we must have

$$\begin{aligned} 4h &= 17 - \kappa, \\ \kappa &= 17 - 4h. \end{aligned}$$

Giving to h the values 1, 2, 3, 4, 5, 6, 7, 8, we find for κ the values 13, 9, 5, 1, -3, -7, -11, -15. Hence

$$\begin{aligned} z_1 &= S_{13}, & z_6 &= -S_7, \\ z_2 &= S_9, & z_7 &= -S_{11}, \\ z_3 &= S_5, & z_8 &= -S_8, \\ z_4 &= -S_{15}, & z_5 &= S_6. \end{aligned}$$

The figure shows that S_κ increases with the subscript; hence the order of increasing magnitude of the periods z is

$$z_4, z_6, z_5, z_7, z_2, z_8, z_3, z_1.$$

Moreover, the chord $A_\kappa A_{\kappa+p}$ subtends p divisions of the semicircle and is equal to S_p ; the triangle $OA_\kappa A_{\kappa+p}$ shows that

$$S_{\kappa+p} < S_\kappa + S_p,$$

and *a fortiori*

$$S_{\kappa+p} < S_{\kappa+r} + S_{p+r}.$$

Calculating the differences two and two of the periods y , we easily find

$$\begin{aligned}
y_1 - y_2 &= S_{13} + S_1 - S_0 + S_{16} > 0, \\
y_1 - y_3 &= S_{18} + S_1 + S_7 + S_{11} > 0, \\
y_1 - y_4 &= S_{13} + S_1 + S_3 - S_6 > 0, \\
y_2 - y_3 &= S_0 - S_{16} + S_7 + S_{11} > 0, \\
y_2 - y_4 &= S_0 - S_{16} + S_3 - S_6 < 0, \\
y_3 - y_4 &= -S_7 - S_{11} + S_3 - S_6 < 0.
\end{aligned}$$

Hence

$$y_3 < y_2 < y_4 < y_1.$$

Finally we obtain in a similar way

$$x_2 < x_1.$$

6. We now propose to calculate $z_1 = 2 \cos \frac{2\pi}{17}$. After making this calculation and constructing z_1 , we can easily deduce the side of the regular polygon of 17 sides. In order to find the quadratic equation satisfied by the periods, we proceed to determine symmetric functions of the periods.

Associating z_1 with the period z_2 and thus forming the period y_1 , we have, first,

$$z_1 + z_2 = y_1.$$

Let us now determine $z_1 z_2$. We have

$$z_1 z_2 = (16 + 1)(13 + 4),$$

where the symbolic product κp represents

$$\epsilon_\kappa \cdot \epsilon_p = \epsilon_{\kappa+p}.$$

Hence it should be represented symbolically by $\kappa + p$, remembering to subtract 17 from $\kappa + p$ as often as possible. Thus,

$$z_1 z_2 = 12 + 3 + 14 + 5 = y_4.$$

Therefore z_1 and z_2 are the roots of the quadratic equation

$$(\xi) \quad z^2 - y_1 z + y_4 = 0,$$

whence, since $z_1 > z_2$,

$$z_1 = \frac{y_1 + \sqrt{y_1^2 - 4y_4}}{2}, \quad z_2 = \frac{y_1 - \sqrt{y_1^2 - 4y_4}}{2}.$$

We must now determine y_1 and y_4 . Associating y_1 with the period y_2 , thus forming the period x_1 , and y_3 with the period y_4 , thus forming the period x_2 , we have, first,

$$y_1 + y_2 = x_1.$$

Then,

$$y_1 y_2 = (13 + 16 + 4 + 1)(9 + 15 + 8 + 2).$$

Expanding symbolically, the second member becomes equal to the sum of all the roots; that is, to -1 . Therefore y_1 and y_2 are the roots of the equation

$$(\eta) \quad y^2 - x_1 y - 1 = 0,$$

whence, since $y_1 > y_2$,

$$y_1 = \frac{x_1 + \sqrt{x_1^2 + 4}}{2}, \quad y_2 = \frac{x_1 - \sqrt{x_1^2 + 4}}{2}$$

Similarly,

$$y_3 + y_4 = x_2$$

and

$$y_3 y_4 = -1.$$

Hence y_3 and y_4 are the roots of the equation

$$(\eta') \quad y^2 - x_2 y - 1 = 0;$$

whence, since $y_4 > y_3$,

$$y_4 = \frac{x_2 + \sqrt{x_2^2 + 4}}{2}, \quad y_3 = \frac{x_2 - \sqrt{x_2^2 + 4}}{2}$$

It now remains to determine x_1 and x_2 . Since $x_1 + x_2$ is equal to the sum of all the roots,

$$x_1 + x_2 = -1.$$

Further,

$$x_1 x_2 = (13 + 16 + 4 + 1 + 9 + 15 + 8 + 2) \\ (10 + 11 + 7 + 6 + 3 + 5 + 14 + 12).$$

Expanding symbolically, each root occurs 4 times, and thus

$$x_1 x_2 = -4.$$

Therefore x_1 and x_2 are the roots of the quadratic

$$(\xi) \quad x^2 + x - 4 = 0;$$

whence, since $x_1 > x_2$,

$$x_1 = \frac{-1 + \sqrt{17}}{2}, \quad x_2 = \frac{-1 - \sqrt{17}}{2}$$

Solving equations ξ, η, η', ζ in succession, z_1 is determined by a series of square roots.

Effecting the calculations, we see that z_1 depends upon the four square roots

$$\sqrt{17}, \sqrt{x_1^2 + 4}, \sqrt{x_2^2 + 4}, \sqrt{y_1^2 - 4y_4}.$$

If we wish to reduce z_1 to the normal form we must see whether any one of these square roots can be expressed rationally in terms of the others.

Now, from the roots of (η) ,

$$\sqrt{x_1^2 + 4} = y_1 - y_3,$$

$$\sqrt{x_2^2 + 4} = y_4 - y_3.$$

Expanding symbolically, we verify that

$$(y_1 - y_2)(y_4 - y_3) = 2(x_1 - x_2),*$$

$$\begin{aligned} * (y_1 - y_2)(y_4 - y_3) &= (13 + 16 + 4 + 1 - 9 - 15 - 8 - 2)(3 + 5 + 14 \\ &\quad + 12 - 10 - 11 - 7 - 6) \\ &= 16 + 1 + 10 + 8 - 6 - 7 - 3 - 2 \\ &\quad + 2 + 4 + 13 + 11 - 9 - 10 - 6 - 5 \\ &\quad + 7 + 9 + 1 + 16 - 14 - 15 - 11 - 10 \\ &\quad + 4 + 6 + 15 + 13 - 11 - 12 - 8 - 7 \\ &\quad - 12 - 14 - 6 - 4 + 2 + 3 + 16 + 15 \\ &\quad - 1 - 3 - 12 - 10 + 8 + 9 + 5 + 4 \\ &\quad - 11 - 13 - 5 - 3 + 1 + 2 + 15 + 14 \\ &\quad - 5 - 7 - 16 - 14 + 12 + 13 + 9 + 8 \\ &= 2(16 + 1 + 8 + 2 + 4 + 13 + 15 + 9 - 10 - 6 - 7 - 3 - 11 - 5 - 14 \\ &\quad - 12) \\ &= 2(x_1 - x_2). \end{aligned}$$

that is,

$$\sqrt{x_1^2 + 4} \sqrt{x_2^2 + 4} = 2 \sqrt{17}.$$

Hence $\sqrt{x_2^2 + 4}$ can be expressed rationally in terms of the other two square roots. This equation shows that if two of the three differences $y_1 - y_2$, $y_4 - y_3$, $x_1 - x_2$ are positive, the same is true of the third, which agrees with the results obtained directly.

Replacing now x_1 , y_1 , y_4 by their numerical values, we obtain in succession

$$x_1 = \frac{-1 + \sqrt{17}}{2},$$

$$y_1 = \frac{-1 + \sqrt{17} + \sqrt{34 - 2\sqrt{17}}}{4},$$

$$y_4 = \frac{-1 - \sqrt{17} + \sqrt{34 + 2\sqrt{17}}}{4},$$

$$z_1 = \frac{-1 + \sqrt{17} + \sqrt{34 - 2\sqrt{17}}}{8}$$

$$+ \frac{\sqrt{68 + 12\sqrt{17} - 16\sqrt{34 + 2\sqrt{17}} - 2(1 - \sqrt{17})\sqrt{34 - 2\sqrt{17}}}}{8}$$

The algebraic part of the solution of our problem is now completed. We have already remarked that there is no known construction of the regular polygon of 17 sides based upon purely geometric considerations. There remains, then, only the geometric translation of the individual algebraic steps.

7. We may be allowed to introduce here a brief historical account of geometric constructions with straight edge and compasses.

In the geometry of the ancients the straight edge and compasses were always used together; the difficulty lay merely in bringing together the different parts of the figure so as not to

draw any unnecessary lines. Whether the several steps in the construction were made with straight edge or with compasses was a matter of indifference.

On the contrary, in 1797, the Italian Mascheroni succeeded in effecting all these constructions with the compasses alone; he set forth his methods in his *Geometria del compasso*, and claimed that constructions with compasses were practically more exact than those with the straight edge. As he expressly stated, he wrote for mechanics, and therefore with a practical end in view. Mascheroni's original work is difficult to read, and we are under obligations to Hutt for furnishing a brief *résumé* in German, *Die Mascheroni'schen Constructionen* (Halle, 1880).

Soon after, the French, especially the disciples of Carnot, the author of the *Géométrie de position*, strove, on the other hand, to effect their constructions as far as possible with the straight edge. (See also Lambert, *Freie Perspective*, 1774.)

Here we may ask a question which algebra enables us to answer immediately: In what cases can the solution of an algebraic problem be constructed with the straight edge alone? The answer is not given with sufficient explicitness by the authors mentioned. We shall say:

With the straight edge alone we can construct all algebraic expressions whose form is rational.

With a similar view Brianchon published in 1818 a paper, *Les applications de la théorie des transversales*, in which he shows how his constructions can be effected in many cases with the straight edge alone. He likewise insists upon the practical value of his methods, which are especially adapted to field work in surveying.

Poncelet was the first, in his *Traité des propriétés projectives* (Vol. I, Nos. 351–357), to conceive the idea that it is sufficient to use a *single fixed circle* in connection with the straight lines

of the plane in order to construct all expressions depending upon square roots, the center of the fixed circle being given.

This thought was developed by Steiner in 1833 in a celebrated paper entitled *Die geometrischen Constructionen, ausgeführt mittels der geraden Linie und eines festen Kreises, als Lehrgegenstand für höhere Unterrichtsanstalten und zum Selbstunterricht.*

8. To construct the regular polygon of 17 sides we shall follow the method indicated by von Staudt (Crelle's *Journal*, Vol. XXIV, 1842), modified later by Schröter (Crelle's *Journal*, Vol. LXXV, 1872). The construction of the regular polygon of 17 sides is made in accordance with the methods indicated by Poncelet and Steiner, inasmuch as besides the straight edge but one fixed circle is used.*

First, we will show *how with the straight edge and one fixed circle we can solve every quadratic equation.*

At the extremities of a diameter of the fixed unit circle (Fig. 4) we draw two tangents, and select the lower as the

axis of X , and the diameter perpendicular to it as the axis of Y . Then the equation of the circle is

$$x^2 + y(y - 2) = 0.$$

Let

$$x^2 - px + q = 0$$




FIG. 4.

be any quadratic equation

with real roots x_1 and x_2 . Required to construct the roots x_1 and x_2 upon the axis of X .

Lay off upon the upper tangent from A to the right, a segment measured by $\frac{4}{p}$; upon the axis of X from O , a segment

* A Mascheroni construction of the regular polygon of 17 sides by L. Gérard is given in *Math. Annalen*, Vol. XLVIII, 1896, pp. 390-392.

measured by $\frac{q}{p}$; connect the extremities of these segments by the line 3 and project the intersections of this line with the circle from A, by the lines 1 and 2, upon the axis of X. The segments thus cut off upon the axis of X are measured by x_1 and x_2 .

Proof. Calling the intercepts upon the axis of X, x_1 and x_2 , we have the equation of the line 1,

$$2x + x_1(y - 2) = 0;$$

of the line 2.

$$2x + x_2(y - 2) = 0.$$

If we multiply the first members of these two equations we get

$$x^2 + \frac{x_1 + x_2}{2} x (y - 2) + \frac{x_1 x_2}{4} (y - 2)^2 = 0$$

as the equation of the line pair formed by 1 and 2. Subtracting from this the equation of the circle, we obtain

$$\frac{x_1 + x_2}{2} x (y - 2) + \frac{x_1 x_2}{4} (y - 2)^2 - y (y - 2) = 0$$

This is the equation of a conic passing through the four intersections of the lines 1 and 2 with the circle. From this equation we can remove the factor $y - 2$, corresponding to the tangent, and we have left

$$\frac{x_1 + x_2}{2} x + \frac{x_1 x_2}{4} (y - 2) - y = 0,$$

which is the equation of the line 3. If we now make $x_1 + x_2 = p$ and $x_1 x_2 = q$, we get

$$\frac{p}{2} x + \frac{q}{4} (y - 2) - y = 0,$$

and the transversal 3 cuts off from the line $y = 2$ the seg-

ment $\frac{4}{p}$, and from the line $y = 0$ the segment $\frac{q}{p}$. Thus the correctness of the construction is established.

9. In accordance with the method just explained, we shall now construct the roots of our four quadratic equations. They are (see pp. 29–31)

$$(\xi) \quad x^2 + x - 4 = 0, \text{ with roots } x_1 \text{ and } x_2; x_1 > x_2,$$

$$(\eta) \quad y^2 - x_1 y - 1 = 0, \text{ with roots } y_1 \text{ and } y_2; y_1 > y_2,$$

$$(\eta') \quad y^2 - x_2 y - 1 = 0, \text{ with roots } y_3 \text{ and } y_4; y_4 > y_3,$$

$$(\xi) \quad z^2 - y_1 z + y_4 = 0, \text{ with roots } z_1 \text{ and } z_2; z_1 > z_2.$$

These will furnish

$$z_1 = 2 \cos \frac{2\pi}{17},$$

whence it is easy to construct the polygon desired. We notice further that to construct z_1 it is sufficient to construct x_1, x_2, y_1, y_4 .

We then lay off the following segments: upon the upper tangent, $y = 2$,

$$-4, \frac{4}{x_1}, \frac{4}{x_2}, \frac{4}{y_1};$$

upon the axis of X ,

$$+4, -\frac{1}{x_1}, -\frac{1}{x_2}, \frac{y_4}{y_1}.$$

This may all be done in the following manner: The straight line connecting the point $+4$ upon the axis of X with the point -4 upon the tangent $y = 2$ cuts the circle in




FIG. 5.

two points, the projection of which from the point A (0, 2), the upper vertex of the circle, gives the two roots x_1, x_2 of the first quadratic equation as intercepts upon the axis of X.

To solve the second equation we have to lay off $\frac{4}{x_1}$ above and $-\frac{1}{x_1}$ below.

To determine the first point we connect x_1 upon the axis of X with A, the upper vertex, and from O, the lower vertex, draw another straight line through the intersection of this line with the circle. This cuts off upon the upper tangent the intercept $\frac{4}{x_1}$. This can easily be shown analytically.

The equation of the line from A to x_1 (Fig. 5),

$$2x + x_1y = 2x_1,$$

and that of the circle,

$$x^2 + y(y - 2) = 0,$$

give as the coördinates of their intersection

$$\frac{4x_1}{x_1^2 + 4}, \frac{2x_1^2}{x_1^2 + 4}.$$

The equation of the line from O through this point becomes

$$y = \frac{x_1}{2}x,$$

cutting off upon $y = 2$ the intercept $\frac{4}{x_1}$.

We reach the same conclusion still more simply by the use of some elementary notions of projective geometry. By our construction we have obviously associated with every point x of the lower range one, and only one, point of the upper, so that to the point $x = \infty$ corresponds the point $x' = 0$, and conversely. Since in such a correspondence there must exist a

linear relation, the abscissa x' of the upper point must satisfy the equation

$$x' = \frac{\text{const.}}{x}$$

Since $x' = 2$ when $x = 2$, as is obvious from the figure, the constant $= 4$.




FIG. 6.

To determine $-\frac{1}{x_1}$ upon the axis of X we connect the point -4 upon the upper with the point $+1$ upon the lower tangent (Fig. 6). The point thus determined upon the vertical diameter we connect with the point $\frac{4}{x_1}$ above. This line cuts off upon the axis of X the intercept $-\frac{1}{x_1}$. For the line from -4 to $+1$,

$$5y + 2x = 2,$$

intersects the vertical diameter in the point $(0, \frac{2}{5})$. Hence the equation of the line from $\frac{4}{x_1}$ to this point is

$$5y - 2x_1x = 2,$$

and its intersection with the lower tangent gives $-\frac{1}{x_1}$

The projection from A of the intersections of the line from $-\frac{1}{x_1}$ to $\frac{4}{x_1}$ with the circle determines upon the axis of X the two roots of the second quadratic equation, of which, as

already noted, we need only the greater, y_1 . This corresponds, as shown by the figure, to the projection of the upper intersection of our transversal with the circle.

Similarly, we obtain the roots of the third quadratic equation. Upon the upper tangent we project from O the intersection of the circle with the straight line which gave upon the axis of X the root $+x_2$. This immediately gives the intercept $\frac{4}{x_2}$, by reason of the correspondence just explained.




FIG. 7.

If we connect this point with the point where the vertical diameter intersects the line joining -4 above and $+1$ below, we cut off upon the axis of X the segment $-\frac{1}{x_2}$, as desired.

If we project that intersection of this transversal with the circle which lies in the positive quadrant from A upon the axis of X, we have constructed the required root y_4 of the third quadratic equation.

We have finally to determine the root z_1 of the fourth quadratic equation and for this purpose to lay off $\frac{4}{y_1}$ above and $\frac{y_4}{y_1}$ below. We solve the first problem in the usual way, by projecting the intersection of the circle with the line connecting A with $+y_1$ below, from O upon the upper tangent, thus obtaining $\frac{4}{y_1}$. For the other segment we connect the point $+4$ above with y_4 below, and then the point thus determined

upon the vertical diameter produced with $\frac{4}{y_1}$. This line cuts off upon the axis of X exactly the segment desired, $\frac{y_4}{y_1}$. For the line a (Fig. 8) has the equation

$$(y_4 - 4)y + 2x = 2y_4.$$




FIG. 8.

It cuts off upon the vertical diameter the segment $\frac{2y_4}{y_4 - 4}$. The equation of the line b is then

$$2y_1x + (y_4 - 4)y = 2y_4,$$

and its intersection with the axis of X has the abscissa $\frac{y_4}{y_1}$.

If we project the upper intersection of the line b with the circle from A upon the axis of X , we obtain $z_1 = 2 \cos \frac{2\pi}{17}$.

If we desire the simple cosine itself we have only to draw a diameter parallel to the axis of X , on which our last projecting ray cuts off directly $\cos \frac{2\pi}{17}$. A perpendicular erected at this point gives immediately the first and sixteenth vertices of the regular polygon of 17 sides.

The period z_1 was chosen arbitrarily; we might construct in the same way every other period of two terms and so find the remaining cosines. These constructions, made on separate figures so as to be followed more easily, have been combined in a single figure (Fig. 9), which gives the complete construction of the regular polygon of 17 sides.




FIG. 9.

CHAPTER V.

General Considerations on Algebraic Constructions.

1. We shall now lay aside the matter of construction with straight edge and compasses. Before quitting the subject we may mention a new and very simple method of effecting certain constructions, *paper folding*. Hermann Wiener* has shown how by paper folding we may obtain the network of the regular polyhedra. Singularly, about the same time a Hindu mathematician, Sundara Row, of Madras, published a little book, *Geometrical Exercises in Paper Folding* (Madras, Addison & Co., 1893), in which the same idea is considerably developed. The author shows how by paper folding we may construct by points such curves as the ellipse, cissoid, etc.

2. Let us now inquire how to solve geometrically problems whose analytic form is an equation of the third or of higher degree, and in particular, let us see how the ancients succeeded. The most natural method is by means of the conics, of which the ancients made much use. For example, they found that by means of these curves they were enabled to solve the problems of the duplication of the cube and the trisection of the angle. We shall in this place give only a general sketch of the process, making use of the language of modern mathematics for greater simplicity.

Let it be required, for instance, to solve graphically the cubic equation

$$x^3 + ax^2 + bx + c = 0,$$

or the biquadratic,

$$x^4 + ax^3 + bx^2 + cx + d = 0.$$

* See Dyck, *Katalog der Münchener mathematischen Ausstellung von 1893*, Nachtrag, p. 52.

Put $x^2 = y$; our equations become

$$xy + ay + bx + c = 0$$

and

$$y^2 + axy + by + cx + d = 0.$$

The roots of the equations proposed are thus the abscissas of the points of intersection of the two conics.

The equation

$$x^2 = y$$

represents a parabola with axis vertical. The second equation,

$$xy + ay + bx + c = 0,$$

represents an hyperbola whose asymptotes are parallel to the axes of reference (Fig. 10). One of the four points of inter-




FIG. 10.




FIG. 11.

section is at infinity upon the axis of Y , the other three at a finite distance, and their abscissas are the roots of the equation of the third degree.

In the second case the parabola is the same. The hyperbola (Fig. 11) has again one asymptote parallel to the axis of X while the other is no longer perpendicular to this axis. The curves now have four points of intersection at a finite distance.

The methods of the ancient mathematicians are given in detail in the elaborate work of M. Cantor, *Geschichte der Mathematik* (Leipzig, 1894, 2d ed.). Especially interesting is Zeuthen, *Die Kegelschnitte im Altertum* (Kopenhagen, 1886, in German edition). As a general compendium we may mention Baltzer, *Analytische Geometrie* (Leipzig, 1882).

3. Beside the conics, the ancients used for the solution of the above-mentioned problems, higher curves constructed for this very purpose. We shall mention here only the *Cissoid* and the *Conchoid*.




FIG. 12.

The *cissoid of Diocles* (c. 150 B.C.) may be constructed as follows (Fig. 12): To a circle draw a tangent (in the figure the vertical tangent on the right) and the diameter perpendicular to it. Draw lines from O, the vertex of the circle thus determined, to points upon the tangent, and lay off from O upon each the segment lying between its intersection with the circle and the tangent. The locus of points so determined is the *cissoid*.

To derive the equation, let r be the radius vector, θ the angle it makes with the axis of X. If we produce r to the tangent on the right, and call the diameter of the circle 1, the total segment equals $\frac{1}{\cos \theta}$. The portion cut off by the circle is $\cos \theta$. The difference of the two segments is r , and hence

$$r = \frac{1}{\cos \theta} - \cos \theta = \frac{\sin^2 \theta}{\cos \theta}.$$

By transformation of coördinates we obtain the Cartesian equation,

$$(x^2 + y^2)x - y^2 = 0.$$

The curve is of the third order, has a cusp at the origin, and is symmetric to the axis of X . The vertical tangent to the circle with which we began our construction is an asymptote. Finally the cissoid cuts the line at infinity in the circular points.

To show how to solve the Delian problem by the use of this curve, we write its equation in the following form :

$$\left(\frac{y}{x}\right)^3 = \frac{y}{1-x}.$$

We now construct the straight line,

$$\frac{y}{x} = \lambda.$$

This cuts off upon the tangent $x = 1$ the segment λ , and intersects the cissoid in a point for which

$$\frac{y}{1-x} = \lambda^3.$$

This is the equation of a straight line passing through the point $y = 0, x = 1$, and hence of the line joining this point to the point of the cissoid.

This line cuts off upon the axis of Y the intercept λ^3 .

We now see how $\sqrt[3]{2}$ may be constructed. Lay off upon the axis of Y the intercept 2, join this point to the point $x = 1, y = 0$, and through its intersection with the cissoid draw a line from the origin to the tangent $x = 1$. The intercept on this tangent equals $\sqrt[3]{2}$.

4. The *conchoid of Nicomedes* (c. 150 B.C.) is constructed as follows : Let O be a fixed point, a its distance from a fixed

line. If we pass a pencil of rays through O and lay off on each ray from its intersection with the fixed line in both directions a segment b , the locus of the points so determined is the *conchoid*. According as b is greater or less than a , the origin is a node or a conjugate point; for $b = a$ it is a cusp (Fig. 13).

Taking for axes of X and Y the perpendicular and parallel through O to the fixed line, we have

$$\frac{r}{x} = \frac{b}{x - a};$$

whence

$$(x^2 + y^2)(x - a)^2 - b^2x^2 = 0.$$

The conchoid is then of the fourth order, has a double point at the origin, and is composed of two branches having for common asymptote the line $x = a$. Further, the factor $(x^2 + y^2)$ shows that the curve passes through the circular points at infinity, a matter of immediate importance.

We may trisect any angle by means of this curve in the following manner: Let $\phi = \text{MOY}$ (Fig. 13) be the angle to be divided into three equal parts. On the side OM lay off $OM = b$, an arbitrary length. With M as a center and radius b describe a circle, and through M perpendicular to the axis of X with origin O draw a vertical line representing the asymptote of the conchoid to be constructed. Construct the




FIG. 13.

conchoid. Connect O with A, the intersection of the circle and the conchoid. Then is $\angle AOY$ one third of $\angle \phi$, as is easily seen from the figure.

Our previous investigations have shown us that the problem of the trisection of the angle is a problem of the third degree. It admits the three solutions

$$\frac{\phi}{3}, \quad \frac{\phi + 2\pi}{3}, \quad \frac{\phi + 4\pi}{3}.$$

Every algebraic construction which solves this problem by the aid of a curve of higher degree must obviously furnish all the solutions. Otherwise the equation of the problem would not be irreducible. These different solutions are shown in the figure. The circle and the conchoid intersect in eight points. Two of them coincide with the origin, two others with the circular points at infinity. None of these can give a solution of the problem. There remain, then, four points of intersection, so that we seem to have one too many. This is due to the fact that among the four points we necessarily find the point B such that $OMB = 2b$, a point which may be determined without the aid of the curve. There actually remain then only three points corresponding to the three roots furnished by the algebraic solution.

5. In all these constructions with the aid of higher algebraic curves, we must consider the practical execution. We need an instrument which shall trace the curve by a continuous movement, for a construction by points is simply a method of approximation. Several instruments of this sort have been constructed; some were known to the ancients. Nicomedes invented a simple device for tracing the conchoid. It is the oldest of the kind besides the straight edge and compasses. (Cantor, I, p. 302.) A list of instruments of more recent construction may be found in Dyck's Katalog, pp. 227-230. 340. and Nachtrag, pp. 42, 43.

PART II

TRANSCENDENTAL NUMBERS AND THE QUADRATURE OF THE CIRCLE

PART II

TRANSCENDENTAL NUMBERS AND THE QUADRATURE OF THE CIRCLE



CHAPTER I.

Cantor's Demonstration of the Existence of Transcendental Numbers.

1. Let us represent numbers as usual by points upon the axis of abscissas. If we restrict ourselves to *rational numbers* the corresponding points will fill the axis of abscissas densely throughout (*überall dicht*), i.e., in any interval no matter how small there is an infinite number of such points. Nevertheless, as the ancients had already discovered, the continuum of points upon the axis is not exhausted in this way; between the rational numbers come in the irrational numbers, and the question arises whether there are not distinctions to be made among the irrational numbers.

Let us define first what we mean by *algebraic numbers*. Every root of an algebraic equation

$$a_0\omega^n + a_1\omega^{n-1} + \dots + a_{n-1}\omega + a_n = 0$$

with integral coefficients is called an algebraic number. Of course we consider only the real roots. Rational numbers occur as a special case in equations of the form

$$a_0\omega + a_1 = 0.$$

PART II

TRANSCENDENTAL NUMBERS AND THE QUADRATURE OF THE CIRCLE

CHAPTER I.

Cantor's Demonstration of the Existence of Transcendental Numbers.

1. Let us represent numbers as usual by points upon the axis of abscissas. If we restrict ourselves to rational numbers the corresponding points will fill the axis of abscissas densely throughout (*überall dicht*), i.e., in any interval no matter how small there is an infinite number of such points. Nevertheless, as the ancients had already discovered, the continuum of points upon the axis is not exhausted in this way; between the rational numbers come in the irrational numbers, and the question arises whether there are not distinctions to be made among the irrational numbers.

Let us define first what we mean by *algebraic numbers*. Every root of an algebraic equation

$$a_0\omega^n + a_1\omega^{n-1} + \cdots + a_{n-1}\omega + a_n = 0$$

with integral coefficients is called an algebraic number. Of course we consider only the real roots. Rational numbers occur as a special case in equations of the form

$$a_0\omega + a_1 = 0.$$

We now ask the question: Does the totality of real algebraic numbers form a continuum, or a discrete series such that other numbers may be inserted in the intervals? These new numbers, the so-called *transcendental* numbers, would then be characterized by this property, that they cannot be roots of an algebraic equation with integral coefficients.

This question was answered first by Liouville (*Comptes rendus*, 1844, and Liouville's *Journal*, Vol. XVI, 1851), and in fact the existence of transcendental numbers was demonstrated by him. But his demonstration, which rests upon the theory of continued fractions, is rather complicated. The investigation is notably simplified by using the developments given by Georg Cantor in a memoir of fundamental importance, *Ueber eine Eigenschaft des Inbegriffes reeller algebraischer Zahlen* (Crelle's *Journal*, Vol. LXXVII, 1873). We shall give his demonstration, making use of a more simple notion which Cantor, under a different form, it is true, suggested at the meeting of naturalists in Halle, 1891.

2. The demonstration rests upon the fact that algebraic numbers form a *countable* mass, while transcendental numbers do not. By this Cantor means that the former can be arranged in a certain order so that each of them occupies a definite place, is numbered, so to speak. This proposition may be stated as follows:

The manifoldness of real algebraic numbers and the manifoldness of positive integers can be brought into a one-to-one correspondence.

We seem here to meet a contradiction. The positive integers form only a portion of the algebraic numbers; since each number of the first can be associated with one and one only of the second, the part would be equal to the whole. This objection rests upon a false analogy. The proposition that the part is always less than the whole is not true for

infinite masses. It is evident, for example, that we may establish a one-to-one correspondence between the aggregate of positive integers and the aggregate of positive even numbers, thus:

$$\begin{array}{ccccccc} 0 & 1 & 2 & 3 & \cdots & n & \cdots \\ 0 & 2 & 4 & 6 & \cdots & 2n & \cdots \end{array}$$

In dealing with infinite masses, the words *great* and *small* are inappropriate. As a substitute, Cantor has introduced the word *power* (*Mächtigkeit*), and says: *Two infinite masses have the same power when they can be brought into a one-to-one correspondence with each other.* The theorem which we have to prove then takes the following form: *The aggregate of real algebraic numbers has the same power as the aggregate of positive integers.*

We obtain the aggregate of real algebraic numbers by seeking the real roots of all algebraic equations of the form

$$a_0\omega^n + a_1\omega^{n-1} + \cdots + a_{n-1}\omega + a_n = 0;$$

all the a 's are supposed prime to one another, a_0 positive, and the equation irreducible. To arrange the numbers thus obtained in a definite order, we consider their *height* N as defined by

$$N = n - 1 + |a_0| + |a_1| + \cdots + |a_n|,$$

$|a_i|$ representing the absolute value of a_i , as usual. To a given number N corresponds a finite number of algebraic equations. For, N being given, the number n has certainly an upper limit, since N is equal to $n - 1$ increased by positive numbers; moreover, the difference $N - (n - 1)$ is a sum of positive numbers prime to one another, whose number is obviously finite.

N	n	$ a_0 $	$ a_1 $	$ a_2 $	$ a_3 $	$ a_4 $	EQUATION.	$\phi(N)$	ROOTS.
1	1	1	0				$x = 0$	1	0
	2	0	0	0			—		
2	1	2	0				—	2	— 1 + 1
		1	1				$x \pm 1 = 0$		
	2	1	0	0			—		
3	1	3	0				—	4	— 2 — $\frac{1}{2}$ + $\frac{1}{2}$ + 2
		2	1				$2x \pm 1 = 0$		
		1	2				$x \pm 2 = 0$		
	2	2	0	0			—		
		1	1	0			—		
		1	0	1			—		
	3	1	0	0	0		—		
4	1	4	0				—	12	— 3 — 1.61803 — 1.41421 — 0.70711 — 0.01803 — 0.33333 + 0.33333 + 0.61803 + 0.70711 + 1.41421 + 1.61803 + 3
		3	1				$3x \pm 1 = 0$		
		2	2				—		
		1	3				$x \pm 3 = 0$		
	2	3	0	0			—		
		2	1	0			—		
		2	0	1			$2x^2 - 1 = 0$		
		1	2	0			—		
		1	1	1			$x^2 \pm x - 1 = 0$		
		1	0	2			$x^2 - 2 = 0$		
		2	0	0	0		—		
		1	1	0	0		—		
	3	1	0	1	0		—		
		1	0	0	1		—		
		1	0	0	0	0	—		
		1	0	0	0	0	—		

Among these equations we must discard those that are reducible, which presents no theoretical difficulty. Since the number of equations corresponding to a given value of

N is limited, there corresponds to a determinate N only a finite mass of algebraic numbers. We shall designate this by $\phi(N)$. The table contains the values of $\phi(1)$, $\phi(2)$, $\phi(3)$, $\phi(4)$, and the corresponding algebraic numbers ω .

We arrange now the algebraic numbers according to their height, N, and the numbers corresponding to a single value of N in increasing magnitude. We thus obtain all the algebraic numbers, each in a determinate place. This is done in the last column of the accompanying table. It is, therefore, evident that algebraic numbers can be counted.

3. We now state the general proposition :

In any portion of the axis of abscissas, however small, there is an infinite number of points which certainly do not belong to a given countable mass.

Or, in other words :

The continuum of numerical values represented by a portion of the axis of abscissas, however small, has a greater power than any given countable mass.

This amounts to affirming the existence of transcendental numbers. It is sufficient to take as the countable mass the aggregate of algebraic numbers.

To demonstrate this theorem we prepare a table of algebraic numbers as before and write in it all the numbers in the form of decimal fractions. None of these will end in an infinite series of 9's. For the equality

$$1 = 0.999 \dots 9 \dots$$

shows that such a number is an exact decimal. If now we can construct a decimal fraction which is not found in our table and does not end in an infinite series of 9's it will certainly be a transcendental number. By means of a very simple process indicated by Georg Cantor we can find not only one but infinitely many transcendental numbers, even

when the domain in which the number is to lie is very small. Suppose, for example, that the first five decimals of the number are given. Cantor's process is as follows.

Take for 6th decimal a number different from 9 and from the 6th decimal of the *first algebraic* number, for 7th decimal a number different from 9 and from the 7th decimal of the *second algebraic* number, etc. In this way we obtain a decimal fraction which will not end in an infinite series of 9's and is certainly not contained in our table. The proposition is then demonstrated.

We see by this that (if the expression is allowable) there are far more transcendental numbers than algebraic. For when we determine the unknown decimals, avoiding the 9's, we have a choice among eight different numbers; we can thus form, so to speak, 8^{∞} transcendental numbers, even when the domain in which they are to lie is as small as we please.

CHAPTER II.

Historical Survey of the Attempts at the Computation and Construction of π .

In the next chapter we shall prove that the number π belongs to the class of transcendental numbers whose existence was shown in the preceding chapter. The proof was first given by Lindemann in 1882, and thus a problem was definitely settled which, so far as our knowledge goes, has occupied the attention of mathematicians for nearly 4000 years, the problem of the quadrature of the circle.

For, if the number π is not algebraic, it certainly cannot be constructed by means of straight edge and compasses. *The quadrature of the circle in the sense understood by the ancients is then impossible.* It is extremely interesting to follow the fortunes of this problem in the various epochs of science, as ever new attempts were made to find a solution with straight edge and compasses, and to see how these necessarily fruitless efforts worked for advancement in the manifold realm of mathematics.

The following brief historical survey is based upon the excellent work of Rudolph: *Archimedes, Huygens, Lambert, Legendre, Vier Abhandlungen über die Kreismessung*, Leipzig, 1892. This book contains a German translation of the investigations of the authors named. While the mode of presentation does not touch upon the modern methods here discussed, the book includes many interesting details which are of practical value in elementary teaching.

1. Among the attempts to determine the ratio of the diameter to the circumference we may first distinguish the *empirical stage*, in which the desired end was to be attained by measurement or by direct estimation.

One of the oldest known mathematical documents, the Rhind Papyrus (c. 1650 B.C.), contains the problem in the well-known form, to transform a circle into a square of equal area. The writer of the papyrus lays down the following rule: Cut off $\frac{1}{8}$ of a diameter and construct a square upon the remainder; this has the same area as the circle. The value of π thus obtained is $(\frac{15}{8})^2 = 3.16 \dots$; not very inaccurate. Much less accurate is the value $\pi = 3$, used in the Bible (1 Kings, 7. 23, 2 Chronicles, 4. 2).

2. The Greeks rose above this empirical standpoint, and especially Archimedes, who, in his work *κύκλου μέτρησις*, computed the area of the circle by the aid of inscribed and circumscribed polygons, as is still done in the schools. His method remained in use till the invention of the differential calculus; it was especially developed and rendered practical by Huygens (d. 1654) in his work, *De circuli magnitudine inventa*.

As in the case of the duplication of the cube and the trisection of the angle the Greeks sought also to effect the quadrature of the circle by the help of higher curves.

Consider for example the curve $y = \sin^{-1} x$, which represents the sinusoid with axis vertical. Geometrically, π appears as a particular ordinate of this curve; from the standpoint of the theory of functions, as a particular value of our transcendental function. Any apparatus which describes a transcendental curve we shall call a transcendental apparatus. A transcendental apparatus which traces the sinusoid gives us a geometric construction of π .

In modern language the curve $y = \sin^{-1} x$ is called an

integral curve because it can be defined by means of the integral of an algebraic function,

$$y = \int_0^x \frac{dx}{\sqrt{1-x^2}}.$$

The ancients called such a curve a *quadratrix* or τετραγωνίζουσα. The best known is the *quadratrix of Dinostratus* (c. 350 B.C.) which, however, had already been constructed by Hippias of Elis (c. 420 B.C.) for the trisection of an angle. Geometrically it may be defined as follows. Having given a circle and two perpendicular radii OA and OB, two points M and L move with constant velocity, one upon the radius OB, the other upon the arc AB (Fig. 14). Starting at the same time at O and A, they arrive simultaneously at B. The point of intersection P of OL and the parallel to OA through M describes the quadratrix.




FIG. 14.

From this definition it follows that y is proportional to θ . Further, since for $y = 1$, $\theta = \frac{\pi}{2}$ we have

$$\theta = \frac{\pi}{2} y;$$

and from $\theta = \tan^{-1} \frac{y}{x}$ the equation of the curve becomes

$$\frac{y}{x} = \tan \frac{\pi}{2} y.$$

It meets the axis of X at the point whose abscissa is

$$x = \lim_{y \rightarrow 0} \frac{y}{\tan \frac{\pi}{2} y}, \text{ for } y = 0;$$

hence

$$x = \frac{2}{\pi}.$$

According to this formula the radius of the circle is the mean proportional between the length of the quadrant and the abscissa of the intersection of the quadratrix with the axis of X . This curve can therefore be used for the rectification and hence also for the quadrature of the circle. This use of the quadratrix amounts, however, simply to a geometric formulation of the problem of rectification so long as we have no apparatus for describing the curve by continuous movement.

Fig. 15 gives an idea of the form of the curve with the branches obtained by taking values of θ greater than π or




FIG. 15.

less than $-\pi$. Evidently the quadratrix of Dinostratus is not so convenient as the curve $y = \sin^{-1} x$, but it does not appear that the latter was used by the ancients.

3. The period from 1670 to 1770, characterized by the names of Leibnitz, Newton, and Euler, saw the rise of modern analysis. Great discoveries followed one another in such an almost unbroken series that, as was natural, critical rigor fell into the background. For our purposes the development

of the theory of series is especially important. Numerous methods were deduced for approximating the value of π . It will suffice to mention the so-called *Leibnitz series* (known, however, before Leibnitz):

$$\frac{\pi}{4} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} \dots$$

This same period brings the discovery of the mutual dependence of e and π . The number e , natural logarithms, and hence the exponential function, are first found in principle in the works of Napier (1614). This number seemed at first to have no relation whatever to the circular functions and the number π until Euler had the courage to make use of imaginary exponents. In this way he arrived at the celebrated formula

$$e^{ix} = \cos x + i \sin x,$$

which, for $x = \pi$, becomes

$$e^{i\pi} = -1.$$

This formula is certainly one of the most remarkable in all mathematics. The modern proofs of the transcendence of π are all based on it, since the first step is always to show the transcendence of e .

4. After 1770 critical rigor gradually began to resume its rightful place. In this year appeared the work of Lambert: *Vorläufige Kenntnisse für die so die Quadratur des Cirkuls suchen*. Among other matters the irrationality of π is discussed. In 1794 Legendre, in his *Éléments de géométrie*, showed conclusively that π and π^2 are irrational numbers.

5. But a whole century elapsed before the question was investigated from the modern point of view. The starting-point was the work of Hermite: *Sur la fonction exponentielle* (*Comptes rendus*, 1873, published separately in 1874). The transcendence of e is here proved.

An analogous proof for π , closely related to that of Hermite, was given by Lindemann: *Ueber die Zahl π* (*Mathematische Annalen*, XX, 1882. See also the Proceedings of the Berlin and Paris academies).

The question was then settled for the first time, but the investigations of Hermite and Lindemann were still very complicated.

The first simplification was given by Weierstrass in the *Berliner Berichte* of 1885. The works previously mentioned were embodied by Bachmann in his text-book, *Vorlesungen über die Natur der Irrationalzahlen*, 1892.

But the spring of 1893 brought new and very important simplifications. In the first rank should be named the memoirs of Hilbert in the *Göttinger Nachrichten*. Still Hilbert's proof is not absolutely elementary: there remain traces of Hermite's reasoning in the use of the integral

$$\int_0^{\infty} z^p e^{-z} dz = p!$$

But Hurwitz and Gordan soon showed that this transcendental formula could be done away with (*Göttinger Nachrichten*; *Comptes rendus*; all three papers are reproduced with some extensions in *Mathematische Annalen*, Vol. XLIII).

The demonstration has now taken a form so elementary, that it seems generally available. In substance we shall follow Gordan's mode of treatment.

CHAPTER III.

The Transcendence of the Number e .

1. We take as the starting-point for our investigation the well-known series

$$e^x = 1 + \frac{x}{1} + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} + \dots,$$

which is convergent for all finite values of x . The difference between practical and theoretical convergence should here be insisted on. Thus, for $x=1000$ the calculation of e^{1000} by means of this series would obviously not be feasible. Still the series certainly converges theoretically; for we easily see that after the 1000th term the factorial $n!$ in the denominator increases more rapidly than the power which occurs in the numerator. This circumstance that $\frac{x^n}{n!}$ has for

any finite value of x the limit zero when n becomes infinite has an important bearing upon our later demonstrations.

We now propose to establish the following proposition:

The number e is not an algebraic number, i.e., an equation with integral coefficients of the form

$$F(e) = C_0 + C_1 e + C_2 e^2 + \dots + C_n e^n = 0$$

is impossible. The coefficients C_i may be supposed prime to one another.

We shall use the indirect method of demonstration, showing that the assumption of the above equation leads to an absurdity. The absurdity may be shown in the following

way. We multiply the members of the equation $F(e) = 0$ by a certain integer M so that

$$MF(e) = MC_0 + MC_1e + MC_2e^2 + \dots + MC_ne^n = 0.$$

We shall show that the number M can be chosen so that

(1) Each of the products Me, Me^2, \dots, Me^n may be separated into an entire part M_κ and a fractional part ϵ_κ , and our equation takes the form

$$MF(e) = MC_0 + M_1C_1 + M_2C_2 + \dots + M_nC_n \\ + C_1\epsilon_1 + C_2\epsilon_2 + \dots + C_n\epsilon_n = 0;$$

(2) The integral part

$$MC_0 + M_1C_1 + \dots + M_nC_n$$

is not zero. This will result from the fact that when divided by a prime number it gives a remainder different from zero;

(3) The expression

$$C_1\epsilon_1 + C_2\epsilon_2 + \dots + C_n\epsilon_n$$

can be made as small a fraction as we please.

These conditions being fulfilled, the equation assumed is manifestly impossible, since the sum of an integer different from zero, and a proper fraction, cannot equal zero.

The salient point of the proof may be stated, though not quite accurately, as follows:

With an exceedingly small error we may assume e, e^2, \dots, e^n proportional to integers which certainly do not satisfy our assumed equation.

2. We shall make use in our proof of a symbol h^r and a certain polynomial $\phi(x)$.

The symbol h^r is simply another notation for the factorial $r!$. Thus, we shall write the series for e^x in the form

$$e^x = 1 + \frac{x}{h} + \frac{x^2}{h^2} + \dots + \frac{x^n}{h^n} + \dots$$

The symbol has no deeper meaning ; it simply enables us to write in more compact form every formula containing powers and factorials.

Suppose, *e.g.*, we have given a developed polynomial

$$f(x) = \sum_r c_r x^r.$$

We represent by $f(h)$, and write under the form $\sum_r c_r h^r$, the sum

$$c_1 \cdot 1 + c_2 \cdot 2! + c_3 \cdot 3! + \cdots + c_n \cdot n!$$

But if $f(x)$ is not developed, then to calculate $f(h)$ is to develop this polynomial in powers of h and finally replace h^r by $r!$. Thus, for example,

$$f(k+h) = \sum_r c_r (k+h)^r = \sum_r c'_r \cdot h^r = \sum_r c'_r \cdot r!,$$

the c'_r depending on k .

The polynomial $\phi(x)$ which we need for our proof is the following remarkable expression

$$\phi(x) = x^{p-1} \frac{[(1-x)(2-x) \cdots (n-x)]^p}{(p-1)!},$$

where p is a prime number, n the degree of the algebraic equation assumed to be satisfied by e . We shall suppose p greater than n and $|C_0|$, and later we shall make it increase without limit.

To get a geometric picture of this polynomial $\phi(x)$ we construct the curve

$$y = \phi(x).$$

At the points $x = 1, 2, \cdots n$ the curve has the axis of X as an inflexional tangent, since it meets it in an odd number of points, while at the origin the axis of X is tangent without inflexion. For values of x between 0 and n the curve remains in the neighborhood of the axis of X ; for greater values of x it recedes indefinitely.

Of the function $\phi(x)$ we will now establish three important properties :

1. x being supposed given and p increasing without limit, $\phi(x)$ tends toward zero, as does also the sum of the absolute values of its terms.

Put $u = x(1-x)(2-x) \cdots (n-x)$; we may then write

$$\phi(x) = \frac{u^{p-1}}{(p-1)!} \frac{u}{x},$$

which for p infinite tends toward zero.

To have the sum of the absolute values of $\phi(x)$ it is sufficient to replace $-x$ by $|x|$ in the undeveloped form of $\phi(x)$. The second part is then demonstrated like the first.

2. h being an integer, $\phi(h)$ is an integer not divisible by p and therefore different from zero.

Develop $\phi(x)$ in increasing powers of x , noticing that the terms of lowest and highest degree respectively are of degree $p-1$ and $np+p-1$. We have

$$\phi(x) = \sum_{r=p-1}^{r=np+p-1} c_r x^r = \frac{c' x^{p-1}}{(p-1)!} + \frac{c'' x^p}{(p-1)!} + \cdots \pm \frac{x^{np+p-1}}{(p-1)!}.$$

Hence

$$\phi(h) = \sum_{r=p-1}^{r=np+p-1} c_r h^r.$$

Leaving out of account the denominator $(p-1)!$, which occurs in all the terms, the coefficients c_r are integers. This denominator disappears as soon as we replace h^r by $r!$, since the factorial of least degree is $h^{p-1} = (p-1)!$. All the terms of the development after the first will contain the factor p . As to the first, it may be written

$$\frac{(1 \cdot 2 \cdot 3 \cdots n)^p \cdot (p-1)!}{(p-1)!} = (n!)^p$$

and is certainly not divisible by p since $p > n$.

Therefore $\phi(h) \equiv (n!)^p \pmod{p}$,
and hence $\phi(h) \not\equiv 0$.

Moreover, $\phi(h)$ is a very large number; even its last term alone is very large, viz.:

$$\frac{(np + p - 1)!}{(p - 1)!} = p(p + 1) \cdot \dots \cdot (np + p - 1).$$

3. h being an integer, and k one of the numbers $1, 2 \cdot \dots \cdot n$, $\phi(h + k)$ is an integer divisible by p .

$$\text{We have } \phi(h + k) = \sum_r c_r (h + k)^r = \sum_r c'_r h^r,$$

a formula in which we are to replace h^r by $r!$ only after having arranged the development in increasing powers of h .

According to the rules of the symbolic calculus, we have first

$$\begin{aligned} \phi(h + k) \\ = (h + k)^{p-1} \frac{[(1-k-h)(2-k-h) \cdot \dots \cdot (-h) \cdot \dots \cdot (n-k-h)]^p}{(p-1)!}. \end{aligned}$$

One of the factors in the brackets reduces to $-h$; hence the term of lowest degree in h in the development is of degree p . We may then write

$$\phi(h + k) = \sum_{r=p}^{r=np+p-1} c'_r h^r.$$

The coefficients still have for numerators integers and for denominator $(p-1)!$. As already explained, this denominator disappears when we replace h^r by $r!$. But now all the terms of the development are divisible by p ; for the first may be written

$$\frac{(-1)^{kp} \cdot k^{p-1} [(k-1)!(n-k)!]^p \cdot p!}{(p-1)!} = (-1)^{kp} k^{p-1} [(k-1)! \cdot (n-k)!]^p \cdot p.$$

$\phi(h + k)$ is then divisible by p .

3. We can now show that the equation

$$F(e) = C_0 + C_1 e + C_2 e^2 + \dots + C_n e^n = 0$$

is impossible.

For the number M , by which we multiply the members of this equation, we select $\phi(h)$, so that

$$\phi(h) F(e) = C_0 \phi(h) + C_1 \phi(h) e + C_2 \phi(h) e^2 + \dots + C_n \phi(h) e^n.$$

Let us try to decompose any term, such as $C_k \phi(h) e^k$, into an integer and a fraction. We have

$$e^k \cdot \phi(h) = e^k \sum_r c_r h^r.$$

Considering the series development of e^k , any term of this sum, omitting the constant coefficient, has the form

$$e^k \cdot h^r = h^r + \frac{h^r \cdot k}{1} + \frac{h^r \cdot k^2}{2!} + \dots + \frac{h^r \cdot k^r}{r!} + \frac{h^r \cdot k^{r+1}}{(r+1)!} + \dots$$

Replacing h^r by $r!$, or what amounts to the same thing, by one of the quantities

$$r h^{r-1}, r(r-1) h^{r-2}, \dots, r(r-1) \dots 3 \cdot h^2, r(r-1) \dots 2 \cdot h,$$

and simplifying the successive fractions,

$$e^k \cdot h^r = h^r + \frac{r}{1} \cdot h^{r-1} k + \frac{r(r-1)}{2!} h^{r-2} k^2 + \dots + \frac{r}{1} h k^{r-1} + k^r \\ + k^r \left[\frac{k}{r+1} + \frac{k^2}{(r+1)(r+2)} + \dots \right].$$

The first line has the same form as the development of $(h+k)^r$; in the parenthesis of the second line we have the series

$$0 + \frac{k}{r+1} + \frac{k^2}{(r+1)(r+2)} + \dots$$

whose terms are respectively less than those of the series

$$e^k = 1 + k + \frac{k^2}{2!} + \frac{k^3}{3!} + \dots$$

The second line in the expansion of $e^k \cdot h^r$ may therefore be represented by an expression of the form

$$q_{r,k} e^k \cdot k^r,$$

$q_{r,k}$ being a proper fraction.

Effecting the same decomposition for each term of the sum

$$e^k \sum_r c_r h^r$$

it takes the form

$$e^k \sum_r c_r h^r = \sum_r c_r (h+k)^r + e^k \sum_r q_{r,k} c_r k^r.$$

The first part of this sum is simply $\phi(h+k)$; this is a number divisible by p (2, 3). Further (2, 1),

$$\phi(k) = \sum_r |c_r k^r|$$

tends toward zero when p becomes infinite: the same is true *a fortiori* of $\sum_r q_{r,k} c_r k^r$, and also, since e^k is a finite quantity, of $e^k \sum_r q_{r,k} c_r k^r$, which we may represent by ϵ_k .

The term under consideration, $C_k e^k \phi(h)$, has then been put under the form of an integer $C_k \phi(h+k)$ and a quantity $C_k \epsilon_k$ which, by a suitable choice of p , may be made as small as we please.

Proceeding similarly with all the terms, we get finally

$$F(e) \phi(h) = C_0 \phi(h) + C_1 \phi(h+1) + \cdots + C_n \phi(h+n) \\ + C_1 \epsilon_1 + C_2 \epsilon_2 + \cdots + C_n \epsilon_n.$$

It is now easy to complete the demonstration. All the terms of the first line after the first are divisible by p ; for the first, $|C_0|$ is less than p ; $\phi(h)$ is not divisible by p ; hence $C_0 \phi(h)$ is not divisible by the prime number p . Consequently the sum of the numbers of the first line is not zero.

The numbers of the second line are finite in number; each of them can be made smaller than any given number by a suitable choice of p ; and therefore the same is true of their sum.

Since an integer not zero and a fraction cannot have zero for a sum, the assumed equation is impossible.

Thus, the transcendence of e , or Hermite's Theorem, is demonstrated.

CHAPTER IV.

The Transcendence of the Number π .

1. The demonstration of the transcendence of the number π given by Lindemann is an extension of Hermite's proof in the case of e . While Hermite shows that an integral equation of the form

$$C_0 + C_1e + C_2e^2 + \dots + C_ne^n = 0$$

cannot exist, Lindemann generalizes this by introducing in place of the powers e, e^2, \dots sums of the form

$$\begin{array}{c} e^{k_1} + e^{k_2} + \dots + e^{k_N} \\ e^{l_1} + e^{l_2} + \dots + e^{l_{N'}} \\ \dots \end{array}$$

where the k 's are associated algebraic numbers, i.e., roots of an algebraic equation, with integral coefficients, of the degree N ; the l 's roots of an equation of degree N' , etc. Moreover, some or all of these roots may be imaginary.

Lindemann's general theorem may be stated as follows:

The number e cannot satisfy an equation of the form

$$(1) \quad C_0 + C_1(e^{k_1} + e^{k_2} + \dots + e^{k_N}) + C_2(e^{l_1} + e^{l_2} + \dots + e^{l_{N'}}) + \dots = 0$$

where the coefficients C_i are integers and the exponents k_i, l_i, \dots are respectively associated algebraic numbers.

The theorem may also be stated:

The number e is not only not an algebraic number and therefore a transcendental number simply, but it is also not an interscendental number and therefore a transcendental number of higher order.*

* Leibnitz calls a function x^λ , where λ is an algebraic irrational, an interscendental function.

Let

$$ax^n + a_1x^{n-1} + \dots + a_n = 0$$

be the equation having for roots the exponents k_1 ;

$$bx^{n'} + b_1x^{n'-1} + \dots + b_{n'} = 0$$

that having for roots the exponents l_1 , etc. These equations are not necessarily irreducible, nor the coefficients of the first terms equal to 1. It follows that the symmetric functions of the roots which alone occur in our later developments need not be integers.

In order to obtain integral numbers it will be sufficient to consider symmetric functions of the quantities

$$ak_1, ak_2, \dots, ak_n, \\ bl_1, bl_2, \dots, bl_{n'}, \text{ etc.}$$

These numbers are roots of the equations

$$y^n + a_1y^{n-1} + a_2ay^{n-2} + \dots + a_n a^{n-1} = 0, \\ y^{n'} + b_1y^{n'-1} + b_2by^{n'-2} + \dots + b_{n'} b^{n'-1} = 0, \text{ etc.}$$

These quantities are integral associated algebraic numbers, and their rational symmetric functions real integers.

We shall now follow the same course as in the demonstration of Hermite's theorem.

We assume equation (1) to be true; we multiply both members by an integer M ; and we decompose each sum, such as

$$M(e^{k_1} + e^{k_2} + \dots + e^{k_n}),$$

into an integral part and a fraction, thus

$$M(e^{k_1} + e^{k_2} + \dots + e^{k_n}) = M_1 + \epsilon_1, \\ M(e^{l_1} + e^{l_2} + \dots + e^{l_{n'}}) = M_2 + \epsilon_2, \\ \dots \dots \dots$$

Our equation then becomes

$$C_0M + C_1M_1 + C_2M_2 + \dots \\ + C_1\epsilon_1 + C_2\epsilon_2 + \dots = 0.$$

We shall show that with a suitable choice of M the sum of the quantities in the first line represents an integer not divisible by a certain prime number p , and consequently different from zero; that the fractional part can be made as small as we please, and thus we come upon the same contradiction as before.

2. We shall again use the symbol $h^r = r!$ and select as the multiplier the quantity $M = \psi(h)$, where $\psi(x)$ is a generalization of $\phi(x)$ used in the preceding chapter, formed as follows:

$$\psi(x) = \frac{x^{p-1}}{(p-1)!} [(k_1 - x)(k_2 - x) \cdots (k_n - x)]^p \cdot a^{np} \cdot a^{n'p} \cdot a^{n''p} \cdots \\ \cdot [(l_1 - x)(l_2 - x) \cdots (l_r - x)]^p \cdot b^{np} \cdot b^{n'p} \cdot b^{n''p} \cdots \\ \cdot \cdot \cdot \cdot \cdot \cdot \cdot \cdot$$

where p is a prime number greater than the absolute value of each of the numbers

$$C_0, a, b, \cdots, a_n, b_n, \cdots$$

and later will be assumed to increase without limit. As to the factors $a^{np}, b^{n'p}, \cdots$, they have been introduced so as to have in the development of $\psi(x)$ symmetric functions of the quantities

$$ak_1, ak_2, \cdots, ak_n, \\ bl_1, bl_2, \cdots, bl_r, \\ \cdot \cdot \cdot \cdot \cdot \cdot$$

that is, rational integral numbers. Later on we shall have to develop the expressions

$$\sum_v \psi(k_v + h), \quad \sum_v \psi(l_v + h), \quad \cdots$$

The presence of these same factors will still be necessary if we wish the coefficients of these developments to be integers each divided by $(p-1)!$.

1. $\psi(h)$ is an integral number, not divisible by p and consequently different from zero.

Arranging $\psi(h)$ in increasing powers of h , it takes the form

$$\psi(h) = \sum_{r=0}^{r=n_1p+n_2p+\dots+p-1} c_r h^r.$$

In this development all the coefficients have integral numerators and the common denominator $(p-1)!$.

The coefficient of the first term h^{p-1} may be written

$$\begin{aligned} & \frac{1}{(p-1)!} (ak_1 \cdot ak_2 \cdot \dots \cdot ak_n)^p a^{n^2p} a^{n^2p} \cdot \dots \cdot \\ & (bl_1 \cdot bl_2 \cdot \dots \cdot bl_n)^p b^{n^2p} b^{n^2p} \cdot \dots \cdot \\ & \dots \dots \dots \dots \dots \dots \\ & = \frac{1}{(p-1)!} (-1)^{n_1p+n_2p+\dots} (a_n a^{n-1})^p a^{n^2p} a^{n^2p} \cdot \dots \cdot (b_n b^{n-1})^p b^{n^2p} b^{n^2p} \cdot \dots \end{aligned}$$

If in this term we replace h^{p-1} by its value $(p-1)!$ the denominator disappears. According to the hypotheses made regarding the prime number p , no factor of the product is divisible by p and hence the product is not.

The second term $c_p h^p$ becomes likewise an integer when we replace h^p by $p!$ but the factor p remains, and so for all of the following terms. Hence $\psi(h)$ is an integer not divisible by p .

2. For x , a given finite quantity, and p increasing without limit, $\psi(x) = \sum_r c_r x^r$ tends toward zero, as does also the sum $\sum_r |c_r x^r|$.

We may write

$$\begin{aligned} \psi(x) &= \sum_r c_r x^r \\ &= \frac{x^{p-1}}{(p-1)!} [a^n a^n \cdot \dots \cdot b^n b^n (k_1 - x)(k_2 - x) \cdot \dots \cdot (k_n - x) \\ & \quad (l_1 - x)(l_2 - x) \cdot \dots \cdot (l_n - x) \cdot \dots]^p. \end{aligned}$$

Since for x of given value the expression in brackets is a constant, we may replace it by K . We then have

$$\psi(x) = \frac{(xK)^{p-1}}{(p-1)!} K,$$

a quantity which tends toward zero as p increases indefinitely.

The same reasoning will apply when each term of $\psi(x)$ is replaced by its absolute value.

3. The expression $\sum_{v=1}^{v=N} \psi(k_v + h)$ is an integer divisible by p .

We have

$$\begin{aligned} \psi(k_v + h) &= \frac{a^p (k_v + h)^{p-1}}{(p-1)!} b^{xp} b^{x^2 p} \dots \\ &\cdot a^{(x-1)p} [(k_1 - k_v - h)(k_2 - k_v - h) \dots (-h) \dots (k_x - k_v - h)]^p \\ &\cdot a^{x^2 p} b^{x^2 p} [(l_1 - k_v - h)(l_2 - k_v - h) \dots (l_N - k_v - h)]^p \\ &\dots \end{aligned}$$

The v th factor of the expression in brackets in the second line is $-h$, and hence the term of lowest degree in h is h^p .

Consequently

$$\psi(k_v + h) = \sum_{r=p}^{r=xp+x^2p+\dots+p-1} c'_r h^r,$$

whence

$$\sum_{v=1}^{v=N} \psi(k_v + h) = \sum_{r=p}^{r=xp+x^2p+\dots+p-1} C'_r h^r.$$

The numerators of the coefficients C'_r are rational and integral, for they are integral symmetric functions of the quantities

$$\begin{array}{cccc} ak_1, & ak_2, & \dots, & ak_x, \\ bl_1, & bl_2, & \dots, & bl_N, \\ \dots & \dots & \dots & \dots \end{array}$$

and their common denominator is $(p-1)!$.

If we replace h^r by $r!$ the denominator disappears from all the coefficients, the factor p remains in every term, and hence the sum is an integer divisible by p .

Similarly for

$$\sum_{v=1}^{v=N} \psi(l_v + h) \dots$$

We have thus established three properties of $\psi(x)$ analogous to those demonstrated for $\phi(x)$ in connection with Hermite's theorem.

3. We now return to our demonstration that the assumed equation

(1) $C_0 + C_1(e^{k_1} + e^{k_2} + \dots + e^{k_n}) + C_2(e^{l_1} + e^{l_2} + \dots + e^{l_n}) + \dots = 0$ cannot be true. For this purpose we multiply both members by $\psi(h)$, thus obtaining

$C_0\psi(h) + C_1[e^{k_1}\psi(h) + e^{k_2}\psi(h) + \dots + e^{k_n}\psi(h)] + \dots = 0$, and try to decompose each of the expressions in brackets into a whole number and a fraction. The operation will be a little longer than before, for k may be a complex number of the form $k = k' + ik''$. We shall need to introduce $|k| = +\sqrt{k'^2 + k''^2}$.

One term of the above sum is

$$e^k \cdot \psi(h) = e^k \sum_r c_r h^r = \sum_r c_r \cdot e^k \cdot h^r.$$

The product $e^k \cdot h^r$ may be written, as shown before,

$$e^k \cdot h^r = (h+k)^r + k^r \left[\frac{k}{r+1} + \frac{k^2}{(r+1)(r+2)} + \dots \right]$$

The absolute value of every term of the series

$$0 + \frac{k}{r+1} + \frac{k^2}{(r+1)(r+2)} + \dots$$

is less than the absolute value of the corresponding term in the series

$$e^k = 1 + \frac{k}{1} + \frac{k^2}{2!} + \dots$$

Hence
$$\left| \frac{k}{r+1} + \frac{k^2}{(r+1)(r+2)} + \dots \right| < e^{|k|}$$

or
$$\frac{k}{r+1} + \frac{k^2}{(r+1)(r+2)} + \dots = q_{r,k} e^{|k|},$$

$q_{r,k}$ being a complex quantity whose absolute value is less than 1.

We may then write

$$\begin{aligned} e^k \cdot \psi(h) &= \sum_r c_r e^k h^r = \sum_r c_r (h+k)^r + \sum_r c_r q_{r,k} k^r e^{|k|} \\ &= \psi(h+k) + \sum_r c_r q_{r,k} k^r \cdot e^{|k|}. \end{aligned}$$

By giving k in succession the indices $1, 2, \dots, n$, and forming the sum the equation becomes

$$\begin{aligned} e^{k_1}\psi(h) + e^{k_2}\psi(h) + \dots + e^{k_n}\psi(h) \\ = \sum_{\nu=1}^{\nu=n} \psi(k_\nu + h) + \sum_{\nu=1}^{\nu=n} \{e^{ik_\nu} \sum_r c_r k_\nu^r q_{r,k_\nu}\}. \end{aligned}$$

Proceeding similarly with all the other sums, our equation takes the form

$$\begin{aligned} (2) \quad C_0\psi(h) + C_1 \sum_{\nu=1}^{\nu=n} \psi(k_\nu + h) + C_2 \sum_{\nu=1}^{\nu=n'} \psi(l_\nu + h) + \dots \\ + C_1 \sum_{\nu=1}^{\nu=n} \sum_r e^{ik_\nu} c_r k_\nu^r q_{r,k_\nu} + C_2 \sum_{\nu=1}^{\nu=n'} e^{il_\nu} c'_r l_\nu^r q_{r,l_\nu} + \dots = 0. \end{aligned}$$

By 2, 2 we can make $\sum_r |c_r k^r|$ as small as we please by taking p sufficiently great. Since $|q_{r,k}| < 1$, this will be true *a fortiori* of

$$\sum_r c_r k^r q_{r,k}$$

and hence also of

$$\sum_{\nu=1}^{\nu=n} \sum_r c_r k_\nu^r q_{r,k} e^{ik_\nu}.$$

Since the coefficients C are finite in value and in number, the sum which occurs in the second line of (2) can, by increasing p , be made as small as we please.

The numbers of the first line are, after the first, all divisible by p (3), but the first number, $C_0\psi(h)$, is not (1). Therefore the sum of the numbers in the first line is not divisible by p and hence is different from zero. The sum of an integer and a fraction cannot be zero. Hence equation (2) is impossible and consequently also equation (1).*

4. We now come to a proposition more general than the preceding, but whose demonstration is an immediate conse-

* The proof for the more general case where $C_0 = 0$ may be reduced to this by multiplication by a suitable factor, or may be obtained directly by a proper modification of $\psi(h)$.

quence of the latter. For this reason we shall call it Lindemann's corollary.

The number e cannot satisfy an equation of the form

$$(3) \quad C_0 + C_1 e^{k_1} + C_2 e^{l_1} + \dots = 0,$$

in which the coefficients are integers even when the exponents k_1, l_1, \dots are unrelated algebraic numbers.

To demonstrate this, let k_2, k_3, \dots, k_π be the other roots of the equation satisfied by k_1 ; similarly for l_2, l_3, \dots, l_π , etc. Form all the polynomials which may be deduced from (3) by replacing k_1 in succession by the associated roots k_2, \dots, l_1 by the associated roots l_2, \dots . Multiplying the expressions thus formed we have the product

$$\prod_{\alpha, \beta, \dots} \{ C_0 + C_1 e^{k_\alpha} + C_2 e^{l_\beta} + \dots \} \quad \left[\begin{array}{l} \alpha = 1, 2, \dots, \pi \\ \beta = 1, 2, \dots, \pi' \\ \dots \end{array} \right]$$

$$= C_0 + C_1 (e^{k_1} + e^{k_2} + \dots + e^{k_\pi}) + C_2 (e^{k_1+k_2} + e^{k_2+k_3} + \dots) \\ + C_3 (e^{k_1+l_1} + e^{k_1+l_2} + \dots) + \dots$$

In each parenthesis the exponents are formed symmetrically from the quantities k_1, l_1, \dots ; and are therefore roots of an algebraic equation with integral coefficients. Our product comes under Lindemann's theorem; hence it cannot be zero. Consequently none of its factors can be zero and the corollary is demonstrated.

We may now deduce a still more general theorem.

The number e cannot satisfy an equation of the form

$$C_0^{(1)} + C_1^{(1)} e^k + C_2^{(1)} e^l + \dots = 0$$

where the coefficients as well as the exponents are unrelated algebraic numbers.

For, let us form all the polynomials which we can deduce from the preceding when for each of the expressions $C_i^{(1)}$ we substitute one of the associated algebraic numbers

$$C_i^{(2)}, C_i^{(3)}, \dots, C_i^{(\pi)}.$$

If we multiply the polynomials thus formed together we get the product

$$\prod_{\alpha, \beta, \gamma, \dots} \{ C_0^{(\alpha)} + C_1^{(\beta)} e^k + C_2^{(\gamma)} e^l + \dots \} \quad \left[\begin{array}{l} \alpha = 1, 2, \dots, N_0 \\ \beta = 1, 2, \dots, N_1 \\ \gamma = 1, 2, \dots, N_2 \\ \dots \end{array} \right]$$

$$= C_0 + C_k e^k + C_l e^l + \dots$$

$$+ C_{k,k} e^{k+k} + C_{k,l} e^{k+l} + \dots$$

$$+ \dots$$

$$+ \dots$$

where the coefficients C are integral symmetric functions of the quantities

$$\begin{array}{cccc} C_0^{(1)}, & C_0^{(2)}, & \dots, & C_0^{(\pi_0)}, \\ C_1^{(1)}, & C_1^{(2)}, & \dots, & C_1^{(\pi_1)}, \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \end{array}$$

and hence are rational. By the previous proof such an expression cannot vanish, and we have accordingly Lindemann's corollary in its most general form :

The number e cannot satisfy an equation of the form

$$C_0 + C_1 e^k + C_2 e^l + \dots = 0$$

where the exponents k, l, \dots as well as the coefficients C_0, C_1, \dots are algebraic numbers.

This may also be stated as follows :

In an equation of the form

$$C_0 + C_1 e^k + C_2 e^l + \dots = 0$$

the exponents and coefficients cannot all be algebraic numbers.

5. From Lindemann's corollary we may deduce a number of interesting results. First, the transcendence of π is an immediate consequence. For consider the remarkable equation

$$1 + e^{i\pi} = 0.$$

The coefficients of this equation are algebraic; hence the exponent $i\pi$ is not. Therefore, π is transcendental.

6. Again consider the function $y = e^x$. We know that $1 = e^0$. This seems to be contrary to our theorems about the transcendence of e . This is not the case, however. We must notice that the case of the exponent 0 was implicitly excluded. For the exponent 0 the function $\psi(x)$ would lose its essential properties and obviously our conclusions would not hold.

Excluding then the special case ($x=0, y=1$), Lindemann's corollary shows that in the equation $y = e^x$ or $x = \log_e y$, y and x , i.e., the number and its natural logarithm, cannot be algebraic simultaneously. To an algebraic value of x corresponds a transcendental value of y , and conversely. This is certainly a very remarkable property.

If we construct the curve $y = e^x$ and mark all the algebraic points of the plane, i.e., all points whose coördinates are algebraic numbers, the curve passes among them without meeting a single one except the point $x=0, y=1$. The theorem still holds even when x and y take arbitrary complex values. The exponential curve is then transcendental in a far higher sense than ordinarily supposed.

7. A further consequence of Lindemann's corollary is the transcendence, in the same higher sense, of the function $y = \sin^{-1} x$ and similar functions.

The function $y = \sin^{-1} x$ is defined by the equation

$$2ix = e^{iy} - e^{-iy}.$$

We see, therefore, that here also x and y cannot be algebraic simultaneously, excluding, of course, the values $x=0, y=0$. We may then enunciate the proposition in geometric form:

The curve $y = \sin^{-1} x$, like the curve $y = e^x$, passes through no algebraic point of the plane, except $x=0, y=0$.

CHAPTER V.

The Integrator and the Geometric Construction of π .

1. Lindemann's theorem demonstrates the transcendence of π , and thus is shown the impossibility of solving the old problem of the quadrature of the circle, not only in the sense understood by the ancients but in a far more general manner. It is not only impossible to construct π with straight edge and compasses, but there is not even a curve of higher order defined by an integral algebraic equation for which π is the ordinate corresponding to a rational value of the abscissa. An actual construction of π can then be effected only by the aid of a transcendental curve. If such a construction is desired, we must use besides straight edge and compasses a "transcendental" apparatus which shall trace the curve by continuous motion.

2. Such an apparatus is the *integrator*, recently invented and described by a Russian engineer, Abdank-Abakanowicz, and constructed by Coradi of Zürich.

This instrument enables us to trace the *integral curve*

$$Y = F(x) = \int f(x) dx$$

when we have given the *differential curve*

$$y = f(x).$$

For this purpose, we move the linkwork of the integrator so that the *guiding point* follows the differential curve; the *tracing point* will then trace the integral curve. For a fuller description of this ingenious instrument we refer to the original memoir (in German, Teubner, 1889; in French, Gauthier-Villars, 1889).

We shall simply indicate the principles of its working. For any point (x, y) of the differential curve construct the auxiliary triangle having for vertices the points (x, y) , $(x, 0)$, $(x-1, 0)$; the hypotenuse of this right-angled triangle makes with the axis of X an angle whose tangent $= y$.

Hence, *this hypotenuse is parallel to the tangent to the integral curve at the point (X, Y) corresponding to the point (x, y) .*




FIG. 16.

The apparatus should be so constructed then that the tracing point shall move parallel to the variable direction of this hypotenuse, while the guiding point describes the differential curve. This is effected by connecting the tracing point with a sharp-edged roller whose plane is vertical and moves so as to be always parallel to this hypotenuse. A weight presses this roller firmly upon the paper so that its point of contact can advance only in the plane of the roller.

The practical object of the integraph is the approximate evaluation of definite integrals; for us its application to the construction of π is of especial interest.

3. Take for differential curve the circle

$$x^2 + y^2 = r^2;$$

the integral curve is then

$$Y = \int \sqrt{r^2 - x^2} dx = \frac{r^2}{2} \sin^{-1} \frac{x}{r} + \frac{x}{2} \sqrt{r^2 - x^2}.$$

This curve consists of a series of congruent branches. The points where it meets the axis of Y have for ordinates

$$0, \quad \pm \frac{r^2 \pi}{2}, \quad \dots$$

Upon the lines $X = \pm r$ the intersections have for ordinates

$$r^2 \frac{\pi}{4}, \quad r^2 \frac{3\pi}{4}, \quad \dots$$

If we make $r=1$, the ordinates of these intersections will determine the number π or its multiples.

It is worthy of notice that our apparatus enables us to trace the curve not in a tedious and inaccurate manner, but with ease and sharpness, especially if we use a tracing pen instead of a pencil.

Thus we have an actual constructive quadrature of the circle along the lines laid down by the ancients, for our curve is only a modification of the quadratrix considered by them.

**NOTES FOR
PART I AND PART II**

NOTES

PART I — CHAPTER III

Gaussian Polygons. Up to the time of Gauss, no one suspected that it was possible to construct, with ruler and compasses, regular polygons other than those the number of whose sides could be expressed in one of the forms: 2^n , $2^n \cdot 3$, $2^n \cdot 5$, $2^n \cdot 15$. All of these were known to the Greeks. But Gauss proved as early as 1801¹ that whenever a prime number F_μ could be expressed in the form $2^{2^\mu} + 1$, the construction of a regular polygon with F_μ sides was possible by Euclidean methods. It was then apparent that regular polygons not included in the Euclidean series, namely 17, 257, 65537, . . . sides, could be constructed under the same imposed conditions. And indeed Gauss's discussion led to the result², that the *only* regular polygons which it is possible to construct with ruler and compasses, are those the number P of whose sides can be expressed in the form

$$2^\alpha \cdot (2^{2^{\alpha_1}} + 1) \cdot (2^{2^{\alpha_2}} + 1) \cdot (2^{2^{\alpha_3}} + 1) \cdots (2^{2^{\alpha_s}} + 1),$$

where $\alpha \dots \alpha_s$ are distinct positive integers and each $2^{2^{\alpha_i}} + 1$ is a prime. The number of such polygons is small in comparison with the number of regular polygons which can not be constructed with the means employed. As Dickson has pointed out³ the number of P's up to 100 is 24; up to 300 is 37 (all noted by Gauss); up to 1000 is 52; up to 1000000 only 206. Kraitichik has remarked⁴ that there are only 30 polygons with an odd number of sides that are known to be constructible with ruler and compasses. These polygons have the following number of sides: 5, 15, 17, 51, 85, 255, 257, 771, 1285, 3855, 4369, 13107, 21845, 65535, 65537, 196611, 327685, 983055, 1114129, 3342387, 5570645, 16711935, 16843009, 50529027, 84215045, 252645135, 286331153, 858993459, 1431655765,

¹ *Disquisitiones arithmeticae*, Leipzig, 1801, p. 664; *Werke*, v. 1., 2. Abdruck, 1870, p. 462; French ed. *Recherches Arithmétiques*, Paris, 1807, p. 488; Ger. ed. by Maser, Berlin, 1889, p. 447.

² This result was, in effect, stated, but not proved, by Gauss.

³ L. E. Dickson, "On the number of inscriptible regular polygons", *Bull. N. Y. Math. Soc.*, Feb., 1894, v. 3, p. 123.

⁴ Kraitichik, *Recherches sur la théorie des nombres*, Paris, 1924, p. 270.

4294967295. This set of numbers, together with 1 and 3, coincides with the divisors of $2^{32} - 1 = 1 \cdot 3 \cdot 5 \cdot 17 \cdot 257 \cdot 65537$.

The determination of the number of regular polygons which can be constructed for P less than a given integer is, then, bound up in the determination of the prime numbers F_μ . Now for only 18 values of μ has it been shown whether F_μ is prime or not, namely for the values of μ from 0 to 9 inclusive, and for 11, 12, 15, 18, 23, 36, 38, 73. In the first five of these cases, and in these alone, is F_μ prime. These five cases were noted by Fermat in the seventeenth century. It may well turn out that F_μ is not prime, for $\mu > 4$, although Eisenstein proposed as a problem¹: "There are an infinity of prime numbers of the form $2^{2^\mu} + 1$ ".

The results already established in this connection may be set forth in tabular form²:

μ	Prime Factors of $F_\mu = 2^{2^\mu} + 1$	Discoverer	Year of Discovery
0-4	3, 5, 17, 257, 65537	Fermat	1640
5	$\{2^7 \cdot 5 + 1 = 641$ $\{2^7 \cdot 52347 + 1 = 6700417\}$	L. Euler	1732
6	$\{ \text{Unknown but composite.}$ $2^8 \cdot 9 \cdot 7 \cdot 17 + 1 = 274177$	Lucas Landry	1878 1880
7	$2^8 \cdot 5 \cdot 52562829149 + 1 + 67280421310721$ Unknown but composite.	Landry and Le Lasseur A. E. Western, J. C. Morehead	1880 1905
8	Unknown but composite.	A. E. Western J. C. Morehead	1909
9	$2^{16} \cdot 37 + 1 = 2424833$	A. E. Western	1903
11	$\{2^{13} \cdot 3 \cdot 13 + 1 = 319489\}$ $\{2^{13} \cdot 7 \cdot 17 + 1 = 974849\}$	A, Cunningham	1899
12	$\{2^{14} \cdot 7 + 1 = 114689$ $\{2^{16} \cdot 397 + 1 = 26017793\}$ $\{2^{16} \cdot 7 \cdot 139 + 1 = 63766529\}$	E. A. Lucas and P. Pervouchine A. E. Western	1877 1903
15	$2^{21} \cdot 579 + 1 = 1214251009$	M. Kraitchik	1925
18	$2^{29} \cdot 13 + 1 = 13631489$	A. E. Western	1903
23	$2^{25} \cdot 5 + 1 = 167772161$	P. Pervouchine	1878
36	$2^{39} \cdot 5 + 1 = 2748779069441$	Seelhoff	1886
38	$2^{41} \cdot 3 + 1 = 6597069766657$	$\{ \text{J. Cullen, A. Cunningham,}$ $\{ \text{A. E. and F. J. Western}$	1903
73	$2^{75} \cdot 5 + 1 = 188894559314785808547841$	J. C. Morehead	1906

¹ G. Eisenstein, "Aufgaben", Crelle's *Journal*, v. 27, 1844, p. 87.

² The sources for the different results, except those of Fermat, are as follows, for the 13 different values of μ :

The labor expended in deriving these results has been enormous; to the layman who knows nothing of congruences in the theory of numbers, the facts found must seem almost to border on the miraculous. For, even when $\mu = 10$, a case not yet solved, F_μ contains 309 digits; but when $\mu = 36$, F_μ is a number of more than twenty trillion digits. Concerning it Lucas remarked¹ "la bande de papier qui le contiendrait ferait le tour de la Terre". For $\mu = 73$, Ball states that the digits in F_μ "are so numerous that, if the number were printed in full with the type and number of pages used in this book [*Mathematical Recreations*, fifth edition, 1911, 508 pages], many more

5. L. Euler, *Commentarii Academiæ Scientiarum Petrop.*, v. 6 (1732—3), 1738, p. 104; laid before the Academy of St. Petersburg, 26. Sept. 1732.

In his autobiography (Springfield, Mass., 1833, p. 38) the American calculator Zora Colburn records that while on exhibition in London, at the age of 8, he found "by the mere operation of his mind" the factors 641 and 6,700,417 of 4,294,967,297 ($= 2^{32} + 1$). Cf. F. D. Mitchell, "Mathematical prodigies", *Amer. Journal of Psychology*, v. 18, 1907, p. 65.

6. Lucas, *Comptes Rendus de l'Académie des Sciences*, Paris, v. 85, 1878, p. 138; *Amer. Jour. Math.*, v. 1, 1878, p. 238; *Recreations mathématiques*, v. 2 (2e éd., 1896), p. 234—5. Landry, *Nouv. Corresp. Math.*, v. 6, 1880, p. 417.

7. Independent discoverers: Western, *Proc. Lond. Math. Soc.*, s. 2, v. 3, p. xxi—xxii. Abstract of paper read, April 13, 1905; Morehead, *Bull. Amer. Math. Soc.*, v. 11, p. 543—545, abstract of paper read April 29, 1905.

8. Western and Morehead, *Bull. Amer. Math. Soc.*, v. 16, 1909, p. 1—6; "each doing half of the whole work".

9, 12 (Western), 13, 16. *Proc. Lond. Math. Soc.*, s. 2, v. 1, 1903, p. 175; abstract of paper read May 14, 1903.

11. A. Cunningham, *Brit. Assoc. Rept.*, 1899, p. 653—4; the factors are here given as 319489 and 974489. The second number is incorrect, 4 and 8 being interchanged. The other forms of the correct factors were given by A. Cunningham and A. E. Western in *Proc. Lond. Math. Soc.*, s. 2, v. 1, 1903, p. 175. It is here noted also that there are no more factors of $F_\mu < 10^6$, and no other factor of $F_\mu < 10^8$, (μ not less than 14).

12, 23. E. Lucas, *Atti Accad. Torino*, v. 13 (1877—8), p. 271 [27 Jan., 1878]. *Mélanges math. ast. acad. Pétersb.*, v. 5, part 5, 1879, p. 505, 519, or *Bull. Acad. Pétersb.*, s. 3, v. 24, 1878, p. 559; s. 3, v. 25, 1879, p. 63; communication of results, for $\mu = 12$ and 23, found by J. Pervouchine, in Nov. 1877 and Jan. 1878. He notes that the integer $2^{2^{\mu}} + 1$ contains 2525223 digits.

15. M. Kraitchik, *Comptes Rendus de l'Académie des Sciences*, Paris, v. 180, p. 800, March, 1925; also *Sphinx-Oedipe*, v. 20, p. 24.

36. P. Seelhoff, *Zeitschrift math. u. Phys.*, v. 31, 1886, p. 174.

73. J. C. Morehead, *Bull. Amer. Math. Soc.*, v. 12, 1906, p. 449—451.

¹ E. Lucas, *Théorie des nombres*, Paris, v. 1, 1891, p. 51.

volumes would be required than are contained in all the public libraries of the world".

In not less than seven places¹, during the years 1640–58, did Fermat refer to $F_\mu = 2^{2^\mu} + 1$ as representing a series of prime numbers; but in no place did he claim that F_μ was always prime.

Gauss's Statement of his Polygon Results. In two passages the implication to be drawn from what Klein has written is, that Gauss published a proof that a regular polygon of p sides can not be constructed by ruler and compasses if p is a prime not of the form $2^k + 1$. The passages to which I refer are (pages 2, 16):

(1) "Gauss added other cases [to Euclid's] by showing the possibility of the division into parts where p is a prime number of the form $p = 2^{2^\mu} + 1$, and the impossibility for all other numbers"; (2) "Gauss extended this series of numbers [Euclid's] by showing that the division is possible for every prime number of the form $p = 2^{2^\mu} + 1$ but impossible for all other prime numbers and their powers". Now the implication referred to above is not correct, as Pierpont interestingly set forth in his paper "On an undemonstrated theorem of the *Disquisitiones Arithmeticae*"². That is, Gauss *did not* give a proof of the "impossibility" referred to in the quotations. But after proving the "possibility" described above he continued as follows:

"As often as $p-1$ contains other prime factors besides 2, we arrive at higher equations³, namely, to one or more cubic equations, if 3 enters

¹ Letter dated Aug. [?] 1640 to Fronicle (*Oeuvres de Fermat*, v. 2, 1894, p. 206); letter dated 18 Oct., 1640, to Fronicle (*Oeuvres*, v. 2, 1894, p. 208); *Varia Opera*, Toulouse, 1679, p. 162; Brassine's *Précis*, Toulouse, 1853, p. 142–3; letter dated 25 Dec., 1640, to Mersenne (*Oeuvres*, v. 2, p. 212–213); "De solutione problematum geometricorum per curvas simplicissimas et unicuique problematum generi proprie convenientes, Dissertatio tripartita" (*Oeuvres de Fermat*, v. 1, 1891, p. 130–131; French translation, v. 3, 1896, p. 120; *Varia Opera*, 1679 [reprint, 1861], p. 115); letter dated 29 August, 1654, to Pascal (*Oeuvres de Pascal*, v. 4, Paris, 1819, p. 384; *Oeuvres de Fermat*, v. 2, 1894, p. 309–310); letter to Sir Kenelm Digby, sent by Digby to Wallis, 19 June, 1658 (*Oeuvres de Fermat*, v. 2, 1894, p. 402, 404–5; French translation of the Latin, v. 3, 1896, p. 314, 316); letter dated August, 1659 to Carcavi, copy sent by Carcavi to Huygens 14 August, 1659 (*Corresp. de Huygens* no. 651; *Oeuvres de Fermat*, v. 2, p. 433–434).

² *Bull. Amer. Math. Soc.*, v. 2, 1895, p. 77–83.

³ In his earlier discussion of an inscribed polygon of p sides, Gauss considers the equation $x^p - 1 = 0$ and the resulting equation got by dividing out the factor $x - 1$, where p is a prime.


once or oftener as a factor of $p-1$, to equations of 5th degree if $p-1$ is divisible by 5, etc. And we can prove with all rigour that these equations cannot be avoided or made to depend upon equations of lower degree; and although the limits of this work do not permit us to give the demonstration here, we still thought it necessary to signal this fact in order that one should not seek to construct other polygons than those given by our theory, as, for example, polygons of 7, 11, 13, 19 sides, and so employ one's time in vain."

Fermat's Theorem. This theorem (p. 17) was indicated by Fermat in a letter, dated 18 October 1640, to B. Frenicle de Bessy (*Oeuvres de Fermat*, v. 2, 1894, p. 209). Euler gave two proofs (*Comment. Acad. Petrop.*, v. 8 for 1736, 1741, p. 141, and *Comment. Nov. Acad. Petrop.*, v. 7 for 1758-59, 1761, p. 49). Other proofs are due to Lagrange (*Nouv. Mém. de l'Acad. de Berlin*, 1771) and to Gauss (*Disquisitiones Arithmeticae*, § 49)

PART I — CHAPTER IV

Geometrical Constructions of the Regular Heptadecagon. The remark of Klein (p. 24, 32) that we possess as yet no method of construction of the regular polygon of seventeen sides, based upon considerations purely geometrical, is a little curious, since several constructions of this kind have been given. One by Erchinger was indeed reported by Gauss in 1825¹. The construction is as follows:

Let D, B, G, A, I, F, C, E be points on a line determined by constructions about to be given. Let AB be a line of any length. Produce it both ways to C and D so that,



$$AC \times BC = AB \times BD = 4 AB^2.$$

¹ *Göttingische gelehrte Anzeigen*, Dec. 19, 1825, no. 203, p. 2025; *Werke*, v. 2, p. 186-7. To Art. 365 of the *Disquisitiones Arithmeticae* Gauss added this note in his handwriting: "Circulum in 17 partes divisibilem esse geometricè, deteximus 1796 Mart. 30". Cf. *Werke* v. 1, p. 476 and v. 10, 1917, p. 3-4, 120-126, 488. The discovery of the result was first announced in the *Intelligenzblatt* of the *Allgemeine Literatur-Zeitung*, no. 66, 1 June, 1796, col. 554.

Further determine the points, E , G , on both sides of CA produced so that,

$$AE \times EC = AG \times CG = \overline{AB^2};$$

and find the point F on the side A of the line BA produced, such that

$$AF \times DF = \overline{AB^2}.$$

Finally divide AE in I so that

$$AI \times EI = AB \times AF,$$

where AI is the smaller, and EI the larger part of AE . Then construct a triangle, in which each of two sides equals AB , the third being equal to AI . About this triangle describe a circle; then AI will be one side of the regular inscribed polygon of seventeen sides.

Gauss particularly remarks that the author gave a purely synthetic proof of this construction.

Another synthetic construction and proof dated "Dublin, 17th October, 1819" was published by Samuel James in the *Transactions of the Irish Academy*¹. Yet another construction was given by John Lowry in *The Mathematical Repository*² for 1819. But the earliest published geometrical construction was given by Huguenin in his *Mathematische Beiträge zur weiteren Ausbildung angehender Geometer*, Königsberg, 1803, p. 283.

A score of geometrical constructions are assembled in A. Goldenring, *Die elementargeometrischen Konstruktionen des regelmässigen Siebzehnecks*, Leipzig, 1915. See also the review of this work in *Bull. Amer. Math. Soc.*, v. 22, 1916, p. 239—246, and my note "Gauss and the regular polygon of seventeen sides" in *Amer. Math. Monthly*, v. 27, 1920, p. 323—326.

The discovery that the regular polygon of seventeen sides could be constructed with ruler and compasses was not only one of which Gauss was vastly proud throughout his life, but also, according to Sartorius von Waltershausen³, the one which decided him to dedicate his life to the study of mathematics. Archimedes expressed the wish that a sphere inscribed in a cylinder be inscribed on his tomb, as Ludolf van Ceulen did in connection with the value of π to 35 decimal

¹ V. 13 (1818), p. 175—187; paper read Jan. 24, 1820.

² N. s., v. 4, p. 160. Lowry's proof occupies p. 160—168.

³ *Gauss zum Gedächtniss*, Leipzig, 1856, p. 16.

places, and Jacques Bernoulli with reference to the logarithmic spiral. So also, according to Weber¹, Gauss requested that the regular polygon of seventeen sides should be engraved on his tombstone. While this request was not granted, as it was in each of the other cases mentioned, it is engraved on the side of a monument to Gauss in Braunschweig, his birthplace.

Constructions in general with Ruler and Compasses. Regarding constructions as effected when intersections of circles with circles or lines, or of lines with lines may be determined, it can be shown that: *Every problem solved with ruler and compasses can be solved with compasses alone.* This was first shown by Georg Mohr in his *Euclides Danicus* published at Amsterdam in 1672; this work was reprinted in 1928 by the Danish Society of Sciences. Klein refers (p. 33) only to Mascheroni's proof of this result 125 years later, in his *Geometria del Compasso*. Of this work there were two French editions *Géométrie du Compas*, Paris, 1798 and 1828. From the first of these a German edition *L. Mascheroni's Gebrauch des Zirkels*, Berlin, 1825, was prepared by J. P. Gruson. The subject is treated in English by: A. Cayley, *Messenger of Math.*, v. 14, 1885, p. 179—181; *Collected Papers*, v. 12, p. 314—317; by E. W. Hobson, in a presidential address, *Mathematical Gazette*, v. 7, 1913, p. 49—54; by H. P. Hudson, *Ruler & Compasses*, London, 1916, p. 131—143; and by J. Coolidge, *Treatise on the Circle and Sphere*, Oxford, 1916, p. 186—188.

Klein has noted (p. 33—34) that Poncelet first conceived the result that *given a circle and its center, every solution of a problem with ruler and compasses can be carried through with ruler alone.* A little later Klein states (p. 34) "we will show how with the straight edge and one fixed circle we can solve every quadratic equation". This is not possible; Klein should have had "with its center" after "one fixed circle". That the center be also given is very essential when only one circle is given. Hilbert suggested the problem: How many given circles in a plane are necessary in order to determine with ruler alone, the center of one of them? In 1912 D. Cauer² showed: (a) If two circles do not intersect in real points it is generally impossible to determine the center of either circle with ruler alone; (b) A center

¹ *Encyclopädie der elementaren Algebra und Analysis* bearbeitet von H. Weber. 2. ed. Leipzig, 1906, p. 362.

² *Mathematische Annalen*, v. 73, 1912, p. 90—94; v. 74, 1913, p. 462—464.

may be determined if the circles cut in real points, touch, or are concentric. About the same time J. Grossmann discovered a result which proved that *Every problem solvable with ruler and compasses can also be solved with ruler alone if we are given, in the plane of construction, three linearly independent circles.* Correct proofs of this result were given by Schur and Mierendorff.

From this it is clear that every construction with ruler and compasses can be effected with a ruler, and compasses with a fixed opening. Constructions of this kind were found already in the tenth century by Abû'l Wefâ of Bagdad¹. With such means, in the sixteenth century, certain problems of Euclid were solved by Cardano, Ferraro, and Tartaglia. At Venice in 1553 G. B. Benedetti published a little treatise, *Resolutio omnium Euclidis problematum, aliorumque ad hoc necessario inventorum, una tantummodo circuli data apertura.* In English the topic is treated in a rare pamphlet translated from the Dutch by Joseph Moxon², and in an article by J. S. Mackay³.

Every problem whose solution is possible by ruler and compasses can be also solved with a two edged ruler alone, whether the edges are parallel or meet in a point. For some of the literature in this connection the following sources may be consulted: *Nouvelle Corresp. Math.*, v. 3, 1877, p. 204—208; v. 5, 1879, p. 439—442; v. 6, 1880, p. 34—35; Akademie der Wissen., Vienna, *Sitzungsberichte*, Abt. IIa, v. 99, 1890, p. 854—858; *Bolletino di Matematiche e di Scienze fisiche e naturali*, v. 2, 1900—01, p. 129—145, 225—237.

PART II — CHAPTER II

Irrationality of π . Klein wrote (p. 59): "After 1770 critical rigour gradually began to resume its rightful place. In this year appeared the work of Lambert: *Vorläufige Kenntnisse für die, so die Quadratur*

¹ "Woepcke" *Analyse et extrait d'un recueil de constructions géométriques par Aboul Wafâ*, *Journal Asiatique*, 1855.

² *Compendium Euclidis Curiosum: or, geometrical operations. Showing how with a single opening of the Compasses and a straight ruler all the propositions of Euclid's first five books are performed.* London, 1677. Moxon does not tell us who the author of the Dutch treatise was.

³ "Solutions of Euclid's problems, with a ruler and one fixed aperture of the compasses, by the Italian geometers of the sixteenth century", *Proc. Edinb. Math. Soc.*, v. 5, 1887, p. 2—22.

...des Cirkuls suchen. Among other matters the irrationality of π is discussed. In 1794 Legendre in his *Éléments de Géométrie* showed conclusively that π and π^2 are irrational numbers." The implication of this note is that Lambert did not discuss the irrationality of π conclusively and that Legendre did. How both of these points of view are essentially incorrect will appear in what follows. Klein was simply reproducing the erroneous statements of Rudio¹; but after Pringsheim's careful study in 1898², Lambert's proof emerged as "ausserordentlich scharfsinnig und im wesentlichen vollkommen einwandfrei", while Legendre's remained "in Bezug auf Strenge hinter Lambert weit zurück".

As in the later proof of the transcendence of π , so here when its irrationality was in question, discussion of e is fundamental. The irrationality of e and e^2 was shown, substantially, by Euler in 1737³ and he gave the expression for e as a continued fraction on which Lambert's proofs of the irrationality of e^x , $\tan x$ and π rest. Starting with Euler's development⁴

$$\frac{e-1}{2} = \frac{1}{1+\frac{1}{6+\frac{1}{10+\frac{1}{14+\frac{1}{18+\dots}}}}}$$

Lambert found

$$\frac{ex-1}{ex+1} = \frac{1}{2/x+\frac{1}{6/x+\frac{1}{10/x+\frac{1}{14/x+\dots}}}}$$

and since

¹ F. Rudio: *Archimedes, Huygens, Lambert, Legendre, vier Abhandlungen über die Kreismessung*. Leipzig, 1892, p. 56f. This error is also reproduced by B. Cald in Enriques's *Fragen der Elementargeometrie*, II. Teil, 1907, p. 315; by D. E. Smith in Young's *Monographs on Topics of Modern Mathematics*, 1911, p. 401. The matter was correctly set forth by T. Vahlen in *Konstruktionen und Approximationen*, Leipzig, 1911, p. 319f.

² A. Pringsheim: "Über die ersten Beweise der Irrationalität von e und π ", Bayerische Akad. der Wissen., *Sitzungsberichte*, mathem.-phys. Cl., v. 28, 1899, p. 325-337.

³ "De fractionibus continuis", *Comment. acad. de Petrop*, v. 9, 1744, p. 108. Presented to St. Petersburg Academy, March. 1737.

⁴ L. Euler: *Introductio in analysin infinitorum*. Tomus Primus, Lausannae, 1748, p. 319. This work was finished in 1745; Cf. G. Eneström, *Verzeichnis* etc., Erste Lieferung, p. 25.

$$\frac{e^x - 1}{e^x + 1} = \frac{e^{x/2} - e^{-x/2}}{e^{x/2} + e^{-x/2}} = \tanh \frac{x}{2} = \frac{1}{i} \tan \frac{ix}{2}, \text{ if } z = \frac{ix}{2},$$

$$\tan z = \frac{1}{1/z} - \frac{1}{3/z} - \frac{1}{5/z} - \frac{1}{7/z} - \frac{1}{9/z} - \dots$$

He then proved the theorems:

1. If x is a rational number different from zero, e^x can never be rational.

For $x = 1$, we have as special case the irrationality of e .

2. If z is a rational number different from zero, $\tan z$ can never be rational.

For $z = \pi/4$, $\tan \pi/4 = 1$, and hence as a special case the irrationality of π .

The part of Lambert's *Vorläufige Kenntnisse* to which Klein refers contains some formulae without proof, and no analytical developments, and was rather intended to serve as a popular survey of the treatment of the topic. With it must be considered the scientifically remarkable "*Mémoire*" of 1767¹. Here "mit minutiöser Genauigkeit" Lambert proves the convergence of the expression for $\tan z$ as a continued fraction. Pringsheim dwells on the "astounding" nature of these considerations at this period in the history of mathematical thought. For of such considerations Legendre was innocent, as well as the great Gauss in his 1812 memoir on hypergeometric series, and others, till a much later period.

"Thus the Lambert memoir contains the *first*, and for many years, the *only* example of what we now consider really rigorous developments of functions as converging continued fractions, in particular, that for $\tan z$ given above."

Measurement of a Circle. By considering inscribed and circumscribed polygons up to 96 sides Archimedes arrived at the result that the ratio of the circumference of a circle to its diameter is less than $3\frac{10}{70}$ but greater than $3\frac{10}{71}$. The following table exhibits the perimeters of regular inscribed and circumscribed polygons of a circle with a unit diameter (Chauvenet, *Treatise on Elementary Geometry*, Philadelphia, 1870, p. 161).

¹ "Mémoire sur quelques propriétés remarquables des quantités transcendentes circulaires et logarithmiques". Lu en 1767. Printed in 1768 in *Hist. de l'acad. royale des sciences et belles-lettres*, Berlin, Année 1761 (1), p. 265—322.

Number of sides	Perimeter of circumscribed polygon	Perimeter of inscribed polygon
4	4.0000000	2.8284271
8	3.3137085	3.0614675
16	3.1825979	3.1214452
32	3.1517249	3.1365485
64	3.1441184	3.1403312
128	3.1422236	3.1412773
256	3.1417504	3.1415138
512	3.1416321	3.1415729
1024	3.1416025	3.1415877
2048	3.1415951	3.1415914
4096	3.1415933	3.1415923
8192	3.1415928	3.1415926

The remarkable approximation $355/113$ for π is correct to six places of decimals. It seems to have been first given by a Chinese, Tsu Ch'ung-ching (5th century), and later by Valentin Otho (16th century) and Adriaen Anthonisz (17th century). Grunert gave a geometrical construction for π based on the fact that $355/113 = 3 + 4^2/(7^2 + 8^2)$, *Archiv der Mathematik und Physik*, v. 12, 1849, p. 98.

Another construction was given by Ramanujan in *Journ. Indian Math. Soc.*, v. 5, 1913, p. 132 (also in *Collected Papers of Srinivasa Ramanujan*, Cambridge, 1927, p. 22, 35).

Euler's Formula. The formula

$$e^{ix} = \cos x + i \sin x$$

was first given by Euler in *Miscellanea Berolinensia*, v. 7, 1743, p. 179 (paper read 6 Sept. 1742), and again in his *Introductio in Analysin*, Lausanne, 1748, v. 1, p. 104. He gave also

$$e^{-ix} = \cos x - i \sin x.$$

The equivalent of the form

$$ix = \log (\cos x + i \sin x$$


was given earlier by Roger Cotes (*Philosophical Transactions*, 1714, v. 29, 1717, p. 32) as: "Si quadrantis circuli quilibet arcus, radio

CE descriptus, sinum habeat CX , sinumque complementi ad quadrantem XE : sumendo radium CE , pro Modulo, arcus erit rationis inter $EX + XC \sqrt{-1}$ & CE mensura ducta in $\sqrt{-1}$." See also Cotes, *Harmonia Mensurarum*, 1722, p. 28.

PART II — CHAPTER IV

In the course of the discussion on pages 61—74 it is assumed that there are an infinite number of prime numbers. One of the neatest proofs of this fact was given by Euclid (about 300 B.C.) in proposition 20, book 9 of his *Elements*.

On page 77, in considering the relation $y = e^x$, Klein made a slight slip when he wrote: "To an algebraic value of x corresponds a transcendental value of y , and conversely." "Conversely" leads to the statement, to a transcendental value of y corresponds an algebraic value of x . But a proof of this has nowhere been given; indeed the result is not true, in general. To correct delete "conversely" and add: "To an algebraic value of y corresponds a transcendental value of x ."



CATALOGUE OF DOVER BOOKS

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

General

WHAT IS SCIENCE?, Norman Campbell. This excellent introduction explains scientific method, role of mathematics, types of scientific laws. Contents: 2 aspects of science, science & nature, laws of science, discovery of laws, explanation of laws, measurement & numerical laws, applications of science. 192pp. 5½ x 8. S43 Paperbound \$1.25

THE COMMON SENSE OF THE EXACT SCIENCES, W. K. Clifford. Introduction by James Newman, edited by Karl Pearson. For 70 years this has been a guide to classical scientific and mathematical thought. Explains with unusual clarity basic concepts, such as extension of meaning of symbols, characteristics of surface boundaries, properties of plane figures, vectors, Cartesian method of determining position, etc. Long preface by Bertrand Russell. Bibliography of Clifford. Corrected, 130 diagrams redrawn. 249pp. 5½ x 8. T61 Paperbound \$1.60

SCIENCE THEORY AND MAN, Erwin Schrödinger. This is a complete and unabridged reissue of **SCIENCE AND THE HUMAN TEMPERAMENT** plus an additional essay: "What is an Elementary Particle?" Nobel laureate Schrödinger discusses such topics as nature of scientific method, the nature of science, chance and determinism, science and society, conceptual models for physical entities, elementary particles and wave mechanics. Presentation is popular and may be followed by most people with little or no scientific training. "Fine practical preparation for a time when laws of nature, human institutions . . . are undergoing a critical examination without parallel," Waldemar Kaempffert, N. Y. TIMES. 192pp. 5½ x 8. T428 Paperbound \$1.35

FADS AND FALLACIES IN THE NAME OF SCIENCE, Martin Gardner. Examines various cults, quack systems, frauds, delusions which at various times have masqueraded as science. Accounts of hollow-earth fanatics like Symmes; Velikovsky and wandering planets; Hoerbiger; Bellamy and the theory of multiple moons; Charles Fort; dowsing, pseudoscientific methods for finding water, ores, oil. Sections on naturopathy, iridagnosis, zone therapy, food fads, etc. Analytical accounts of Wilhelm Reich and orgone sex energy; L. Ron Hubbard and Dianetics; A. Korzybski and General Semantics; many others. Brought up to date to include Bridey Murphy, others. Not just a collection of anecdotes, but a fair, reasoned appraisal of eccentric theory. Formerly titled **IN THE NAME OF SCIENCE**. Preface. Index. x + 384pp. 5½ x 8. T394 Paperbound \$1.50

A DOVER SCIENCE SAMPLER, edited by George Barkin. 64-page book, sturdily bound, containing excerpts from over 20 Dover books, explaining science. Edwin Hubble, George Sarton, Ernst Mach, A. d'Abro, Galileo, Newton, others, discussing island universes, scientific truth, biological phenomena, stability in bridges, etc. Copies limited; no more than 1 to a customer, FREE

POPULAR SCIENTIFIC LECTURES, Hermann von Helmholtz. Helmholtz was a superb expositor as well as a scientist of genius in many areas. The seven essays in this volume are models of clarity, and even today they rank among the best general descriptions of their subjects ever written. "The Physiological Causes of Harmony in Music" was the first significant physiological explanation of musical consonance and dissonance. Two essays, "On the Interaction of Natural Forces" and "On the Conservation of Force," were of great importance in the history of science, for they firmly established the principle of the conservation of energy. Other lectures include "On the Relation of Optics to Painting," "On Recent Progress in the Theory of Vision," "On Goethe's Scientific Researches," and "On the Origin and Significance of Geometrical Axioms." Selected and edited with an introduction by Professor Morris Kline. xii + 286pp. 5½ x 8½. T799 Paperbound \$1.45

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

Physics

CONCERNING THE NATURE OF THINGS, Sir William Bragg. Christmas lectures delivered at the Royal Society by Nobel laureate. Why a spinning ball travels in a curved track; how uranium is transmuted to lead, etc. Partial contents: atoms, gases, liquids, crystals, metals, etc. No scientific background needed; wonderful for intelligent child. 32pp. of photos, 57 figures. xii + 232pp. 5½ x 8. T31 Paperbound \$1.50

THE RESTLESS UNIVERSE, Max Born. New enlarged version of this remarkably readable account by a Nobel laureate. Moving from sub-atomic particles to universe, the author explains in very simple terms the latest theories of wave mechanics. Partial contents: air and its relatives, electrons & ions, waves & particles, electronic structure of the atom, nuclear physics. Nearly 1000 illustrations, including 7 animated sequences. 325pp. 6 x 9. T412 Paperbound \$2.00

Catalogue of Dover Books

FROM EUCLID TO EDDINGTON: A STUDY OF THE CONCEPTIONS OF THE EXTERNAL WORLD, Sir Edmund Whittaker. A foremost British scientist traces the development of theories of natural philosophy from the western rediscovery of Euclid to Eddington, Einstein, Dirac, etc. The inadequacy of classical physics is contrasted with present day attempts to understand the physical world through relativity, non-Euclidean geometry, space curvature, wave mechanics, etc. 5 major divisions of examination: Space; Time and Movement; the Concepts of Classical Physics; the Concepts of Quantum Mechanics; the Eddington Universe. 212pp. 5½ x 8. T491 Paperbound \$1.35

PHYSICS, THE PIONEER SCIENCE, L. W. Taylor. First thorough text to place all important physical phenomena in cultural-historical framework; remains best work of its kind. Exposition of physical laws, theories-developed chronologically, with great historical, illustrative experiments diagrammed, described, worked out mathematically. Excellent physics text for self-study as well as class work. Vol. 1: Heat, Sound; motion, acceleration, gravitation, conservation of energy, heat engines, rotation, heat, mechanical energy, etc. 211 illus. 407pp. 5½ x 8. Vol. 2: Light, Electricity: images, lenses, prisms, magnetism, Ohm's law, dynamos, telegraph, quantum theory, decline of mechanical view of nature, etc. Bibliography. 13 table appendix. Index. 551 illus. 2 color plates. 508pp. 5½ x 8.

Vol. 1 S565 Paperbound \$2.25

Vol. 2 S566 Paperbound \$2.25

The set \$4.50

A SURVEY OF PHYSICAL THEORY, Max Planck. One of the greatest scientists of all time, creator of the quantum revolution in physics, writes in non-technical terms of his own discoveries and those of other outstanding creators of modern physics. Planck wrote this book when science had just crossed the threshold of the new physics, and he communicates the excitement felt then as he discusses electromagnetic theories, statistical methods, evolution of the concept of light, a step-by-step description of how he developed his own momentous theory, and many more of the basic ideas behind modern physics. Formerly "A Survey of Physics." Bibliography. Index. 128pp. 5½ x 8. S650 Paperbound \$1.15

THE ATOMIC NUCLEUS, M. Korsunsky. The only non-technical comprehensive account of the atomic nucleus in English. For college physics students, etc. Chapters cover: Radioactivity, the Nuclear Model of the Atom, the Mass of Atomic Nuclei, the Disintegration of Atomic Nuclei, the Discovery of the Positron, the Artificial Transformation of Atomic Nuclei, Artificial Radioactivity, Mesons, the Neutrino, the Structure of Atomic Nuclei and Forces Acting Between Nuclear Particles, Nuclear Fission, Chain Reaction, Peaceful Uses, Thermocuclear Reactions. Slightly abridged edition. Translated by G. Yankovsky. 65 figures. Appendix includes 45 photographic illustrations. 413 pp. 5½ x 8. S1052 Paperbound \$2.00

PRINCIPLES OF MECHANICS SIMPLY EXPLAINED, Morton Mott-Smith. Excellent, highly readable introduction to the theories and discoveries of classical physics. Ideal for the layman who desires a foundation which will enable him to understand and appreciate contemporary developments in the physical sciences. Discusses: Density, The Law of Gravitation, Mass and Weight, Action and Reaction, Kinetic and Potential Energy, The Law of Inertia, Effects of Acceleration, The Independence of Motions, Galileo and the New Science of Dynamics, Newton and the New Cosmos, The Conservation of Momentum, and other topics. Revised edition of "This Mechanical World." Illustrated by E. Kosa, Jr. Bibliography and Chronology. Index. xiv + 171pp. 5½ x 8½. T1067 Paperbound \$1.35

THE CONCEPT OF ENERGY SIMPLY EXPLAINED, Morton Mott-Smith. Elementary, non-technical exposition which traces the story of man's conquest of energy, with particular emphasis on the developments during the nineteenth century and the first three decades of our own century. Discusses man's earlier efforts to harness energy, more recent experiments and discoveries relating to the steam engine, the engine indicator, the motive power of heat, the principle of excluded perpetual motion, the bases of the conservation of energy, the concept of entropy, the internal combustion engine, mechanical refrigeration, and many other related topics. Also much biographical material. Index. Bibliography. 33 illustrations. ix + 215pp. 5½ x 8½. T1071 Paperbound \$1.25

HEAT AND ITS WORKINGS, Morton Mott-Smith. One of the best elementary introductions to the theory and attributes of heat, covering such matters as the laws governing the effect of heat on solids, liquids and gases, the methods by which heat is measured, the conversion of a substance from one form to another through heating and cooling, evaporation, the effects of pressure on boiling and freezing points, and the three ways in which heat is transmitted (conduction, convection, radiation). Also brief notes on major experiments and discoveries. Concise, but complete, it presents all the essential facts about the subject in readable style. Will give the layman and beginning student a first-rate background in this major topic in physics. Index. Bibliography. 50 illustrations. x + 165pp. 5½ x 8½. T978 Paperbound \$1.15

THE STORY OF ATOMIC THEORY AND ATOMIC ENERGY, J. G. Feinberg. Wider range of facts on physical theory, cultural implications, than any other similar source. Completely non-technical. Begins with first atomic theory, 600 B.C., goes through A-bomb, developments to 1959. Avogadro, Rutherford, Bohr, Einstein, radioactive decay, binding energy, radiation danger, future benefits of nuclear power, dozens of other topics, told in lively, related, informal manner. Particular stress on European atomic research. "Deserves special mention . . . authoritative." Saturday Review. Formerly "The Atom Story." New chapter to 1959. Index. 34 illustrations. 251pp. 5½ x 8. T625 Paperbound \$1.60

Catalogue of Dover Books

THE STRANGE STORY OF THE QUANTUM, AN ACCOUNT FOR THE GENERAL READER OF THE GROWTH OF IDEAS UNDERLYING OUR PRESENT ATOMIC KNOWLEDGE, B. Hoffmann. Presents lucidly and expertly, with barest amount of mathematics, the problems and theories which led to modern quantum physics. Dr. Hoffmann begins with the closing years of the 19th century, when certain trifling discrepancies were noticed, and with illuminating analogies and examples takes you through the brilliant concepts of Planck, Einstein, Pauli, de Broglie, Bohr, Schroedinger, Heisenberg, Dirac, Sommerfeld, Feynman, etc. This edition includes a new, long postscript carrying the story through 1958. "Of the books attempting an account of the history and contents of our modern atomic physics which have come to my attention, this is the best," H. Margenau, Yale University, in "American Journal of Physics." 32 tables and line illustrations. Index. 275pp. 5½ x 8. T518 Paperbound \$1.75

THE EVOLUTION OF SCIENTIFIC THOUGHT FROM NEWTON TO EINSTEIN, A. d'Abro. Einstein's special and general theories of relativity, with their historical implications, are analyzed in non-technical terms. Excellent accounts of the contributions of Newton, Riemann, Weyl, Planck, Eddington, Maxwell, Lorentz and others are treated in terms of space and time, equations of electromagnetics, finiteness of the universe, methodology of science. 21 diagrams. 482pp. 5½ x 8. T2 Paperbound \$2.25

THE RISE OF THE NEW PHYSICS, A. d'Abro. A half-million word exposition, formerly titled **THE DECLINE OF MECHANISM**, for readers not versed in higher mathematics. The only thorough explanation, in everyday language, of the central core of modern mathematical physical theory, treating both classical and modern theoretical physics, and presenting in terms almost anyone can understand the equivalent of 5 years of study of mathematical physics. Scientifically impeccable coverage of mathematical-physical thought from the Newtonian system up through the electronic theories of Dirac and Heisenberg and Fermi's statistics. Combines both history and exposition; provides a broad yet unified and detailed view, with constant comparison of classical and modern views on phenomena and theories. "A must for anyone doing serious study in the physical sciences," JOURNAL OF THE FRANKLIN INSTITUTE. "Extraordinary faculty . . . to explain ideas and theories of theoretical physics in the language of daily life," ISIS. First part of set covers philosophy of science, drawing upon the practice of Newton, Maxwell, Poincaré, Einstein, others, discussing modes of thought, experiment, interpretations of causality, etc. In the second part, 100 pages explain grammar and vocabulary of mathematics, with discussions of functions, groups, series, Fourier series, etc. The remainder is devoted to concrete, detailed coverage of both classical and quantum physics, explaining such topics as analytic mechanics, Hamilton's principle, wave theory of light, electromagnetic waves, groups of transformations, thermodynamics, phase rule, Brownian movement, kinetics, special relativity, Planck's original quantum theory, Bohr's atom, Zeeman effect, Broglie's wave mechanics, Heisenberg's uncertainty, Eigen-values, matrices, scores of other important topics. Discoveries and theories are covered for such men as Alembert, Born, Cantor, Debye, Euler, Foucault, Galois, Gauss, Hadamard, Kelvin, Kepler, Laplace, Maxwell, Pauli, Rayleigh, Volterra, Weyl, Young, more than 180 others. Indexed. 97 illustrations. ix + 982pp. 5½ x 8. T3 Volume 1, Paperbound \$2.25
T4 Volume 2, Paperbound \$2.25

SPINNING TOPS AND GYROSCOPIC MOTION, John Perry. Well-known classic of science still unsurpassed for lucid, accurate, delightful exposition. How quasi-rigidity is induced in flexible and fluid bodies by rapid motions; why gyrostap falls, top rises; nature and effect on climatic conditions of earth's precessional movement; effect of internal fluidity on rotating bodies, etc. Appendixes describe practical uses to which gyroscopes have been put in ships, compasses, monorail transportation. 62 figures. 128pp. 5½ x 8. T416 Paperbound \$1.25

THE UNIVERSE OF LIGHT, Sir William Bragg. No scientific training needed to read Nobel Prize winner's expansion of his Royal Institute Christmas Lectures. Insight into nature of light, methods and philosophy of science. Explains lenses, reflection, color, resonance, polarization, x-rays, the spectrum, Newton's work with prisms, Huygens' with polarization, Crookes' with cathode ray, etc. Leads into clear statement of 2 major historical theories of light, corpuscle and wave. Dozens of experiments you can do. 199 illus., including 2 full-page color plates. 293pp. 5½ x 8. S538 Paperbound \$1.85

THE STORY OF X-RAYS FROM RÖNTGEN TO ISOTOPES, A. R. Bleich. Non-technical history of x-rays, their scientific explanation, their applications in medicine, industry, research, and art, and their effect on the individual and his descendants. Includes amusing early reactions to Röntgen's discovery, cancer therapy, detections of art and stamp forgeries, potential risks to patient and operator, etc. Illustrations show x-rays of flower structure, the gall bladder, gears with hidden defects, etc. Original Dover publication. Glossary. Bibliography. Index. 55 photos and figures. xiv + 186pp. 5½ x 8. T662 Paperbound \$1.50

ELECTRONS, ATOMS, METALS AND ALLOYS, Wm. Hume-Rothery. An introductory-level explanation of the application of the electronic theory to the structure and properties of metals and alloys, taking into account the new theoretical work done by mathematical physicists. Material presented in dialogue-form between an "Old Metallurgist" and a "Young Scientist." Their discussion falls into 4 main parts: the nature of an atom, the nature of a metal, the nature of an alloy, and the structure of the nucleus. They cover such topics as the hydrogen atom, electron waves, wave mechanics, Brillouin zones, co-valent bonds, radioactivity and natural disintegration, fundamental particles, structure and fission of the nucleus, etc. Revised, enlarged edition. 177 illustrations. Subject and name indexes. 407pp. 5½ x 8½. S1046 Paperbound \$2.25

Catalogue of Dover Books

TEACH YOURSELF MECHANICS, P. Abbott. The lever, centre of gravity, parallelogram of force, friction, acceleration, Newton's laws of motion, machines, specific gravity, gas, liquid pressure, much more. 280 problems, solutions. Tables. 163 illus. 271pp. 6 $\frac{3}{4}$ x 4 $\frac{1}{4}$. Clothbound \$2.00

MATTER & MOTION, James Clerk Maxwell. This excellent exposition begins with simple particles and proceeds gradually to physical systems beyond complete analysis: motion, force, properties of centre of mass of material system, work, energy, gravitation, etc. Written with all Maxwell's original insights and clarity. Notes by E. Larmor. 17 diagrams. 178pp. 5 $\frac{3}{8}$ x 8. \$188 Paperbound \$1.35

SOAP BUBBLES, THEIR COLOURS AND THE FORCES WHICH MOULD THEM, C. V. Boys. Only complete edition, half again as much material as any other. Includes Boys' hints on performing his experiments, sources of supply. Dozens of lucid experiments show complexities of liquid films, surface tension, etc. Best treatment ever written. Introduction. 83 illustrations. Color plate. 202pp. 5 $\frac{3}{8}$ x 8. T542 Paperbound 95¢

MATTER & LIGHT, THE NEW PHYSICS, L. de Broglie. Non-technical papers by a Nobel laureate explain electromagnetic theory, relativity, matter, light and radiation, wave mechanics, quantum physics, philosophy of science. Einstein, Planck, Bohr, others explained so easily that no mathematical training is needed for all but 2 of the 21 chapters. Unabridged. Index. 300pp. 5 $\frac{3}{8}$ x 8. T35 Paperbound \$1.85

SPACE AND TIME, Emile Borel. An entirely non-technical introduction to relativity, by world-renowned mathematician, Sorbonne professor. (Notes on basic mathematics are included separately.) This book has never been surpassed for insight, and extraordinary clarity of thought, as it presents scores of examples, analogies, arguments, illustrations, which explain such topics as: difficulties due to motion; gravitation a force of Inertia; geodesic lines; wave-length and difference of phase; x-rays and crystal structure; the special theory of relativity; and much more. Indexes. 4 appendixes. 15 figures. xvi + 243pp. 5 $\frac{3}{8}$ x 8. T592 Paperbound \$1.45

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

Astronomy

THE FRIENDLY STARS, Martha Evans Martin. This engaging survey of stellar lore and science is a well-known classic, which has introduced thousands to the fascinating world of stars and other celestial bodies. Descriptions of Capella, Sirius, Arcturus, Vega, Polaris, etc.—all the important stars, with informative discussions of rising and setting of stars, their number, names, brightness, distances, etc. in a non-technical, highly readable style. Also: double stars, constellations, clusters—concentrating on stars and formations visible to the naked eye. New edition, revised (1963) by D. H. Menzel, Director Harvard Observatory. 23 diagrams by Prof. Ching-Sung Yu. Foreword by D. H. Menzel and W. W. Morgan. 2 Star Charts. Index. xii + 147pp. 5 $\frac{3}{8}$ x 8 $\frac{1}{2}$. T1099 Paperbound \$1.00

AN ELEMENTARY SURVEY OF CELESTIAL MECHANICS, Y. Ryabov. Elementary exposition of gravitational theory and celestial mechanics. Historical introduction and coverage of basic principles, including: the elliptic, the orbital plane, the 2- and 3-body problems, the discovery of Neptune, planetary rotation, the length of the day, the shapes of galaxies, satellites (detailed treatment of Sputnik I), etc. First American reprinting of successful Russian popular exposition. Elementary algebra and trigonometry helpful, but not necessary; presentation chiefly verbal. Appendix of theorem proofs. 58 figures. 165pp. 5 $\frac{3}{8}$ x 8. T756 Paperbound \$1.25

THE SKY AND ITS MYSTERIES, E. A. Beet. One of most lucid books on mysteries of universe; deals with astronomy from earliest observations to latest theories of expansion of universe, source of stellar energy, birth of planets, origin of moon craters, possibility of life on other planets. Discusses effects of sunspots on weather; distances, ages of several stars; master plan of universe; methods and tools of astronomers; much more. "Eminently readable book," London Times. Extensive bibliography. Over 50 diagrams. 12 full-page plates, fold-out star map. Introduction. Index. 5 $\frac{1}{4}$ x 7 $\frac{1}{2}$. T627 Clothbound \$3.50

THE REALM OF THE NEBULAE, E. Hubble. One of the great astronomers of our time records his formulation of the concept of "island universes," and its impact on astronomy. Such topics are covered as the velocity-distance relation; classification, nature, distances, general field of nebulae; cosmological theories; nebulae in the neighborhood of the Milky Way. 39 photos of nebulae, nebulae clusters, spectra of nebulae, and velocity distance relations shown by spectrum comparison. "One of the most progressive lines of astronomical research," The Times (London). New introduction by A. Sandage. 55 illustrations. Index. iv + 201pp. 5 $\frac{3}{8}$ x 8. S455 Paperbound \$1.50

Catalogue of Dover Books

OUT OF THE SKY, H. H. Nininger. A non-technical but comprehensive introduction to "meteoritics", the young science concerned with all aspects of the arrival of matter from outer space. Written by one of the world's experts on meteorites, this work shows how, despite difficulties of observation and sparseness of data, a considerable body of knowledge has arisen. It defines meteors and meteorites; studies fireball clusters and processions, meteorite composition, size, distribution, showers, explosions, origins, craters, and much more. A true connecting link between astronomy and geology. More than 175 photos, 22 other illustrations. References. Bibliography of author's publications on meteorites. Index. viii + 336pp. 5 3/8 x 8. T519 Paperbound \$1.85

SATELLITES AND SCIENTIFIC RESEARCH, D. King-Hele. Non-technical account of the manmade satellites and the discoveries they have yielded up to the autumn of 1961. Brings together information hitherto published only in hard-to-get scientific journals. Includes the life history of a typical satellite, methods of tracking, new information on the shape of the earth, zones of radiation, etc. Over 60 diagrams and 6 photographs. Mathematical appendix. Bibliography of over 100 items. Index. xii + 180pp. 5 3/8 x 8 1/2. T703 Paperbound \$2.00

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

Mathematics

CHANCE, LUCK AND STATISTICS: THE SCIENCE OF CHANCE, Horace C. Levinson. Theory of probability and science of statistics in simple, non-technical language. Part I deals with theory of probability, covering odd superstitions in regard to "luck," the meaning of betting odds, the law of mathematical expectation, gambling, and applications in poker, roulette, lotteries, dice, bridge, and other games of chance. Part II discusses the misuse of statistics, the concept of statistical probabilities, normal and skew frequency distributions, and statistics applied to various fields—birth rates, stock speculation, insurance rates, advertising, etc. "Presented in an easy humorous style which I consider the best kind of expository writing," Prof. A. C. Cohen, Industry Quality Control. Enlarged revised edition. Formerly titled "The Science of Chance." Preface and two new appendices by the author. Index. xiv + 365pp. 5 3/8 x 8. T1007 Paperbound \$1.85

PROBABILITIES AND LIFE, Emile Borel. Translated by M. Baudin. Non-technical, highly readable introduction to the results of probability as applied to everyday situations. Partial contents: Fallacies About Probabilities Concerning Life After Death; Negligible Probabilities and the Probabilities of Everyday Life; Events of Small Probability; Application of Probabilities to Certain Problems of Heredity; Probabilities of Deaths, Diseases, and Accidents; On Poisson's Formula. Index. 3 Appendices of statistical studies and tables. vi + 87pp. 5 3/8 x 8 1/2. T121 Paperbound \$1.00

GREAT IDEAS OF MODERN MATHEMATICS: THEIR NATURE AND USE, Jagjit Singh. Reader with only high school math will understand main mathematical ideas of modern physics, astronomy, genetics, psychology, evolution, etc., better than many who use them as tools, but comprehend little of their basic structure. Author uses his wide knowledge of non-mathematical fields in brilliant exposition of differential equations, matrices, group theory, logic, statistics, problems of mathematical foundations, imaginary numbers, vectors, etc. Original publication. 2 appendices. 2 indexes. 65 illustr. 322pp. 5 3/8 x 8. S587 Paperbound \$1.75

MATHEMATICS IN ACTION, O. G. Sutton. Everyone with a command of high school algebra will find this book one of the finest possible introductions to the application of mathematics to physical theory. Ballistics, numerical analysis, waves and wavelike phenomena, Fourier series, group concepts, fluid flow and aerodynamics, statistical measures, and meteorology are discussed with unusual clarity. Some calculus and differential equations theory is developed by the author for the reader's help in the more difficult sections. 88 figures. Index. viii + 236pp. 5 3/8 x 8. T440 Clothbound \$3.50

THE FOURTH DIMENSION SIMPLY EXPLAINED, edited by H. P. Manning. 22 essays, originally Scientific American contest entries, that use a minimum of mathematics to explain aspects of 4-dimensional geometry: analogues to 3-dimensional space, 4-dimensional absurdities and curiosities (such as removing the contents of an egg without puncturing its shell), possible measurements and forms, etc. Introduction by the editor. Only book of its sort on a truly elementary level, excellent introduction to advanced works. 82 figures. 251pp. 5 3/8 x 8. T711 Paperbound \$1.35

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

Engineering, technology, applied science etc.

TEACH YOURSELF ELECTRICITY, C. W. Wilman. Electrical resistance, inductance, capacitance, magnets, chemical effects of current, alternating currents, generators and motors, transformers, rectifiers, much more. 230 questions, answers, worked examples. List of units. 115 illus. 194pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$. Clothbound \$2.00

ELEMENTARY METALLURGY AND METALLOGRAPHY, A. M. Shrager. Basic theory and descriptions of most of the fundamental manufacturing processes involved in metallurgy. Partial contents: the structure of metals; slip, plastic deformation, and recrystallization; iron ore and production of pig iron; chemistry involved in the metallurgy of iron and steel; basic processes such as the Bessemer treatment, open-hearth process, the electric arc furnace—with advantages and disadvantages of each; annealing, hardening, and tempering steel; copper, aluminum, magnesium, and their alloys. For freshman engineers, advanced students in technical high schools, etc. Index. Bibliography. 177 diagrams. 17 tables. 284 questions and problems. 27-page glossary. ix + 389pp. 5 $\frac{3}{8}$ x 8. S138 Paperbound \$2.25

BASIC ELECTRICITY, Prepared by the Bureau of Naval Personnel. Originally a training course text for U.S. Navy personnel, this book provides thorough coverage of the basic theory of electricity and its applications. Best book of its kind for either broad or more limited studies of electrical fundamentals . . . for classroom use or home study. Part 1 provides a more limited coverage of theory: fundamental concepts, batteries, the simple circuit, D.C. series and parallel circuits, conductors and wiring techniques, A.C. electricity, inductance and capacitance, etc. Part 2 applies theory to the structure of electrical machines—generators, motors, transformers, magnetic amplifiers. Also deals with more complicated instruments, synchros, servo-mechanisms. The concluding chapters cover electrical drawings and blueprints, wiring diagrams, technical manuals, and safety education. The book contains numerous questions for the student, with answers. Index and six appendices. 345 illustrations. x + 448pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$. S973 Paperbound \$3.00

BASIC ELECTRONICS, prepared by the U.S. Navy Training Publications Center. A thorough and comprehensive manual on the fundamentals of electronics. Written clearly, it is equally useful for self-study or course work for those with a knowledge of the principles of basic electricity. Partial contents: Operating Principles of the Electron Tube; Introduction to Transistors; Power Supplies for Electronic Equipment; Tuned Circuits; Electron-Tube Amplifiers; Audio Power Amplifiers; Oscillators; Transmitters; Transmission Lines; Antennas and Propagation; Introduction to Computers; and related topics. Appendix. Index. Hundreds of illustrations and diagrams. vi + 471pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$. S1076 Paperbound \$2.75

BASIC THEORY AND APPLICATION OF TRANSISTORS, Prepared by the U.S. Department of the Army. An introductory manual prepared for an army training program. One of the finest available surveys of theory and application of transistor design and operation. Minimal knowledge of physics and theory of electron tubes required. Suitable for textbook use, course supplement, or home study. Chapters: Introduction; fundamental theory of transistors; transistor amplifier fundamentals; parameters, equivalent circuits, and characteristic curves; bias stabilization; transistor analysis and comparison using characteristic curves and charts; audio amplifiers; tuned amplifiers; wide-band amplifiers; oscillators; pulse and switching circuits; modulation, mixing, and demodulation; and additional semiconductor devices. Unabridged, corrected edition. 240 schematic drawings, photographs, wiring diagrams, etc. 2 Appendices. Glossary. Index. 263pp. 6 $\frac{1}{2}$ x 9 $\frac{1}{4}$. S380 Paperbound \$1.25

TEACH YOURSELF HEAT ENGINES, E. De Ville. Measurement of heat, development of steam and internal combustion engines, efficiency of an engine, compression-ignition engines, production of steam, the ideal engine, much more. 318 exercises, answers, worked examples. Tables. 76 illus. 220pp. 6 $\frac{7}{8}$ x 4 $\frac{1}{4}$. Clothbound \$2.00

BOOKS EXPLAINING SCIENCE AND MATHEMATICS

Miscellaneous

ON THE SENSATIONS OF TONE, Hermann Helmholtz. This is an unmatched coordination of such fields as acoustical physics, physiology, experiment, history of music. It covers the entire gamut of musical tone. Partial contents: relation of musical science to acoustics, physical vs. physiological acoustics, composition of vibration, resonance, analysis of tones by sympathetic resonance, beats, chords, tonality, consonant chords, discords, progression of parts, etc. 33 appendixes discuss various aspects of sound, physics, acoustics, music, etc. Translated by A. J. Ellis. New introduction by Prof. Henry Margenau of Yale. 68 figures. 43 musical passages analyzed. Over 100 tables. Index. xix + 576pp. 6 $\frac{3}{8}$ x 9 $\frac{1}{4}$. S114 Paperbound \$3.00

Catalogue of Dover Books

THE NATURE OF LIGHT AND COLOUR IN THE OPEN AIR, M. Minnaert. Why is falling snow sometimes black? What causes mirages, the fata morgana, multiple suns and moons in the sky? How are shadows formed? Prof. Minnaert of the University of Utrecht answers these and similar questions in optics, light, colour, for non-specialists. Particularly valuable to nature, science students, painters, photographers. Translated by H. M. Kremer-Priest, K. Jay. 202 illustrations, including 42 photos. xvi + 362pp. 5½ x 8. T196 Paperbound \$2.00

THE PHYSICS OF MUSIC, Alexander Wood. Introduction for musicians to the physical aspect of sound. No scientific training necessary to understand concepts, etc. Wealth of material on origin and development of instruments, physical principles involved in the production of their sounds, pitch, intensity and loudness, mechanism of the ear, dissonance and consonance, sound reproduction and recordings, concert halls, etc. Extensively revised by Dr. J. M. Bowscher. Indices. Bibliography. 16 plates. 114 illustrations. 270pp. 5½ x 8½. T322 Paperbound \$2.25

GREAT IDEAS AND THEORIES OF MODERN COSMOLOGY, Jagjit Singh. The theories of Jeans, Eddington, Milne, Kant, Bondi, Gold, Newton, Einstein, Gamow, Hoyle, Dirac, Kuiper, Hubble, Weizsäcker and many others on such cosmological questions as the origin of the universe, space and time, planet formation, "continuous creation," the birth, life, and death of the stars, the origin of the galaxies, etc. By the author of the popular "Great Ideas of Modern Mathematics." A gifted popularizer of science, he makes the most difficult abstractions crystal-clear even to the most non-mathematical reader. Index. xii + 276 pp. 5½ x 8½. T925 Paperbound \$1.85

PIONEERS OF SCIENCE, O. Lodge. Eminent scientist-expositor's authoritative, yet elementary survey of great scientific theories. Concentrating on individuals—Copernicus, Brahe, Kepler, Galileo, Descartes, Newton, Laplace, Herschel, Lord Kelvin, and other scientists—the author presents their discoveries in historical order adding biographical material on each man and full, specific explanations of their achievements. The clear and complete treatment of the post-Newtonian astronomers is a feature seldom found in other books on the subject. Index. 120 illustrations. xv + 404pp. 5½ x 8. T716 Paperbound \$1.65

BIOGRAPHY OF SCIENTISTS

ISAAC NEWTON: A BIOGRAPHY, Louis Trenchard More. The definitive biography of Newton, his life and work. Presents Newton as a living man, with a critical, objective analysis of his character as well as a careful survey of his manifold accomplishments, scientific, theological, etc. The author, himself a professor of physics, has made full use of all of Newton's published works and all material in the Portsmouth Collection of Newton's personal and unpublished papers. The text includes numerous letters by Newton and his acquaintances, and many other of his papers—some translated from Latin to English by the author. A universally-esteemed work. Unabridged republication. 1 full-page plate. Index. xiii + 675pp. 5½ x 8½. T579 Paperbound \$2.50

PIERRE CURIE, Marie Curie. Mme. Curie, Nobel Prize winner, creates a memorable portrait of her equally famous husband and his lifelong scientific researches. She brings to life the determined personality of a great scientist at work. Her own autobiographical notes, included in this volume, reconstruct her own work on radiation which resulted in the isolation of radium. "A delightful book. It marks one of the few instances in which the proverbially humdrum life of the student of physical science, together with the austere ideals, has been made intelligible," New York Times. Unabridged reprint. Translated by Charlotte and Vernon Kellogg. Introduction by Mrs. Wm. Brown Meloney. 8 halftones. viii + 120pp. 5½ x 8½. T199 Paperbound \$1.00

THE BOOK OF MY LIFE (DE VITA PROPRIA LIBER), Jerome Cardan. The remarkable autobiography of an important Renaissance mathematician, physician, and scientist, who at the same time was a paranoid, morbid, superstitious man, consumed with ambition and self-love (and self-pity). These chronicles of his fortunes and misfortunes make absorbing reading, giving us an extremely insightful view of a man's reactions and sensations—the first psychological autobiography. Through his eyes we can also see the superstitions and beliefs of an age, Renaissance medical practices, and the problems that concerned a trained mind in the 16th century. Unabridged republication of original English edition, translated by Jean Stoner. Introduction. Notes. Bibliography. xviii + 331pp. 5½ x 8½. T345 Paperbound \$1.60

THE AUTOBIOGRAPHY OF CHARLES DARWIN, AND SELECTED LETTERS, edited by Francis Darwin. Darwin's own record of his early life; the historic voyage aboard the "Beagle"; the furor surrounding evolution, and his replies; reminiscences of his son. Letters to Henslow, Lyell, Hooker, Huxley, Wallace, Kingsley, etc., and thoughts on religion and vivisection. We see how he revolutionized geology with his concept of ocean subsidence; how his great books on variation of plants and animals, primitive man, the expression of emotion among primates, plant fertilization, carnivorous plants, protective coloration, etc., came into being. Appendix. Index. 365pp. 5½ x 8. T479 Paperbound \$2.00

MATHEMATICAL PUZZLES AND RECREATIONS

AMUSEMENTS IN MATHEMATICS, Henry Ernest Dudeney. The foremost British originator of mathematical puzzles is always intriguing, witty, and paradoxical in this classic, one of the largest collections of mathematical amusements. More than 430 puzzles, problems, and paradoxes. Mazes and games, problems on number manipulation, unicursal and other route problems, puzzles on measuring, weighing, packing, age, kinship, chessboards, joining, crossing river, plane figure dissection, and many others. Solutions. More than 450 illustrations. vii + 258pp. 5½ x 8. T473 Paperbound \$1.25

SYMBOLIC LOGIC and THE GAME OF LOGIC, Lewis Carroll. "Symbolic Logic" is not concerned with modern symbolic logic, but is instead a collection of over 380 problems posed with charm and imagination, using the syllogism, and a fascinating diagrammatic method of drawing conclusions. In "The Game of Logic," Carroll's whimsical imagination devises a logical game played with 2 diagrams and counters (included) to manipulate hundreds of tricky syllogisms. The final section, "Hit or Miss" is a lagniappe of 101 additional puzzles in the delightful Carroll manner. Until this reprint edition, both of these books were rarities costing up to \$15 each. Symbolic Logic: Index, xxxi + 199pp. The Game of Logic: 96pp. Two vols. bound as one. 5½ x 8. T492 Paperbound \$1.50

MAZES AND LABYRINTHS: A BOOK OF PUZZLES, W. Shepherd. Mazes, formerly associated with mystery and ritual, are still among the most intriguing of intellectual puzzles. This is a novel and different collection of 50 amusements that embody the principle of the maze: mazes in the classical tradition; 3-dimensional, ribbon, and Möbius-strip mazes; hidden messages; spatial arrangements; etc.—almost all built on amusing story situations. 84 illustrations. Essay on maze psychology. Solutions. xv + 122pp. 5½ x 8. T731 Paperbound \$1.00

MATHEMATICAL RECREATIONS, M. Kraitchik. Some 250 puzzles, problems, demonstrations of recreational mathematics for beginners & advanced mathematicians. Unusual historical problems from Greek, Medieval, Arabic, Hindu sources: modern problems based on "mathematics without numbers," geometry, topology, arithmetic, etc. Pastimes derived from figurative numbers, Mersenne numbers, Fermat numbers; fairy chess, latruncles, reversi, many topics. Full solutions. Excellent for insights into special fields of math. 181 illustrations. 330pp. 5½ x 8. T163 Paperbound \$1.75

MATHEMATICAL PUZZLES OF SAM LOYD, Vol. I, selected and edited by M. Gardner. Puzzles by the greatest puzzle creator and innovator. Selected from his famous "Cyclopedia of Puzzles," they retain the unique style and historical flavor of the originals. There are posers based on arithmetic, algebra, probability, game theory, route tracing, topology, counter, sliding block, operations research, geometrical dissection. Includes his famous "14-15" puzzle which was a national craze, and his "Horse of a Different Color" which sold millions of copies. 117 of his most ingenious puzzles in all, 120 line drawings and diagrams. Solutions. Selected references. xx + 167pp. 5½ x 8. T498 Paperbound \$1.00

MY BEST PUZZLES IN MATHEMATICS, Hubert Phillips ("Caliban"). Caliban is generally considered the best of the modern problemists. Here are 100 of his best and wittiest puzzles, selected by the author himself from such publications as the London Daily Telegraph, and each puzzle is guaranteed to put even the sharpest puzzle detective through his paces. Perfect for the development of clear thinking and a logical mind. Complete solutions are provided for every puzzle. x + 107pp. 5½ x 8½. T91 Paperbound \$1.00

MY BEST PUZZLES IN LOGIC AND REASONING, H. Phillips ("Caliban"). 100 choice, hitherto unavailable puzzles by England's best-known problemist. No special knowledge needed to solve these logical or inferential problems, just an unclouded mind, nerves of steel, and fast reflexes. Data presented are both necessary and just sufficient to allow one unambiguous answer. More than 30 different types of puzzles, all ingenious and varied, many one of a kind, that will challenge the expert, please the beginner. Original publication. 100 puzzles, full solutions. x + 107pp. 5½ x 8½. T119 Paperbound \$1.00

MATHEMATICAL PUZZLES FOR BEGINNERS AND ENTHUSIASTS, G. Mott-Smith. 188 mathematical puzzles to test mental agility. Inference, interpretation, algebra, dissection of plane figures, geometry, properties of numbers, decimation, permutations, probability, all enter these delightful problems. Puzzles like the Odic Force, How to Draw an Ellipse, Spider's Cousin, more than 180 others. Detailed solutions. Appendix with square roots, triangular numbers, primes, etc. 135 illustrations. 2nd revised edition. 248pp. 5½ x 8. T198 Paperbound \$1.00

MATHEMATICS, MAGIC AND MYSTERY, Martin Gardner. Card tricks, feats of mental mathematics, stage mind-reading, other "magic" explained as applications of probability, sets, theory of numbers, topology, various branches of mathematics. Creative examination of laws and their applications with scores of new tricks and insights. 115 sections discuss tricks with cards, dice, coins; geometrical vanishing tricks, dozens of others. No sleight of hand needed; mathematics guarantees success. 115 illustrations. xii + 174pp. 5½ x 8. T335 Paperbound \$1.00

Catalogue of Dover Books

RECREATIONS IN THE THEORY OF NUMBERS: THE QUEEN OF MATHEMATICS ENTERTAINS, Albert H. Beiler. The theory of numbers is often referred to as the "Queen of Mathematics." In this book Mr. Beiler has compiled the first English volume to deal exclusively with the recreational aspects of number theory, an inherently recreational branch of mathematics. The author's clear style makes for enjoyable reading as he deals with such topics as: perfect numbers, amicable numbers, Fermat's theorem, Wilson's theorem, interesting properties of digits, methods of factoring, primitive roots, Euler's function, polygonal and figurate numbers, Mersenne numbers, congruence, repeating decimals, etc. Countless puzzle problems, with full answers and explanations. For mathematicians and mathematically-inclined laymen, etc. New publication. 28 figures. 9 illustrations. 103 tables. Bibliography at chapter ends. vi + 247pp. 5½ x 8½. T1096 Paperbound \$1.85

PAPER FOLDING FOR BEGINNERS, W. D. Murray and F. J. Rigney. A delightful introduction to the varied and entertaining Japanese art of origami (paper folding), with a full crystal-clear text that anticipates every difficulty; over 275 clearly labeled diagrams of all important stages in creation. You get results at each stage, since complex figures are logically developed from simpler ones. 43 different pieces are explained: place mats, drinking cups, bonbon boxes, sailboats, frogs, roosters, etc. 6 photographic plates. 279 diagrams. 95pp. 5½ x 8½. T713 Paperbound \$1.00

1800 RIDDLES, ENIGMAS AND CONUNDRUMS, Darwin A. Hindman. Entertaining collection ranging from hilarious gags to outrageous puns to sheer nonsense—a welcome respite from sophisticated humor. Children, toastmasters, and practically anyone with a funny bone will find these zany riddles tickling and eminently repeatable. Sample: "Why does Santa Claus always go down the chimney?" "Because it soots him." Some old, some new—covering a wide variety of subjects. New publication. iii + 154pp. 5½ x 8½. T1059 Paperbound \$1.00

EASY-TO-DO ENTERTAINMENTS AND DIVERSIONS WITH CARDS, STRING, COINS, PAPER AND MATCHES, R. M. Abraham. Over 300 entertaining games, tricks, puzzles, and pastimes for children and adults. Invaluable to anyone in charge of groups of youngsters, for party givers, etc. Contains sections on card tricks and games, making things by paperfolding—toys, decorations, and the like; tricks with coins, matches, and pieces of string; descriptions of games; toys that can be made from common household objects; mathematical recreations; word games; and 50 miscellaneous entertainments. Formerly "Winter Nights Entertainments." Introduction by Lord Baden Powell. 329 illustrations. v + 186pp. 5½ x 8. T921 Paperbound \$1.00

DIVERSIONS AND PASTIMES WITH CARDS, STRING, PAPER AND MATCHES, R. M. Abraham. Another collection of amusements and diversion for game and puzzle fans of all ages. Many new paperfolding ideas and tricks, an extensive section on amusements with knots and splices, two chapters of easy and not-so-easy problems, coin and match tricks, and lots of other parlor pastimes from the agile mind of the late British problemist and gamester. Corrected and revised version. Illustrations. 160pp. 5½ x 8½. T1127 Paperbound \$1.00

STRING FIGURES AND HOW TO MAKE THEM: A STUDY OF CAT'S-CRADLE IN MANY LANDS, Caroline Furness Jayne. In a simple and easy-to-follow manner, this book describes how to make 107 different string figures. Not only is looping and crossing string between the fingers a common youthful diversion, but it is an ancient form of amusement practiced in all parts of the globe, especially popular among primitive tribes. These games are fun for all ages and offer an excellent means for developing manual dexterity and coordination. Much insight also for the anthropological observer on games and diversions in many different cultures. Index. Bibliography. Introduction by A. C. Haddon, Cambridge University. 17 full-page plates. 950 illustrations. xxiii + 407pp. 5½ x 8½. T152 Paperbound \$2.00

CRYPTANALYSIS, Helen F. Gaines. (Formerly ELEMENTARY CRYPTANALYSIS.) A standard elementary and intermediate text for serious students. It does not confine itself to old material, but contains much that is not generally known, except to experts. Concealment, Transposition, Substitution ciphers; Vigenere, Kasiski, Playfair, multafid, dozens of other techniques. Appendix with sequence charts, letter frequencies in English, 5 other languages, English word frequencies. Bibliography. 167 codes. New to this edition: solution to codes. vi + 230pp. 5½ x 8. T97 Paperbound \$2.00

MAGIC SQUARES AND CUBES, W. S. Andrews. Only book-length treatment in English, a thorough non-technical description and analysis. Here are nasik, overlapping, pandiagonal, serated squares; magic circles, cubes, spheres, rhombuses. Try your hand at 4-dimensional magical figures! Much unusual folklore and tradition included. High school algebra is sufficient. 754 diagrams and illustrations. viii + 419pp. 5½ x 8. T658 Paperbound \$1.85

CALIBAN'S PROBLEM BOOK: MATHEMATICAL, INFERENTIAL, AND CRYPTOGRAPHIC PUZZLES, H. Phillips ("Caliban"), S. T. Shovelton, G. S. Marshall. 105 ingenious problems by the greatest living creator of puzzles based on logic and inference. Rigorous, modern, piquant, and reflecting their author's unusual personality, these intermediate and advanced puzzles all involve the ability to reason clearly through complex situations; some call for mathematical knowledge, ranging from algebra to number theory. Solutions. xi + 180pp. 5½ x 8. T736 Paperbound \$1.25

FICTION

THE LAND THAT TIME FORGOT and THE MOON MAID, Edgar Rice Burroughs. In the opinion of many, Burroughs' best work. The first concerns a strange island where evolution is individual rather than phylogenetic. Speechless anthropoids develop into intelligent human beings within a single generation. The second projects the reader far into the future and describes the first voyage to the Moon (in the year 2025), the conquest of the Earth by the Moon, and years of violence and adventure as the enslaved Earthmen try to regain possession of their planet. "An imaginative tour de force that keeps the reader keyed up and expectant," *NEW YORK TIMES*. Complete, unabridged text of the original two novels (three parts in each). 5 illustrations by J. Allen St. John. vi + 552pp. 5¾ x 8½.

T1020 Clothbound \$3.75
T358 Paperbound \$2.00

AT THE EARTH'S CORE, PELLUCIDAR, TANAR OF PELLUCIDAR: THREE SCIENCE FICTION NOVELS BY EDGAR RICE BURROUGHS. Complete, unabridged texts of the first three Pellucidar novels. Tales of derring-do by the famous master of science fiction. The locale for these three related stories is the inner surface of the hollow Earth where we discover the world of Pellucidar, complete with all types of bizarre, menacing creatures, strange peoples, and alluring maidens—guaranteed to delight all Burroughs fans and a wide circle of adventure lovers. Illustrated by J. Allen St. John and P. F. Berdanier. vi + 433pp. 5¾ x 8½.

T1051 Paperbound \$2.00

THE PIRATES OF VENUS and LOST ON VENUS: TWO VENUS NOVELS BY EDGAR RICE BURROUGHS. Two related novels, complete and unabridged. Exciting adventure on the planet Venus with Earthman Carson Napier broken-field running through one dangerous episode after another. All lovers of swashbuckling science fiction will enjoy these two stories set in a world of fascinating societies, fierce beasts, 5000-ft. trees, lush vegetation, and wide seas. Illustrations by Fortunino Matania. Total of vi + 340pp. 5¾ x 8½.

T1053 Paperbound \$1.75

A PRINCESS OF MARS and A FIGHTING MAN OF MARS: TWO MARTIAN NOVELS BY EDGAR RICE BURROUGHS. "Princess of Mars" is the very first of the great Martian novels written by Burroughs, and it is probably the best of them all; it set the pattern for all of his later fantasy novels and contains a thrilling cast of strange peoples and creatures and the formula of Olympian heroism amidst ever-fluctuating fortunes which Burroughs carries off so successfully. "Fighting Man" returns to the same scenes and cities—many years later. A mad scientist, a degenerate dictator, and an indomitable defender of the right clash—with the fate of the Red Planet at stake! Complete, unabridged reprinting of original editions. Illustrations by F. E. Schoonover and Hugh Hutton. v + 356pp. 5¾ x 8½.

T1140 Paperbound \$1.75

THREE MARTIAN NOVELS, Edgar Rice Burroughs. Contains: Thuvia, Maid of Mars; The Chessmen of Mars; and The Master Mind of Mars. High adventure set in an imaginative and intricate conception of the Red Planet. Mars is peopled with an intelligent, heroic human race which lives in densely populated cities and with fierce barbarians who inhabit dead sea bottoms. Other exciting creatures abound amidst an inventive framework of Martian history and geography. Complete unabridged reprintings of the first edition. 16 illustrations by J. Allen St. John. vi + 499pp. 5¾ x 8½.

T39 Paperbound \$1.85

THREE PROPHETIC NOVELS BY H. G. WELLS, edited by E. F. Bleiler. Complete texts of "When the Sleeper Wakes" (1st book printing in 50 years), "A Story of the Days to Come," "The Time Machine" (1st complete printing in book form). Exciting adventures in the future are as enjoyable today as 50 years ago when first printed. Predict TV, movies, intercontinental airplanes, prefabricated houses, air-conditioned cities, etc. First important author to foresee problems of mind control, technological dictatorships. "Absolute best of imaginative fiction," *N. Y. Times*. Introduction. 335pp. 5¾ x 8.

T605 Paperbound \$1.50

28 SCIENCE FICTION STORIES OF H. G. WELLS. Two full unabridged novels, *MEN LIKE GODS* and *STAR BEGOTTEN*, plus 26 short stories by the master science-fiction writer of all time. Stories of space, time, invention, exploration, future adventure—an indispensable part of the library of everyone interested in science and adventure. **PARTIAL CONTENTS:** Men Like Gods, The Country of the Blind, In the Abyss, The Crystal Egg, The Man Who Could Work Miracles, A Story of the Days to Come, The Valley of Spiders, and 21 more! 928pp. 5¾ x 8.

T265 Clothbound \$4.50

THE WAR IN THE AIR, IN THE DAYS OF THE COMET, THE FOOD OF THE GODS: THREE SCIENCE FICTION NOVELS BY H. G. WELLS. Three exciting Wells offerings bearing on vital social and philosophical issues of his and our own day. Here are tales of air power, strategic bombing, East vs. West, the potential miracles of science, the potential disasters from outer space, the relationship between scientific advancement and moral progress, etc. First reprinting of "War in the Air" in almost 50 years. An excellent sampling of Wells at his storytelling best. Complete, unabridged reprintings. 16 illustrations. 645pp. 5¾ x 8½.

T1135 Paperbound \$2.00

Catalogue of Dover Books

SEVEN SCIENCE FICTION NOVELS, H. G. Wells. Full unabridged texts of 7 science-fiction novels of the master. Ranging from biology, physics, chemistry, astronomy to sociology and other studies, Mr. Wells extrapolates whole worlds of strange and intriguing character. "One will have to go far to match this for entertainment, excitement, and sheer pleasure . . ." *NEW YORK TIMES*. Contents: *The Time Machine*, *The Island of Dr. Moreau*, *First Men in the Moon*, *The Invisible Man*, *The War of the Worlds*, *The Food of the Gods*, *In the Days of the Comet*. 1015pp. 5½ x 8. T264 Clothbound \$4.50

BEST GHOST STORIES OF J. S. LE FANU, Selected and introduced by E. F. Bleiler. LeFanu is deemed the greatest name in Victorian supernatural fiction. Here are 16 of his best horror stories, including 2 nouvelles: "Carmilla," a classic vampire tale couched in a perverse eroticism, and "The Haunted Baronet." Also: "Sir Toby's Will," "Green Tea," "Schalken the Painter," "Ullor de Lacy," "The Familiar," etc. The first American publication of about half of this material: a long-overdue opportunity to get a choice sampling of LeFanu's work. New selection (1964). 8 illustrations. 5¾ x 8¾. T415 Paperbound \$1.85

THE WONDERFUL WIZARD OF OZ, L. F. Baum. Only edition in print with all the original W. W. Denslow illustrations in full color—as much a part of "The Wizard" as Tenniel's drawings are for "Alice in Wonderland." "The Wizard" is still America's best-loved fairy tale, in which, as the author expresses it, "The wonderment and joy are retained and the heartaches and nightmares left out." Now today's young readers can enjoy every word and wonderful picture of the original book. New introduction by Martin Gardner. A Baum bibliography. 23 full-page color plates. viii + 268pp. 5¾ x 8. T691 Paperbound \$1.50

GHOST AND HORROR STORIES OF AMBROSE BIERCE, Selected and introduced by E. F. Bleiler. 24 morbid, eerie tales—the cream of Bierce's fiction output. Contains such memorable pieces as "The Moonlit Road," "The Damned Thing," "An Inhabitant of Carcosa," "The Eyes of the Panther," "The Famous Gilson Bequest," "The Middle Toe of the Right Foot," and other chilling stories, plus the essay, "Visions of the Night" in which Bierce gives us a kind of rationale for his aesthetic of horror. New collection (1964). xxii + 199pp. 5¾ x 8¾. T767 Paperbound \$1.00

HUMOR

MR. DOOLEY ON IVRYTHING AND IVRYBODY, Finley Peter Dunne. Since the time of his appearance in 1893, "Mr. Dooley," the fictitious Chicago bartender, has been recognized as America's most humorous social and political commentator. Collected in this volume are 102 of the best Dooley pieces—all written around the turn of the century, the height of his popularity. Mr. Dooley's Irish brogue is employed wittily and penetratingly on subjects which are just as fresh and relevant today as they were then: corruption and hypocrisy of politicians, war preparations and chauvinism, automation, Latin American affairs, superbombs, etc. Other articles range from Rudyard Kipling to football. Selected with an introduction by Robert Hutchinson. xii + 244pp. 5¾ x 8½. T626 Paperbound \$1.00

RUTHLESS RHYMES FOR HEARTLESS HOMES and MORE RUTHLESS RHYMES FOR HEARTLESS HOMES, Harry Graham ("Col. D. Streamer"). A collection of Little Willy and 48 other poetic "disasters." Graham's funniest and most disrespectful verse, accompanied by original illustrations. Nonsensical, wry humor which employs stern parents, careless nurses, uninhibited children, practical jokers, single-minded golfers, Scottish lairds, etc. in the leading roles. A precursor of the "sick joke" school of today. This volume contains, bound together for the first time, two of the most perennially popular books of humor in England and America. Index. vi + 69pp. 5¾ x 8. T930 Paperbound 75¢

A WHIMSEY ANTHOLOGY, Collected by Carolyn Wells. 250 of the most amusing rhymes ever written. Acrostics, anagrams, palindromes, alphabetical jingles, tongue twisters, echo verses, alliterative verses, riddles, mnemonic rhymes, interior rhymes, over 40 limericks, etc. by Lewis Carroll, Edward Lear, Joseph Addison, W. S. Gilbert, Christina Rossetti, Chas. Lamb, James Boswell, Hood, Dickens, Swinburne, Leigh Hunt, Harry Graham, Poe, Eugene Field, and many others. xiv + 221pp. 5¾ x 8½. T195 Paperbound \$1.25

MY PIOUS FRIENDS AND DRUNKEN COMPANIONS and MORE PIOUS FRIENDS AND DRUNKEN COMPANIONS, Songs and ballads of Conviviality Collected by Frank Shay. Magnificently illuminated by John Held, Jr. 132 ballads, blues, vaudeville numbers, drinking songs, cowboy songs, sea chanteys, comedy songs, etc. of the Naughty Nineties and early 20th century. Over a third are reprinted with music. Many perennial favorites such as: *The Band Played On*, *Frankie and Johnnie*, *The Old Grey Mare*, *The Face on the Bar-room Floor*, etc. Many others unlocatable elsewhere: *The Dog-Catcher's Child*, *The Cannibal Maiden*, *Don't Go in the Lion's Cage Tonight*, *Mother*, etc. Complete verses and introductions to songs. Unabridged republication of first editions, 2 Indexes (song titles and first lines and choruses). Introduction by Frank Shay. 2 volumes bounds as 1. Total of xvi + 235pp. 5¾ x 8½. T946 Paperbound \$1.25

Catalogue of Dover Books

MAX AND MORITZ, Wilhelm Busch. Edited and annotated by H. Arthur Klein. Translated by H. Arthur Klein, M. C. Klein, and others. The mischievous high jinks of Max and Moritz, Peter and Paul, Ker and Plunk, etc. are delightfully captured in sketch and rhyme. (Companion volume to "Hypocritical Helena.") In addition to the title piece, it contains: Ker and Plunk; Two Dogs and Two Boys; The Egghead and the Two Cut-ups of Corinth; Deceitful Henry; The Boys and the Pipe; Cat and Mouse; and others. (Original German text with accompanying English translations.) Afterword by H. A. Klein. vi + 216pp. 5½ x 8½.
T181 Paperbound \$1.15

THROUGH THE ALIMENTARY CANAL WITH GUN AND CAMERA: A FASCINATING TRIP TO THE INTERIOR, Personally Conducted by George S. Chappell. In mock-travelogue style, the amusing account of an imaginative journey down the alimentary canal. The "explorers" enter the esophagus, round the Adam's Apple, narrowly escape from a fierce Amoeba, struggle through the impenetrable Nerve Forests of the Lumbar Region, etc. Illustrated by the famous cartoonist, Otto Soglow, the book is as much a brilliant satire of academic pomposity and professional travel literature as it is a clever use of the facts of physiology for supremely comic purposes. Preface by Robert Benchley. Author's Foreword. 1 Photograph. 17 illustrations by O. Soglow. xii + 114pp. 5½ x 8½.
T376 Paperbound \$1.00

THE BAD CHILD'S BOOK OF BEASTS, MORE BEASTS FOR WORSE CHILDREN, and A MORAL ALPHABET, H. Belloc. Hardly an anthology of humorous verse has appeared in the last 50 years without at least a couple of these famous nonsense verses. But one must see the entire volumes—with all the delightful original illustrations by Sir Basil Blackwood—to appreciate fully Belloc's charming and witty verses that play so subacutely on the platitudes of life and morals that beset his day—and ours. A great humor classic. Three books in one. Total of 157pp. 5½ x 8.
T749 Paperbound \$1.00

THE DEVIL'S DICTIONARY, Ambrose Bierce. Sardonic and irreverent barbs puncturing the pomposities and absurdities of American politics, business, religion, literature, and arts, by the country's greatest satirist in the classic tradition. Epigrammatic as Shaw, piercing as Swift, American as Mark Twain, Will Rogers, and Fred Allen. Bierce will always remain the favorite of a small coterie of enthusiasts, and of writers and speakers whom he supplies with "some of the most gorgeous witticisms of the English language." (H. L. Mencken) Over 1000 entries in alphabetical order. 144pp. 5½ x 8.
T487 Paperbound \$1.00

THE COMPLETE NONSENSE OF EDWARD LEAR. This is the only complete edition of this master of gentle madness available at a popular price. A BOOK OF NONSENSE, NONSENSE SONGS, MORE NONSENSE SONGS AND STORIES in their entirety with all the old favorites that have delighted children and adults for years. The Dong With A Luminous Nose, The Jumbies, The Owl and the Pussycat, and hundreds of other bits of wonderful nonsense. 214 limericks, 3 sets of Nonsense Botany, 5 Nonsense Alphabets. 546 drawings by Lear himself, and much more.
T167 Paperbound \$1.00

SINGULAR TRAVELS, CAMPAIGNS, AND ADVENTURES OF BARON MUNCHAUSEN, R. E. Raspe, with 90 illustrations by Gustave Doré. The first edition in over 150 years to reestablish the deeds of the Prince of Liars exactly as Raspe first recorded them in 1785—the genuine Baron Munchausen, one of the most popular personalities in English literature. Included also are the best of the many sequels, written by other hands. Introduction on Raspe by J. Carswell. Bibliography of early editions. xliv + 192pp. 5½ x 8.
T698 Paperbound \$1.00

HOW TO TELL THE BIRDS FROM THE FLOWERS, R. W. Wood. How not to confuse a carrot with a parrot, a grape with an ape, a puffin with nuffin. Delightful drawings, clever puns, absurd little poems point out farfetched resemblances in nature. The author was a leading physicist. Introduction by Margaret Wood White. 106 illus. 60pp. 5½ x 8.
T523 Paperbound 75¢

JOE MILLER'S JESTS OR, THE WITS VADE-MECUM. The original Joe Miller jest book. Gives a keen and pungent impression of life in 18th-century England. Many are somewhat on the bawdy side and they are still capable of provoking amusement and good fun. This volume is a facsimile of the original "Joe Miller" first published in 1739. It remains the most popular and influential humor book of all time. New introduction by Robert Hutchinson. xxi + 70pp. 5½ x 8½.
T423 Paperbound \$1.00

Prices subject to change without notice.

Dover publishes books on art, music, philosophy, literature, languages, history, social sciences, psychology, handicrafts, orientalia, puzzles and entertainments, chess, pets and gardens, books explaining science, intermediate and higher mathematics, mathematical physics, engineering, biological sciences, earth sciences, classics of science, etc. Write to:

Dept. catrr.

Dover Publications, Inc.

180 Varick Street, N.Y. 14, N.Y.

(continued from front flap)

Chance, Luck and Statistics, Horace C. Levinson. \$2.00

The Fourth Dimension Simply Explained, edited by Henry P. Manning. \$1.50

Introductory Non-Euclidean Geometry, Henry P. Manning. \$1.00

Higher Mathematics for Students of Chemistry and Physics, Joseph W. Mellor. \$2.50

How Do You Use a Slide Rule? Arthur A. Merrill. 75¢

Mathematical Excursions, Helen Abbot Merrill. \$1.25

On Mathematics and Mathematicians, Robert E. Moritz. \$2.00

Cardano: The Gambling Scholar, Oystein Ore. \$1.60

Guide to the Literature of Mathematics and Physics, Nathan G. Parke III. \$2.75

Mathematics and Science: Last Essays, Henri Poincaré. \$1.25

Geometric Exercises in Paper Folding, T. Sundara Row. \$1.00

Great Ideas of Modern Mathematics: Their Nature and Use, Jagjit Singh. \$2.00

Applied Mathematics for Radio and Communication Engineers, Carl E. Smith. \$1.75

Teach Yourself the Slide Rule, Burns Snodgrass. Clothbound \$2.00

How to Calculate Quickly, Henry Sticker. \$1.00

Mathematics in Action, O. G. Sutton. Clothbound \$3.50

Concepts and Methods of Arithmetic, Marvin C. Volpel. \$2.50

An Introduction to Projective Geometry, Roy M. Winger. \$2.00

Monographs on Topics of Modern Mathematics, edited by J. W. A. Young. \$2.00

Paperbound unless otherwise indicated. Prices subject to change without notice. Available at your book dealer or write for free catalogues to Dept. TF 4, Dover Publications, Inc., 180 Varick St., N. Y., N. Y. 10014. Please indicate field of interest. Each year Dover publishes over 150 classical records and books on art, music, languages, philosophy, science, engineering, advanced mathematics, chess, humor, puzzles and other areas.

Felix Klein

Famous Problems of Elementary Geometry

This expanded version of the famous 1894 Easter lectures at Göttingen University has been accepted as a modern mathematical classic, and has been translated into four different languages. Using techniques of modern mathematics, it examines three famous problems which were intensely investigated in premodern mathematics: doubling the volume of a cube, trisecting an angle, squaring a circle.

Written with all Felix Klein's mathematical breadth, clarity, and profundity, this volume provides answers to modern problems connected with these three problems of the past. It is especially interesting to the modern student in answering such questions as: Under what circumstances is a geometric construction possible? By what means can a geometric construction be effected? What are transcendental numbers, and how can you prove that e and π are transcendental? Treatment is simple, and no knowledge of higher mathematics is required.

Contents. I. The Possibility of the Construction of Algebraic Expressions. Algebraic equations solvable by square roots. The delian problem and the trisection of the angle. The division of the circle into equal parts. The construction of the regular polygon of 17 sides. General considerations of algebraic constructions. II. Transcendental Numbers and the Quadrature of the Circle. Cantor's demonstration of the existence of transcendental numbers. Historical survey of the attempts at the computation of e . Transcendence of the number e . Trisection of the angle and the geometric construction of π . Discussion in detail of Gaussian polygons, Fermat's Last Theorem, and the impossibility of squaring the circle. Finiteness of π , and similar topics.

Unabridged, unaltered republication
Translated by W. W. Beman and D. L. E. Jones
Prefaces. xi + 92pp. 5 7/8 x 8.



Library

IAS, Shimla

513 K 672 F



00039358

1298 Paperbound \$1.25

A DOVER EDITION DESIGNED FOR YEARS OF USE!

We have made every effort to make this the best book possible. Our paper is opaque, with minimal show-through; it will not discolor or become brittle with age. Pages are sewn in signatures, in the method traditionally used for the best books, and will not drop out, as often happens with paperbacks held together with glue. Books open flat for easy reference. The binding will not crack or split. This is a permanent book.

