

NDIAN INSTITUTE OF ADVANCED STUDY LIBRARY * SIMLA

HUMAN SKELETAL REMAINS FROM HARAPPA

MEMOIR NO 9: 1962

3.

HUMAN SKELETAL REMAINS FROM HARAPPA

(WITH SEVENTYTWO PLATES AND ONE HUNDRED ELEVEN FIGURES)

FOREWORD

NIRMAL KUMAR BOSE

DIRECTOR

ANTHROPOLOGICAL SURVEY OF INDIA

THE ARCHAEOLOGICAL BACKGROUND

A. GHOSH
DIRECTOR-GENERAL
ARCHAEOLOGICAL SURVEY OF INDIA

DEMOGRAPHIC NOTES ON HARAPPA SKELETONS J. M. DATTA

HUMAN REMAINS FROM HARAPPA

P. GUPTA

P. C. DUTTA

A. Basu



ANTHROPOLOGICAL SURVEY OF INDIA GOVERNMENT OF INDIA CALCUTTA 13 WITHOUT THE EXPRESSED PERMISSION FROM THE DIRECTOR,
ANTHROPOLOGICAL SURVEY OF INDIA
ANY ARTICLE, PLATES OR FIGURES OF THIS MEMOIR
CANNOT BE REPRODUCED PARTLY OR WHOLLY.



Price: Rupees Twentytwo and Seventyfive naye Paise only



Published by Nirmal Kumar Rose, Director, Anthropological Survey of India, Indian Museum, 27 Chowringhee Road, Calcutta 13 and Printed by Sri Praboth Chandra Mitra, M.A. from Venus Printing Works, 52-7 Bipin Beharl Ganguli Street, Calcutta 12

CONTENTS

FOREWORD	V	NIRMAL KUMAR BOSE
THE ARCHAEOLOGICAL BACKGROUND	1	A. GHOSH
DEMOGRAPHIC NOTES ON HARAPPA SK	ELETONS 6	J. M. DATTA
HUMAN REMAINS FROM HARAPPA	13	P. GUPTA P. C. DUTTA A. BASU
ANTHROPOLOGICAL INTRODUCTION	13	
SQUARE (CEMETERY) R 37	18	
AREA G 289	81	
AREA (MOUND) AB	113	
CEMETERY H STRATUM II (Open Burials)	118	
CEMETERY H STRATUM I (Jar Burials)	149	
GENERAL OBSERVATIONS	177	·
REFERENCES	181	
APPENDIX	18 3	
COLLECTIVE TABLES	i - xe	

LIST OF ILLUSTRATIONS

HALF-TONES

Plates

I Tarkhanwala Dera: Cremation ground II Skl. 10 Norma Lateralis (Wash on dioptographic contour) Ш Skl. H 793 (B) Norma Frontalis (Wash on dioptographic contour) IV Skl. H 793 (B) Norma Lateralis (Wash on dioptographic contour) \mathbf{v} Skl. H 794 Norma Frontalis (Wash on dioptographic contour) VISkl. H 794 Norma Lateralis (Wash on dioptographic contour) VII Skl. H 796 (B) Nørma Frontalis (Wash on dieptographic contour) VIII Skl. H 796 (B) Norma Lateralis (Wash on dioptographic contour) IX Skl. H 798 (a) Norma Frontalis (Wash on dioptographic contour) X Skl. H 798 (a) Norma Lateralis (Wash on dioptographic contour) Mandibles showing attrition of incisor teeth : Cemetery R 37XI Skl. 10 Norma Frontalis (Wash on dioptographic contour) XII Photographs of Type A Male [Skl. H 820 (iii)] : Cemetery R 37XIII Photographs of Type A Female [Skl. H 798 (a)]: Cemetery R 37 XIV Photographs of Type A₁ Male [Skl. H 793 (B)]: Cemetery R 37 XVPhotographs of Type A1 Female [Skl. H 801/A]: Cemetery R 37 XVI Anterior aspect of Tibia and Femur : Cemetery R 37XVII Normae of Skl. H 779 (a) XVIII Normae of Skl. H 793 (A) XIX Normae of Skl. H 794 XXNormae of Skl. H 805 XXI Normae of Skl. H 806 XXII Normae of Skl. H 806/A XXIII Figs. 1-3 Normae of Skl. H 811; Figs. 4-6 Normae of Skl. H 818-XXIV Normae of Skl. H 816 XXV Normae of Skl. 2 XXVI Normae of Skl. 10 XXVII Occlusal and Left Lateral aspect of Mandible : Cemetery R 37 XXVIII Femur, Tibia and Fibula: Cemetery R 37 XXIX Skl. II S 18 Norma Lateralis (Wash on dioptographic contour) XXXSkulls showing cranial injuries : Area G XXXI Skl. III S 47 Norma Lateralis (Wash on dioptographic contour) MXXII Skl. III S 47 Norma Frontalis (Wash en dieptegraphic centeur) IIIXXX Photographs of Type A, Male (Skl. III S 23) : Area G XXXIV Photographs of Type A, Female (Skl. III S 21) : Area G XXXV Photographs of Type B. Male (Skl. III S 47) : Area G XXXVI Photographs of Type B, Female (Skl. II S 5) : Area G XXXVII Normae of Skl. I S 11 XXXVIII Normae of Skl. III S 22 XXXXIX Normae of Skl. III S 2 XLOcclusal and Left Lateral aspect of mandible : Area G XLI Skl. H 344 Norma Frontalis (Wash on dioptographic contour) XLII Skl. H 344 Norma Lateralis (Wash on dieptegraphic contour) XLIII Ski. Pet. No. 12 (a) Norma Frontalis (Wash on dioptographic content) XLIV Skl. Pot. No. 12 (a) Norma Lateralis (Wash on dioptographic contour) XLV

HALF-TONES

Р			

Tates	
XLVI	Photographs of Type A Male [Skl. H 255 (a)] of Cemetery H St. I (Jar burials)
XLVII	Photographs of Type A_2 Female (Skl. Pot. No. 11) of Cemetery H St. I (Jar burials)
XLVIII	Normae of Skl. H 344
XLIX	Normae of Skl. H 154 (a)
L	Skl. H 11635 Norma Frontalis (Wash on dioptographic contour)
LI	Skl. H 11635 Norma Lateralis (Wash on dioptographic contour)
LII	Normae of Skl. H 11635
LIII	Skl. H 698 Norma Frontalis (Wash on dioptographic contour)
LIV	Skl. H 698 Norma Lateralis (Wash on dioptographic contour)
LV	Skl. H 699 Norma Frontalis (Wash on dioptographic contour)
LVI	Skl. H 699 Norma Lateralis (Wash on dioptographic contour)
LVII	Photographs of Type B_2 Male (Skl. H 698) of Cemetery H St. II (Open burials)
LVIII	Photographs of Type A Female (Skl. H 488) of Cemetery H St. II (Open burials)
LIX(a)	Photographs of Type A ₁ Male [H 487 (a)] of Cemetery H St. II (Open burials)
LIX(b)	Photographs of Type A_1 Female (Skl. H 88) of Cemetery H St. II (Open burials)
LX	Anterior or Flexor aspect of the Humures (Cemetery H St. II)
LXI	Figs. 1-3 Radial aspect of Ulna; Figs. 4-7 Volar aspect of Radius : Cemetery H St. II (Open burials)
LXII	Anterior aspect of Tibia and Femur : Cemetery H St. II
LXIII	Normae of Skl. H 695
LXIV	Occlusal and Left lateral aspect of Mandible: Cemetery H St. II (Open burials)
LXV	Flexor and Extensor aspect of Humerus : Cemetery H St. II (Open burials)
LXVI	Normae of Skl. H 247 (b)
LXVII	Occlusal and Left lateral aspect of Mandible : Cemetery H St. I (Jar burials)
LXVIII	Fig. 1 Skeleton of an infant in embryonic position (Jar buriats) Fig. 2 Occipital view showing cranial injury (Skl. H 698) Fig. 3 Endocranial cast of Skl. H 5440 D (Mound AB)
LXIX	Fig. 1 Flexor aspect of Humerus (Jar burials) Fig. 2 Radial aspect of Ulna (Jar burials) Figs 3. & 4 Volar aspect of articulated radius and Ulna (Open burials)
LXX	Fig. 1. Flexor aspect of humerus: Skl. 10, 796/B and H 818 Fig. 2 Anterior aspect of femora: H 698, H 710 Fig. 3 Anterior view of tibia: H 710, H 698 Fig. 4 Anterior view of tibia & fibula in articulation H 710
LXXI	Hand and Foot : Cemetery H St. II (Open burials)

LINE DRAWINGS

Figures

1	Skl. H 779 (a)	Norma Frontalis	22
2	Skl. H 779 (a)	Norma Lateralis	22
3	Skl H. 779 (e)	Norma Frontalis	.23
4	Skl. H 779 (e)	Norma Lateralis	23
5	Skl. H 779 (c)	Mid-Sagittal Curve	24
6	Skl. H 779 (e)	Mid-Sagittal Curve	24
7	Skl. H 780	Norma Frontalis	25
8	Skl. H 780	Norma Lateralis	25
9	Skl. H 793 (A)	Norma Frentalis	26
10	Skl. H 793 (A)	Norma Lateralis	26
11	Skl. H 780	Mid-Sagittal Curve	27
12	Skl. H 793 (A)	Mid-Sagittal Curve	27
13	Skl. H 793 (B)	Mid-Sagittal Curve	29
14	Skl. H 794	Mid-Sagittal Curve	29
15	Skl. H 798 (A)	Norma Frentalis	31
16	Skl. H 798 (A)	Norma Lateralis	31
17	Skl. H 798 (A)	Mid-Sagittal Curve	32
18	Skl. H 796 (B)	Mid-Sagittal Curve	32
19	Skl. H 798/A1	Norma Frontalis	33
20	Skl. H 798/A1	Norma Lateralis	33
21	Skl. H 798 (a)	Mid-Sagittal Curve	34
22	Skl. H 798/A1	Mid-Sagittal Curve	34
23	Skl. H 801/A	Mid-Sagittal Curve	36
24	Skl. H 805	Mid-Sagittal Curve	36
25	Skl. H 806	Norma Frontalis	37
26	Skl. H 806	Norma Lateralis	37
27	Skl. H 806	Mid-Sagittal Curve	38
28	Skl. H 806/A	Mid-Sagittal Curve	38
20	Skl. H 806/A	Norma Frontalis	39
3 0	Skl. H 806/A	Nørma Lateralis .	39 40
31	Skl. H 811	Mid-Sagittal Curve	40
32	Skl. H 812	Mid-Sagittal Curve	41
33	Skl. H 816	Norma Frontalis	41
٠4	Skl. H 816	Norma Lateralis	42
35	Skl. H 816	Mid-Sagittal Curve	42
36	Skl. H 818	Mid-Sagittal Curve	
37	Skl. H 820 (I)	Norma Frontalis	44
38	Skl. H 820 (1)	Norma Lateralis	44
39	Skl. H 820 (I)	Mid-Sagittal Curve	45
40	Skl. H-820 (iii)	Mid-Sagittal Curve	45

		. '	
41	Skl. 1	Norma Frontalis	46
42	Skl. 1	Norma Lateralis	46
43	Skl. 1	Mid-Sagittal Curve	47
44	Skl. 2	Mid-Sagittal Curve	48
45	Skl. 10	Mid-Sagittal Curve	48
46,	47, 48	Dioptographic Contours of Type A Male [Skl. H 820 (iii)] of Cemetery R 37	52
49,	50, 51	Dioptographic Contours of Type A Female [Skl. H 798 (a)] of Cemetery R 37	53
52,	53, 54	Dioptographic Contours of Type A ₁ Male [Skl. H 793 (B)] of Cemetery R 37	56
55,	56, 57	Dioptographic Contours of Type A ₁ Female [Skl. H 801/A of Cemetery R 37	57
58	Skl. I S 11	Norma Lateralis	82
59	Skl. I S 11	Mid-Sagittal Curve	83
60	Skl. II S 18	Mid-Sagittal Curve	83
61	Skl. II S 42	Norma Lateralis	84
6 2	Skl. III S 22	Norma Frontalis	85
6 3	Skl. III S 22	Norma Lateralis	85
64	Skl. II S 42	Mid-Sagittal Curve	86
65	Skl. III S 47	Mid-Sagittal Curve	86
66	Skl. II S 5	Mid-Sagittal Curve	88
67	Skl. III S 21	Mid-Sagittal Curve	88
68,	69, 70	Dioptographic Contours of Type A ₁ Male [Skl. III S 23] of Area G	92
71,	72, 73	Dioptographic Contours of Type A ₁ Female [Skl. III S 21] of Area G	93
	75, 76	Dioptographic Contours of Type B ₁ Male [Skl. III S 47] of Area G	96
77,	78, 79	Dioptographic Contours of Type B ₁ Female [Skl. II S 5] of Area G	97
80	Skl. 11635	Norma Verticalis	114
81	Skl. 11635	Mid-Sagittal Curve	115
82	Skl. Mound AB (Rec.) Mid-Sagittal Curve	115
8 3		Rec.) Norma Frontalis	116
84	Skl. Mound AB (1	Rec.) Norma Lateralis	116
85	Skl. H 307 (a)	Mid-Sagittal Curve	120
86	Skl. H 488	Mid-Sagittal Curve	120
87	Skl. H 698	Mid-Sagittal Curve	121
88	Skl. H 488	Norma Frontalis	122
89	Skl. H 699	Norma Lateralis	123
90	Skl. H 710	Nørma Verticalis	123
91,	92, 9 3	Dieptographic Centours of Type B ₂ Male	
	+	[Skl. H 698] of Cemetery H St. II	100
		(Open burials)	128

Figures

94, 95, 96	Dioptographic Contours of Type A Female [Skl. H 488] of Cemetery H St. II	100
	(Open burials)	129
97 Skl. H 255 (a)	Mid-Sagittal Curve	152
98 Skl. H 344	Mid-Sagittal Curve	152
99 Skl. H 245 (e)	Norma Lateralis	153
100 Skl. H 7435 (b)	Norma Verticalis	154
101 Skl. H 7435 (b)	Norma Lateralis	154
102 Skl. H 246 (c)	Mid-Sagittal Curve	155
103 Skl. Pot. No. 11	Mid-Sagittal Curve	155
104, 105, 106	Dioptographic Contours of Type A Male	
	[Skl. H 255 (a)] of Cemetery H St. I	
	(Jar burials)	-00
107, 108, 109	Dioptographic Contours of Type A2 Female	
	[Skl. Pot. No. 11] of Cemetery H St. I	
	(Jar burials)	161
110	Volar projection of an Ulna [Skl. H 206 (d)]	162
111	Lateral projection of an Ulna [Skl. H 206 (d)]	162

913.031 Ar22

The Anthropological Survey of India has great pleasure in submitting the first report on the Harappān skeletal remains.

The Archaeological Survey of India began excavation in Haradda in 1921 and carried it on till after 1946. Dr B. S. Guha, who was originally attached to the Zoological Survey of India as Anthropologist, was entrusted with the task of reporting on the skeletal remains. He was responsible for the work till his retirement from the Anthropological Survey in 1954.

During his tenure of office as Director, Dr Guha succeeded in training up a fairly large number of very able assistants through whom the major portion of the work was actually conducted. Mr H. K. Bose who was originally employed by the Archaeological Survey was, in the main, responsible for the lifting up of the skeletons from 1930 to 1947. Later on, as a member of the staff of the Anthropological Survey of India, he was also responsible for reconstruction. The main work of reconstruction, however, was entrusted by Dr Guha to Mr M. Biswas, Senior Technical Assistant in the Survey. Mr Biswas is a skilled craftsman and had the advantage of training under Dr B. S. Guha, Dr A. K. Mitra, Dr B. K. Chatterjee, and Mr H. K. Bose. In the work of cleaning, preservation and partly of reconstruction others who helped were Messrs P. Gupta, B. N. Chatterji, J. N. Biswas, and H. N. Das.

FOREWORD

There is a story behind the delay after which it has been possible to present the scholarly world with at least the first report on Harappān remains.

The Harappān remains were removed to Banaras on account of the war. Unfortunately, during the heavy floods of 1943, several boxes containing the precious material were slightly damaged. It was only in 1948 that the remains were once more removed to Calcutta. By that time the Anthropological Survey of India had come into being as an independent department. Since 1948 the work has gone on steadily in the Osteology Laboratory of the Anthropological Survey in Calcutta.

In the Census Report of 1931, Vol. I, Part III, Dr B. S. Guha described briefly the important features of the Harappān skeletons which had been recovered up to that time. It is obvious from a foot-note at page laviii of the same report that a description of the Harappān human remains had already been sent to the press. But nothing is known of what happened about its printing. No copy was also available in the office of the Survey. Some notes prepared by Dr Guha were, however, incorporated in *Exacavation at Harappa*, Vol. I, by M. S. Vats. Sir Mortimer Wheeler's paper entitled 'Harappa 1946: The Defences and Cemetery R 37', published in *Ancient India*, No. 3, 1947, contains descriptive notes on the graves of Cemetery R 37 and Cemetery H st I, both exposed in 1946.

It appears from our office records, that some dioptographic drawings were prepared at Banaras; but these are not traceable. Later on, Dr B. K. Chatterjee and Mr G. D. Kumar also prepared the draft of a report on Harappān skeletons, which, could not however be utilized, for several reasons, during the preparation of the present report.

All measurements and drawings presented here are the result of the labours of Messrs P. Gupta, P. C. Dutta, A. Basu, A. Pal, Miss B. Sinha and Mrs A. Ray. This group worked on the skeletal material reconstructed by their predecessors from the 1st of August 1961 to the end of December 1961. The report was prepared by Messrs P. Gupta, P. C. Dutta and A. Basu between January 1962 and July 1962.

It may be pointed out here that the report on such a precious collection as Harappā ought to contain basic data as well as a comparison with skeletal material recovered from other contemporary sites. A comparison is also possible with the physical characteristics of different populations inhabiting India and the neighbouring countries at the present time. However valuable the latter part of the work may be, it was considered advisable to publish the basic data first of all for the use of the scholarly world. The work of comparison or inference drawn from them can be undertaken at any time later on by scholars either in India or abroad.

It has been with this end in view that the basic data are now being published. It will also be noted from the report that the scholars responsible for it have been very guarded in the nomenclature of various physical types described. Under advice, the types have been designated as A and B, split up into several sub-types. It was decided that it would be better not to give names to types as has already been done by previous authors; for that might suggest inferences which may not even-

tually be borne out by later discoveries or comparisons. But the present writers have been careful in indicating what names have been given to their types or sub-types by other scholars; so that the reader may easily equate one with the other.

As organizational head of the Anthropological Survey of India, I beg to place on record my appreciation of the enthusiasm and expeditiousness with which young scholars have prepared the report; and the skill and accuracy of craftsmanship with which Mr M. Biswas and others have actually succeeded in reconstructing the skeletal remains. My thanks are also due to a great measure to Stenographer Mr B. N. De, Photographer Mr S. Chattopadhyay, Artists Messrs R. C. Dey and B. N. Bagchi and Statisticians Messrs H. K. Nag, D. P. Mukherjee. M. N. Kaul and S. Chatterjee. Mr H. N. Mukherjee has been of assistance in proof-correction and preparation of the copy. Mr Arun Chakrabarti has been very largely responsible for efficient printing of the report. Messrs Venus Printing Works, Calcutta, to whom the task was entrusted, have shown admirable patience and courtesy: and it has also been on account of their appreciation of the importance of the work that it has been possible for the report to be printed within a reasonably short period of time.

Last, but not least, my thanks are due to my friend Mr J. M. Datta, formerly a Fellow of the Royal Statistical Society and of the Royal Economic Society, who has contributed a brief, but significant chapter on the population of Harappa. The data placed at his disposal have not naturally been sufficient; yet, he has thrown out a few suggestions with regard to the population which are certainly thought-provoking.

Anthropological Survey of India, Calcutta. Dated the 15th of August, 1962.

NIRMAL KUMAR BOSE
Director

The Archaeological Background

A. GHOSH
Director General of Archaeology in India

INTRODUCTION

How the excavations at Harappa in Montgomery District and at Mohenjo-daro in Larkana District, both now in West Pakistan, carried out by the Archaeological Survey of India, mostly in the twenties of the present century, revolutionized the concept of the origins of civilization in the Indo-Pakistan sub-continent is well-known and need not be dwelt upon here. It would suffice to say that later explorations have shown that the culture revealed at these two sites, initially named after the Indus but now more commonly called the Harappa culture, extended, variously in its pure state or in its ramifications, from Baluchistan on the west to beyond Delhi on the east, and from the foot-hills of the Himalayas in the north to the Tapti estuary in the south. Spatially, therefore, it was the most extensive of the Bronze-Age¹ cultures of south-western and southern Asia. On the temporal side, the latest view is that for Panjab and Sind 'a provisional dating of 2500-1500 B.C....responds consistently to current tests' 2.

To come directly to the subject-matter of the present memoir. The human skeletal remains found at Mohenjo-daro have been reported on and conclusions drawn therefrom³. In the excavations

I The Harappa culture has often been called 'chalcolithic' on account of the use of tools and other objects of stone along with those of copper by the people in their daily life.

² Mortimer Wheeler, The Indus Civilization, 2nd ed. (Cambridge, 1960), p. 99. It may be recorded here that the very recent (1962) Carbon-14 dating of objects from the late Harappan levels of Kalibangan, District Ganganagar (north Rajasthan), and of Lothal, District Ahmadabad (Gujarat), is, in both cases, 4080 ± 125 Before Present. Even making allowance for all limitations of this method of dating, one might wonder if 1500 B.C. for the

end of the Harappa culture does not err too much on the conservative side, though it may be argued that the suggestion holds good only for Panjab and Sind and has the advantage of establishing a possible causal connexion between the advent of the Aryans and the disappearance of the Harappans. Cf. Wheeler (1960), pp. 96-99. Our view is that nothing that has been said and excavated till now has established any connexion between the Harappans and the Aryans at any stage.

³ John Marshall, Mohenjo-daro and the Indus Civilization (London, 1931), I, pp. 107-08; R. B. Seymour Sewell and B. S. Guha in ibid., II, pp. 599-648.

at Harappa4 as well, human remains were encountered in various contexts, with which it is necessary for the reader of the present memoir to be acquainted and which are given below, to explain the circumstances of their discovery and the burial-customs they reveal.

AREA (MOUND) G5

In the southern portion of Trench II in this Area, there was a group of human skeletal remains, which were found tightly packed together between 4 ft. and 5 ft. 10 in. below surface and consisted of twenty complete human skulls and fragments of a few others, ten lower jaws, parts of vertebral columns and bones of hips, legs and arms, besides some animal-bones and a considerable quantity of pottery. Except in one case, in which the skull lay alongside the trunk, all bones were disarticulated and lay in a disorderly condition6. While regular burials are thus ruled out, at best fractional postexposure deposits, possibly within undetected pits, are uncertainly indicated. The evidence is too illdefined to be of help in determining the burial-custom(s) of the Harappans.

The pottery found with the bones cannot be regarded as real grave-furniture in that fragments of pots lay without any regular order. However, the pottery was typically Harappan but included two dishes-on-stand supposed to be of an aberrant type'.

AREA (MOUND) AB

Human remains, consisting of two fragmentary human skulls, a lower jaw and some other bones, a tubular terracotta bead and a fragment of the base of a dish-on-stand were found 'between the IIIrd and IVth strata's in this Area. Here also, while a post-exposure fractional burial seems to be indicated, the details are not clear.

SQUARE (CEMETERY) R 37

In 1938 and 1939, twentyseven regularly-disposed skeletons were excavated in Square R 37 with typical Harappan pottery as grave-furniture. A few more skeletons were excavated in the next two or three years10. As there could be no mistake that they lay in a regular cemetery, systematic

4 The excavations were carried out by Daya Ram Sahni from 1921 to 1924-25, by Madho Sarup Vats from 1926-27 to 1933-34 and by R.E.M. Wheeler in 1946. See also n. 10. Sahni's report appears in Annual Report, Archaeological Survey of India, 1920-21 (1923), pp. 15-17. 1923-24 (1926), pp. 52-54, and 1924-25 (1927), pp. 73-80. Vats' in his Excavations at Harappa, 2 vols. (Delhi, 1940), and Wheeler's in 'Harappa 1946: the Defences and Cemetery R 37', Ancient India, no. 3 (1947), pp. 58-130.

It may be necessary to explain here such sitelabels as G. AB, R 37, etc. Alexander Cunningham, who visited Harappa in 1872-73, numbered the mounds as A E. Archaeological Survey of India Reports, V (Calcutta, 1875), pl. XXXII. Later on, it must have been noticed that his A and B together represented a single mound': that mound was, therefore, called AB. At the same time, Cunningham's letters were extended by F to J to cover unnumbered mounds, Vats, II, pl. I. Simultaneously, the whole site of Harappa was gridded into 100-ft. squares for purposes of co-ordination, 'those running west to east being numbered A, B, C, etc., and similarly those going from north to south, 1, 2, 3, and so on', ibid., I, p. 8. This would explain 'R 37'.

- 6 Vats, I, pp. 198-202.
- 7 lbid., I, p. 233; II, pl. LXX, 10 and 14.
- 8 Ibid., I, pp. 161-62.
- 9 Ibid., I, p. 200 n.

10 The excavations of 1938-41 were carried by Sastri, Custodian, Archaeological Museum, Harappa, and H. K. Bose of the Anthropological Survey

excavation was undertaken in 1946 with a view to exposing a few more burials under closely-observed conditions and to linking stratigraphically Cemetery R 37 with the previously-discovered Cemetery H^{II} . As a result of the excavation, ten additional graves were identified; it was also established that 'the cemetery belongs to one and the same general stratum and was in continuous use.

The body was normally extended, occasionally on one side or the other, with the head to the north (between north-west and north-east, but usually within a few degrees of north)... Grave-pits varied in dimensions, ranging from 10 to 15 feet in length, $2\frac{1}{2}$ to 10 feet in width, and dug to a depth of 2 to 3 feet from the contemporary surface. An average grave measured superficially 10 by 3-4 feet, with a depth of 2 feet. The pit was generally wider towards the head. Its large size was due to the custom of including large quantities of pottery, mostly near the head but some also at the feet and along the sides and occasionally below the body. The number of pots accompanying a burial ranged from two to forty, with an average of fifteen to twenty. Most of the types were such as occur on habitation-sites of the mature Harappa culture.

'Personal ornaments were sometimes worn by the dead...

'Besides pottery and personal ornaments, toilet objects occasionally formed a part of the grave-furniture. From the total number of graves found in 1937-46, twelve yielded each a handled copper mirror; others produced mother-of-pearl shells; one an antimony stick; and one, a large shell spoon.

'It may be noted that some of the graves contained, besides a human skeleton, a few decayed animal bones. One grave included the bones of a fowl, together with a small handled lamp, placed at the feet of the dead' 12.

In one case, the body, probably of a female, was placed within a wooden coffin. There were also traces of some light-green substance over and around it¹³.

Extended inhumation, then, was the normal funeral practice of the Harappans at Harappa (no cemetery has as yet been identified at Mohenjo-daro); evidently, this was their practice elsewhere too, '4 to judge from two recently-excavated sites within the Indian border—Rupar in District Ambala

ing cremation-ground, marked off by flatly-laid mudbricks, in which there had been at least five cremations (pl. I). After each cremation, marked by ashes and bits of charred bone (sometimes collected in pots), the ground within the enclosure was levelled by a coating of clay or mud-bricks for the next cremation to take place. That the Harappans cremated at least five of their dead at this place seems established; but the conclusion that extended inhumation was the normal practice of the Harappans need not be prejudiced by this single isolated instance, particularly as in the neighbouring Harappan instance, particular, site of Binjor-3 (lat. 29°10' N.; long 73°05' E.), 6 miles to the west of Tarkhanwala-Dera and 3 miles to the east of the Indo-Pakistan border, the existence of a cemeof the industration of the course of the same exploration, in the flat land to the west of the mound, where large number of complete Harappan pots were found near the surface.

¹¹ Wheeler (1947), particularly pp. 83-90 and 101-120.

¹² Ibid., p. 86.

¹³ Ibid., p. 87. This burial 'represents a mode customary in Sumer during a considerable period which overlapped and probably preceded that of Cemetery R 37', ibid., p. 88. At the time of the excavation it was suspected that there was a reed-shroud below a part of the body, but this has now been disproved, K. A. Chowdhury and S. S. Ghosh in Ancient India, no. 7 (1951), p. 13.

¹⁴ It would be unfair if I do not mention here a case of Harappan cremation observed by me at Tarkhanwala-Dera (lat. 29910' N.; long 73°10' E.) in Ganganagar District (north Rajasthan), 4 miles to the north of Anupgarh, in the course of my exploration (1950-53) in the dried-up valley of the Sarasvati. Here, at the top of a made-up platform, on which stood a modest Harappan settlement, was noticed, by excavation, an oblong stand-

(Panjab)¹⁵ and Lothal in District Ahmadabad (Gujarat)¹⁶—which also have cemeteries with similar burials.

AREA (CEMETERY) H

Excavated between the years 1929-30 and 1933-34, Area H was, like R 37, very fruitful for human skeletal remains¹⁷. Here there were two 'strata' of burials (respectively called in this note Early and Late Cemetery H¹⁸, each characterized by its own method of burial: it was earth-burial in Early Cemetery H and pot-burial in Late. In the former, in the eastern section of the Area, complete bodies were seen laid in an extended way, while in the western section only fractional burials were met with. In Late Cemetery H, disarticulated bones, evidently collected after the exposure of the bodies, were found placed without any order at the bottom of the jars, too small to take complete bodies, except those of babies, which were found completely enclosed in an embryonic position.

In shape, fabric, slip and painted decoration, the pottery of Late Cemetery H is entirely different from that of the true Harappa pottery and is very easily distinguishable from the latter. It may be mentioned that this pottery also occurs here and there in the upper levels of the habitational mounds of Harappa. Even the pottery of Early Cemetery H had hardly anything in common with the typical Harappa, though the excavator hints at its possible derivation out of the latter as a result of a long period of development—a hypothesis difficult to sustain, notwithstanding his attempt to substantiate it on the basis of craniological data¹⁹.

It has been stated above (p. 3) that one of the objectives of the excavation of 1946 was to establish the stratigraphic relation between Cemetery R 37 and Cemetery H by means of a long trench. The excavation was pre-eminently successful in this, in that it clearly demonstrated that Late Cemetery H 'was not only later than R 37 but was subsequent to a deep intervening deposit of potsherds and other débris which indicated a considerable alteration of the site between the two cultures' 20. Though no burial of Early Cemetery H was encountered in the trench, observation showed that this cemetery too post-dated the débris.

That Cemetery H represents the influx of a new people at Harappa is amply established, but there is no evidence in regard to the archaeological affiliation of the people. Their identity with

- 16 Indian Archaeology 1954-55—A Review (1955), p. 9.
 16 Ibid., 1957-58 (1958), p. 10; 1958-59 (1959), pp. 14-15;
 1959-60 (1960), p. 19.
- 17 Vats, I. pp. 203-45. Three jars of Late Cemetery H were also exposed in 1946, Wheeler (1947), pp. 89-90.
- 18 These were respectively designated as Stratum II and Stratum I by Vats, and this system has been followed in the subsequent pages.
- 19 Vats, I., pp. 283-35. Wheeler (1947), p. 118, regards the two groups Early and Late Cemetery H. as successive phases of the same culture. Cf. Stuart Piggott, Prefilstoric India (Harmondsworth, 1950), p. 233. But the radical difference between the methods of burial in the
- two cemeteries remains unexplained by this single-culture hypothesis.
- 20 Wheeler (1947), p. 85. The published section, ibid., pl. XXXIX, graphically demonstrates this. At the same time, this, together with the total absence everywhere at Harappa of any overlap between the Harappa and Cemetery H material, precludes any contact between the two cultures as the conquered and the conqueror or as of peaceful co-existence. Cf. B. B. Lal in Ancient India, nos. 10 and 11 (1954 and 1955), p. 151 n. As Wheeler now cautiously observes (1960), p. 56: 'At least it is wise at present not to assume a temporal continuity between the Harappan culture and that of cemetery H'.



TARKHANWALA-DERA : CREMATION-GROUND

the Aryans has been hesitatingly suggested²¹, but this lacks plausibility, for the relics of the Aryans should be widespread and not localized, as the Cemetery H relics are.

POST-CREMATION BURIALS

Both at Harappa and Mohenjo-daro were found, in different levels, large wide-mouthed pottery urns, containing smaller vessels, bones of animals and other objects. It has been surmised that they represent burials in which human bones surviving cremation were deposited after powdering²². As, however, out of over two hundred and thirty urns found at Harappa²³, only one contained an unburnt human bone (tibia), their evidence may be discounted in the present context.

CONCLUSION

Such, in brief, are the circumstances of the discovery of human remains in the excavations at Harappa. In the exposing of the skeletons and in their field-treatment and lifting, the Archaeological Survey of India invariably received help from the officers of its sister-organization, the Anthropological Survey of India, to which the skeletons were removed from the field for joining, final preservation, study and scientific report. It is a matter of gratification that the long-awaited report is now ready.

From the archaeological point of view, the following are the prerequisites to any conclusions that the anthropologist may draw on the ethnic affinities of the people whose remains are reported on in the following pages. The remains from R 37 are to be studied first, individually and as a whole, for they indisputably belong to the real authors of the Harappa culture and therefore provide the most reliable material for racially affiliating them. The results thus obtained should then be applied to the remains from Area G, which also seemingly belong to the mature Harappans. The (later) Early Cemetery H material may thereafter be compared and contrasted with the Harappan material, and finally should come the (latest) Late Cemetery H material. As more than one archaeological culture is involved, any mixing up of the material will doubtless produce erroneous results.

Harappa has the unique advantage of having two protohistoric cemeteries successively superimposed upon an authentic Harappan cemetery. The comparative results of the anthropological study of the remains of these cemeteries will be of immense importance from all points of view.

Finally, the results, in so far as they relate to the real Harappan remains, have to be correlated with those already obtained from Mohenjo-daro and those to be obtained from two Indian sites with Harappan cemeteries, viz. Rupar and Lothal, which, I understand, are receiving the expeditious attention of the Anthropological Survey of India.

Demographic Notes on Harappa Skeletons

J. M. DATTA

NATURE OF THE POPULATION

The total number of skeletons is 260 only, the primary data have been given in page 13 of the 'Anthropological Introduction'.

The number whose age has been more precisely determined, i.e. within a certain range of time, is 107.

The skeletons obtained were not buried at one time. They belong to different periods of time. Their distribution has been given in $Table\ I$ (Anthropological Introduction) page 13.

Sundbarg's age-categories of population are:

	A G E 0-15	I _N 15-50	Υ	Ε	A	R S
Progressive Stationary	400 330	500 500				100 170
Regressive	200	500				300

This empirical relationship holds good all over the world both in space and in time.

Increasing the number of those who are between 0-12 years by one-fifth to get the number of 0-15 we find:

	AGE	G R	0 U P
	0-15		over 15
Cemetery H stratum I	39.6 (508)	38.4	(492) = 1000
Cemetery H stratum II	2.4 (92)	23.6	(908) = 1000
Mound Area	9.6 (384)	15.4	(616) = 1000
Area @ 289	7.2 (313)	15.8	(687) = 1000
Cemetery R 37	0	-	108

The population of Mound Area is of the progressive type; and that of Area G 289 of the stationary type. The other populations do not conform to any of these types. But having regard to the sex-ratio probably Area G 289 people were of the progressive type also.

Let us find out the sex-proportions as determined from skeletons. Unfortunately the sex of many subjects could not be determined; and that of children has not been possible.

	MALE	FEMALE	UNDETERMINED
Cemetery H stratum I	11	21	13
Cemetery H stratum II	11	9	4
Mound Area	6	6	5
Area G 289	9	4	4
Cemetery R 37	38	55	15

The sex-ratios are given below. As the number of subjects, whose sex has been determined is very small, any accidental ommission or failure to determine its sex, even of one, would effect the ratio seriously.

	SEX-RATIO PER 1000 MALES
Cemetery H stratum I	1909
Cemetery H stratum II	818
Mound Area	1000
Area G 289	444
Cemetery R 37	1450

The sex-proportions, except in the Mound Area, are abnormal. The excess or shortage of females may be explained by supposing that as Harrappa was a central city—one of the joint capitals, the structure, both age-structure and sex-ratio, were artificial. Normally there would be shortage of females; but if females are employed as slaves or impressed as labourers in certain periods, there would be an excess.

The proportion of older men i.e. 50 years and above is very small; it is 5 out of 107. Even lowering the age-limit to 40 years, the total is 12. This fact also goes to show that the population of the city of Harappa was not representative or a good sample of 'the Harappa people'.

An attempt may be made to find out the general fertility rate, expressed as the number of children per 1000 females aged 15-44 years. Here all those who are above 12 and below 50 have been taken as the base. As sex of as many as 17 per cent has not been determined the rates found are necessarily inaccurate. But they give some idea about the population. They tend to confirm our conclusions stated above.

Females of reproductive age period i.e. all those who are between 12 and 50 years.

	(1) (2) I 2 18 II 1 8 0 6 0 4	TOTAL	CHILDREN	FERTILITY RATE	
	(1)	(2)	(3)	0-12 YEARS (4)	(4) x 1000/(3) x 12
Cemetery H stratum I	2	18	20	33	137
Cemetery H stratum II	1	8	9	2	19
Mound Area	0	6	6	8	111
Area G 289	0	4	4	6	125
Cemetery R 37	3	52	55	0	0

With the data available we have not been able to reconstruct the age-pyramid, and find out the mean age.

AN ESTIMATE OF POPULATION IN HARAPPA AND MOHENJO-DARO

The capital cities of Harappa and Mohenjo-daro have been described in some detail by Stuart Piggot (*Prehistoric India*, 1950) and Sir Mortimer Wheeler (*The Indus Civilization*, 1953). In certain areas excavated beyond the workmen's quarters there appears to have been arrangements for pounding wheat into flour. It has been argued that much of the labour for this purpose was slavelabour. Both Wheeler and Stuart Piggot have described the elaborate bureaucratic machinery which was capable of organizing and distributing surplus wealth. Grain was stored as part of the governmental agricultural policy and municipal granaries or the preparation of flours was part of the controlled economy using organized labour.

One may therefore infer that there was considerable concentration of population at Harappa as well as at Mohenjo-daro. The question is, how many people actually lived either at Harappa or at Mohenjo-daro? We believe it is possible to make some kind of a rather rough guess.

'The combined floor-space of the twelve granaries (at Harappa) was something over 9000 sq. ft., and approximates closely to that of the Mohenjo-daro granary as originally planned' (Wheeler 1953, p. 22). The gross area of the Harappa granary is $12' \times 50' \times 20' = 12000$ sq. ft. From the plan of the granary at Mohenjo-daro as given by Wheeler at page 232 of his book, it appears that originally planned it extended over 8000 sq. ft. Later on additions were made and it occupied a space of 11260 sq. ft.

We shall take 12,000 sq. ft. as the area of the granary at Harappa and 11,260 sq. ft. as the area at Mohenjo-daro.

The height of the State granary has not been given by either of the authors, but it is possible to make some kind of an estimate.

Stuart Piggot says (p. 161) that the defensive wall of the citadel at Harappa was roughly 40 feet wide at the base and had an approximate height of 35 feet. That the walls of the granaries are buttressed shows that they were of considerable height. If the defensive walls of the citadel so

thick at the bottom rose to 35 feet only, the height of the granary walls certainly did not exceed that figure. One may take 35 feet to be the upper limit of its height.

Whatever may have been the total height, we have to deduct from it the thickness of the roof, some air space above the stacks of bags of grain or milled flour for ventilation and easy handling of the bags, and the height of podium to get the 'effective' height of the grain-stacks. The height of the podium being 4 feet and the roof, say, a foot or one and half feet thick; the air space being roughly $4\frac{1}{2}$ to 5 feet, we get an approximation of 25 feet as the 'effective height'.

The height of the granary at Mohenjo-daro can be roughly estimated from plate IX of Wheeler's book. It is about 5 times the height of the man standing at the bottom with out-stretched legs. So, up to the passage for ventilation, it is approximately 25 feet high. The breadth of the sub-structure as shown in figure VIII of the same book is about 15 feet.

We may now attempt to calculate the effective volumes of the two granaries at Harappa and Mohenjo-daro. At Harappa the volume is $12 \times 50' \times 20' \times 25' = 300,000$ cubic feet; at Mohenjo-daro 11,260 sq. ft. \times 24 = 270,240 cubic feet. The two figures are close to one another having regard to the estimates of height made already; though the height at Mohenjo-daro is 10% less than that at Harappa.

The aisle in the middle of the granary at Harappa is as broad as 23 feet; and there are open spaces between the sub-structures. We infer that grain could not have been stored in heaps, but were stored in bags. The presence of the circular working floors near the granary goes to show that grain was milled before storage.

As milled flour was stored in bags, some space is wasted at the corners, and in the interspaces between bags. We have been informed by several godown-keepers here in Bengal, that the volume occupied by flour is five-sixths of the total volume.

The volume occupied by milled flour is:

at Harappa

2.50,000 eu. ft.

at Mohenjo-daro

2,25,200 eu. ft.

The Punjab and Sind are wheat-producing countries. The Harappa people used wheat and barley (cf. Wheeler, p. 62). We have been informed by Sri Asoka Mitra, I.C.S., Registrar General of India that—

1 cu. ft. of imported common wheat weighs 21-22 seers.

1 cu. ft. of common Indian wheat weighs 25 seers.

1 cu. ft. of barley weighs

22 seers.

MEMOIR: No. 9

This is the weight as grain; but when it is milled and tightly packed, 1 cubic foot of common Indian wheat weighs as much as 29-30 seers.

As milling in ancient times was not as fine as in the present day, 1 cubic foot of wheat must have weighed less. We take it to have weighed 28 seers, or 0.7 maunds.

Consumption of food varies with age, sex and nature of daily labour. Adults consume more than old men or young ones; in the case of infants it is almost nil. Female consumption has been taken by Mulhall in the *Dictionary* of *Statistics* to be two-thirds that of a male. Modern research in England shows it not to exceed 85 per cent that of a male at certain ages.

Annual consumption of paddy per head, irrespective of age, sex and nature of work, has been taken to be 9 maunds (*Bengal Land Revenue Commission Report*, Vol. II, p. 106). And 20 manuals of paddy produce 13.5 maunds of rice. So the annual consumption of rice is 6 maunds per head.

Annual consumption of wheat per head is a little less, as wheat is more nutritious than rice. The standard diet which keeps a man fit prescribes 17 ounces of cereals a day, i.e. some 4.71 maunds per head per annum.

The actual consumption per head of the total population is likely to be less, as it includes children. But the standard diet includes many other items, which may not have been available in those early times. Further the consumption was likely to have been greater. For all these reasons, we take 4.71 maunds to be the annual consumption per head.

Drs R. C. Majumdar, H. C. Rai Chaudhuri and K. K. Datta in their Advanced History of India, say: 'Wheat was the principal article of food, but barley and palm-date were also familiar. They also used mutton, pork, fish and eggs' (p. 18).

Assuming that the grain stored was for a year's consumption (as milled wheat or barley is not likely to remain unaffected by mildews and other fungi and insects for more than a year) the population of the two cities of Harappa and Mohenjo-daro may be estimated thus:

Volume of milled grain at Harappa 2,50,000 cu. ft
Weight of milled grain 2,50,000 \times 0.7 mds

Number of persons at Harappa 2,50,000 \times 0.7 mds $2,50,000 \times 0.7 \\
4.71 = 37,155$ Similarly the population at Mohenjo-daro is $\frac{2,25,200 \times 0.7}{4.71} = 33,469$

We do not lay stress on the particular figures; but they do give us an idea of the sizes of the population.

'In the matter of actual plan and lay-out the two cities are strikingly similar, and Wheeler's recent work at Harappa has thrown much light on the significance of this uniform planning' (Stuart Piggot, p. 159).

The suggested street-plan of Mohenjo-daro as shown at page 25 of *The Indus Civilization* shows it to be almost a square. The site plan of Harappa at page 17 shows it to be a square.

Wheeler says: 'It (Harappa) overlies and adjoins the mounds of the ancient city, which appears to have had a circuit of not less than 3 miles'..... (p. 16).

Stuart Piggot says 'The mounds which represent the city of Mohenjo-daro today cover a square mile.....' (p. 165).

The area of Harappa is a little more than 3/4 sq. miles or 480 acres. We take it to be 500 acres in round numbers, as the full city cannot be recovered on account of brick-robbing during the construction of the Lahore-Multan Railway.

The area of Mohenjo-daro, we take to be 640 acres.

The density of population is at-

Harappa Mohenjo-daro 74 persons per acre. 52 persons per acre.

26.1

It appears that Harappa was more populous, and the greater density indicates that it was the older city. At Mohenjo-daro the granary was expanded by some 38 per cent between Period I and Period II B—which shows that it was a growing city. Non-extension of the granary and high density at Harappa indicate that the authorities for some reasons unknown to us did not like the extension of the city.

What this high density at such a remote time (Circa 2000 B.C.) means, we have not been able to guess. But to appreciate the problem, we give certain data below:


(a) Calcutta: The density of persons per acre at Calcutta during the census period has been:

444	CALCUTTA INCLUDING	MUNICIPAL CALC
	FORT AND MAIDAN	9.7
1872	25	35
1881	24	33
1891	27	37
1901	33	45
1911		50
	36	52
1921	38	58
1931	42	
1941	76	107
1951	92	130
1961	106	148
1002	700	

- (b) *Gaur*: In the 15th century, 'the capital city of Gaur (in Bengal) was beautified and expanded to an extent of over 20 sq. miles with a population of over 600,000' (M. N. Gupta, *Land System of Bengal*, p. 156). The density is 47 per acre.
- (c) Vijayanagar: Abd-er-Razzak visited the city in 1443. He says 'from the northern gate of the outer fortress to the southern is a distance of two statutory parsangs (about 7 or 8 miles) and the same with respect to the distance between the eastern and western gates' (Vincent Smith, Oxford History of India, p. 309).

The area enclosed is more than 50 sq. miles. Domingos Paes, a Portuguese, visited it about 1522 in the reign of Krishna Dev Raya when the Empire of Vijayanagar was at the height of its glory. Paes found a difficulty in estimating the size of the city, because the hills prevented him from seeing the whole at once. So far as he could judge, it was as large as Rome. The houses were said to exceed 100,000 in number. If that guess be near the truth, the population cannot be less than half a million (ibid p. 310).

As there is no reason to think of the diminution of the city in extent between 1443 and 1522, the density works out to 16 per acre. As hills and dales and rivulets were enclosed within the outer wall, the density for residential areas was greater, say 2 or 3 times the above figure. Even then it does not approach that of Harappa.



Norma Lateralis

Human Remains from Harappa

P. GUPTA, P. C. DUTTA & A. BASU

ANTHROPOLOGICAL INTRODUCTION

The human remains of altogether 260 individuals were recovered by excavations in Harappa between the years 1925-26 and 1946-47. Some individuals are represented by tiny fragments alone; but in many cases the entire skeleton is available, although often badly damaged by chemical action or earth-pressure. The sites yielding bones are as follows:

- (a) Cemetery H stratum I—jar burials
- (b) Cemetery H stratum II—open burials
- (c) Mound AB, F and Area J
- (d) Area G 289
- (e) Cemetery R 37

Their archaeological position has been indicated by Mr A. Ghosh, Director-General of Archaeology in India, in an introductory chapter entitled 'The Archaeological Background'.

Table 1 gives details of site, sex and age of all skeletal remains recovered.

TABLE 1 SITE, SEX AND AGE DISTRIBUTION OF ALL SKELETONS

Sites	Infant (Birth to 3 years)	Child (3 to 12	Juvenile*	Male Adult (Above 20 years)	Female Adult (Above 18 years)	Adult Sex indeter- minable	Total
material control of the second control of th	years)	years)		11	19	11	78
Cemetery H St I	20	13	4	11	8	4	26
Cemetery H St II		2	1	6	6	5	25
Mound Area	2	6	_	9	4	4	23
Area G 289		6	_	38	55	13	108
Cemetery R 37		provide.	2				
				75	92	37	260
Total	22	27	7				

^{*} Juveniles are not shown sexwise since all of them could not be properly determined.

While estimating the stature of the living from dry long bones Pearson's (1899: 196) regression formulae, Dupertuis and Hadden's (1951: 15-53) 'general formulae' and Manouvrier's* (1893: 347-402) tables were employed.

AGE AND SEX

Estimation of age at death of an individual was based on the extent of closure of sutures of skull and degree of ossification of bones. Eruption of tooth was also one of the criteria in the case of non-adults.

As regards determination of sex, there was difficulty as all relevant bones were not often available. As has already been mentioned, the majority of skeletons were in a fragmentary condition. Certain anatomical features of bones have a high degree of diagnostic value in regard to sex, and Krogman (1946: 154) is of opinion that when portions of skeletons are available, the degree of accuracy is as follows:

D. 1		
Pelvis alone	95%	accuracy
Skull alone		accuracy
Skull alone	92%	accuracy
Pelvis + skull	000	
	98%	accuracy
Long bones alone	00 0=0	
	80-85%	accuracy
Long bones + skull		-
	95%	accuracy
Long bones + pelvis	98%	
2	JO 70	accuracy

In the present Report, utilizing Trevor's (Mukherjee, Rao and Trevor 1955:9-10) principle (modyfying slightly), laboratory sexing of the entire skeletal material is based on anatomical appreciation of all the available bones pertaining to an individual. Only adult specimens were considered and the result is shown in Table 4. In many cases, bones were broken beyond repair and unsuitable for measurement, but they could still be used for sexing.

It is apparent from Table 4 that when the total skeletons are considered, laboratory sexing shows slightly higher proportion of females (55.09%) than males (44.91%), when M and M? are taken together and compared with F and F? also taken together.

When different stratum is taken into consideration it is observed that in Cemetery R 37 female is predominant (59.14%) over male (40.86%). Similarly in Cemetery H jar burials female population was numerically greater (63.33%) than male (36.67%). Reverse is the case of cemetery H open burials, i.e. 57.89% male and 42.10% female. In Area G 289 male was in overwhelming majority (69.23%) over female (30.77%).

The present study being the anthropological complement of Vat's and Wheeler's Reports on archaeology, its primary object is to determine the physical characteristics of the Harappāns.

Following Dwight's (1894:16) recommendation subtraction of 2 cm. from the mean of the total approximations was not done.

TABLE 4

SEX COMPOSITION* OF ADULT INDIVIDUALS

Basis of sexing		Cemetery H Stratum I (Jar burial)					Cemetery H Stratum II (Open burial)				Mound AB					Area G 289						Cemeiery R 37				
-	M	M ?	F	F ? T	'otal	M	M ?	F	F?T	otal	М	M ?	F	F ? T	otal	M	M ?	F	F?	Total	M	М?	F	F?	Total	
alvarium	3		3	_	6	1	_	_	_	1	1	_		1	2	6	1	4	_	11	8	1	10	1	20	
alvarium and mandible	_		_	_	_	_	_	_	_		_	_		_	_	2	_	_	-	2	1	_	3	1	5	
alvarium and limb bones	_	_	_	_	-	4	_	_	_	4	1	_	_	_	1	_		_	_	_	3	_	1	1	5	
alvarium, mandible and limb bones	_	_	4	1	5	_	_	3	1	4	1	_	2	1	4	_	_	_	_	_	5	1	7	3	16	
alvarium, pelvis and limb bones	1	_	2	_	3		_	_	_	_	_	_	_	_	_	_	_	_	_	_	3	_	1		4	
alvarium, mandible, pelvis and limb bones	5	_	5	1	11	5	·	4	_	9	-	_	_	_	· —	-	_	_	_	_	5	_	5	1	11	
Mandible	_	_	_	_	_	_	-	-	_	_	1	_	_	_	1	_	_	-	_	-	1	2	_	-	3	
Mandible and limb bones	_	, —	_		_	}	_	_	-	-	_	_	_		_	_	_	_	_	_	_	_	2	_	2	
Mandible, pelvis and limb bones	_	_	1	_	1	-	_	_	1-	_	_	_	_		_	_	_ ,	_	_	_	1		_	_	1	
Pelvis			_		-	_			_	_	1	-	1	_	2		_	_	_	_	_	_	2	1-7	2	
Pelvis and limb bones		_	_			_	_		_	-		_		_	5	_	_	_	_	_	_1	-	4	-	5	
Limb bones	_	2		2	4	1	. T			1	1		1	_	2	-			2 7		5	1	- 11	2	. Ì	
Total	9	2	15	4	30	11		7	1	19	6		4	2	12		8			13	33	 5	46	9	93	

^{*} Skeletons which could not be sexed by anatomical appreciation have been omitted.

SQUARE (CEMETERY) R 37

The cemetery in Square R 37 at Harappā, located on a slightly raised ground to the south of the main habitation-area brought to light the only regular cemetery of the inhabitants of Harappā belonging to the true Harappān period. Altogether, skeletons of 108 individuals were recovered from fifty-seven burials of the cemetery in different seasons between the years 1938-41 and 1946-47.

Of these graves forty-seven were exposed in earlier seasons and the remaining ten in 1946, short notes on the burials of the latter ones have been furnished by Mr H. K. Bose (Wheeler 1947; 86-89). The graves exposed in earlier seasons, however, have not been described previously, and, therefore, short notes on them are presented below:

- 1 Burial No. H. 779 and H. 783: Multiple burial; North South orientation; laid in supine position; head turned towards the East; No. of skulls five; bones generally in decayed condition.
- 2 Burial No. H. 780: Fractional burial; No. of skull one.
- 3 Burial No. H. 781: Fractional burial; North South orientation; No. of skull one; in extremely decayed condition.
- 4 Burial No. H. 782: Complete burial; North South orientation; laid in supine; head turned to the East; No. of skull one; bones in decayed condition. Two animal ribs are on each side of the head.
- 5 Burial No. H. 791: Fragmentary skeleton; North South orientation; laid in supine; bones in extremely decayed condition.
- 6 Burial No. H. 791: Fragmentary skeleton; North South orientation; laid in supine; head turned towards East; No. of skull one; bones generally in decayed condition.
- 7 Burial No. H. 792: Fractional burial; bones in decayed condition.
- 8 Burial No. H. 793: Complete burial; North South orientation; laid on left side; head facing East; No. of skull one; bones in distorted condition.
- 9 Burial No. H. 793/A: Complete burial; North South orientation; lay prostrate; head facing downward; large skull one in number; bones generally well preserved.
- 10 Burial No. H. 793/B: Fractional burial; head tilted on right cheek; No. of skull one; bones in fair condition.
- Burial No. H. 794: Complete burial; North South orientation; laid on right side; head facing West; No. of skull one; well preserved condition. Animal rib near foot.

- 12 (i) Burial No. H. 794/A: Complete burial; North South orientation; laid on supine; head tilted westward; No. of skull one; bones in decayed condition; arm bone of another individual present.

 (ii) Burial No. H. 794/B: One piece of human bone only represented this burial.
- 13 Burial No. H. 795: Fractional burial.
- 14 (i) $Burial\ No.\ H.\ 795/A$: Complete burial; North South orientation; laid on right side; head tilted westerward; No. of skull one; bones in poor state of preservation.
 - (ii) Burial No. H. 795/B: No bones found; treated as part of H. 795/A.
- 15 Burial No. H. 796: Fractional burial; comprised three fragments of long bones only.
- 16 Burial No. H. 796/A: Fragmentary skeleton; North South orientation: laid in supine; this burial treated as a part of H. 796.
- 17 Burial No. H. 796/B: Fractional burial; head turned on cheek; No. of skull one; bones in fair preservation. Burial Nos. H. 796, H. 796/A, H. 796/B could form a single burial.
- 18 Burial No. H. 797: Fractional burial; only one piece of long bone represented in this burial.
- 19 Burial No. H. 798: Multiple burial; No. of skulls six. Animal bones present.
- 20 (i) Burial No. H. 798/A: Fragmentary skeleton; North South orientation; laid in supine; head resting on frontal region; No. of skulls two.
 - (ii) Burial No. H. 798: No bones found. Burial Nos. H. 798/A and H. 798/B treated as one burial.
- 21 Burial No. H. 799: Fractional burial; North South orientation; laid on left side; head facing East.
- 22 (i) Burial No. H. 800: Fragmentary skeleton; North South orientation; laid in supine.
 - (ii) Burial No. H. 800/A: Represented by animal rib only.
 - (iii) Burial No. H. 800/B: No bones found. Burial Nos. H. 800, H. 800/A and H. 800/B treated as a single burial.
- 23 Burial No. H. 801: Fractional burial; only two pieces of bones found.
- 24 Burial No. H. 800/A: Complete burial; North South orientation; laid in supine; head lay detached from the body; No. of skull one. Animal rib present.
- 25 Burial No. H. 801/B: Multiple burial; No. of skulls two. Two animal bones present of which one was identified as cattle horn.
- 26 Burial Nos. (i) H. 802 and (ii) H. 802/A: Fractional burial; comprised a large number of big bones. The latter burial had a fragment of a skull.
 (iii) Burial No. H. 802/B: Fractional burial; only a few pieces of human bones represented. Burial Nos. H. 802, H. 802/A and H. 802/B treated as a single burial.
- 27 Burial No. H. 803: Fractional burial; North-Northeast to South-Southwest orientation; laid on right side; head facing west; bones in extremely fragmentary condition. Animal bone present.
- 28 Burial No. H. 803/A: Fractional burial; only two pieces of bones and a piece of a skull present.
- 29 Burial No. H. 804: Complete burial; North South orientation; laid on right side; head facing West; only part of a skull survived.
- 30 Burial No. H. 805: Fractional burial; head tilted: No. of skull one; bones in fair preservation.

- 31 Burial No. H. 805/A: Fractional burial; North South orientation; laid prostrate; fragentary skeleton; piece of skull only.
- 32 Burial No. H. 806: Complete burial; North South orientation; laid supine; head tilted slightly eastward; one well preserved skull; bones in extremely decayed condition. Animal ribs and two fragments of leg bones of another person found.
- 33 Burial No. H. 806/A: Complete burial; North South orientation; laid in supine; head facing East; No. of skull one.
- 34 Burial No. H. 807: Fractional burial.
- 35 Burial No. H. 808: Fractional burial; North South orientation; laid in supine; fragmentary skeleton.
- 36 (i) Burial No. H. 310: Complete burial; North South orientation; laid in supine; head facing East; No. of skull one; bones in extremely decayed condition.
 - (ii) Burial No. H. 810/A: No bones found. Burial H. 810 and H. 810/A treated as a single burial.
- 37 (i) Burial No. H. 811: Fractional burial; head tilted on right cheek; No. of skull one.
 - (ii) Burial No. H. 811/A: No bones found, Burials H. 811 and H. 811/A treated as one burial.
- 38 (i) Burial No. H. 812: Fractional burial; North South orientation; laid in supine; head facing East; No. of skull one; bones in decayed condition.
 - (ii) Burial No. H. 812/A: No bones found. Nos. (i) and (ii) treated as one burial.
- 39 (i) Burial No. H. 813: Fragmentary body; North South orientation; laid in supine; head facing East.
 - (ii) Burial No. H. 813/A: No bones found. This burial treated as part of burial H. 813.
- 40 Burial No. H. 814: Fragmentary skeleton; North south orientation; laid in supine; lower jaw pointing eastward; No. of skulls two. Some stray human bones found.
- 41 Burial No. H. 815: Fragmentary skeleton; North South orientation; laid in supine.
- 42 (i) Burial No. H. 816: Fragmentary skeleton; Southeast to Northwest orientation; laid prostrate; head facing downward; No. of skull one. Bones of other individual present.
 - (ii) Burial No. H. 816/A: No bones found. This burial treated as part of H. 816.
- 43 Burial No. H. 817: Complete burial; North South orientation: laid in supine; head facing skyward; No. of skull one; bones in extremely decayed condition. Bones of bird present.
- 44 Burial No. H. 318: Complete burial; North South orientation; laid in supine; head facing sky-ward; bones well preserved.
- 45 Burial No. H. 819: Complete burial; North South orientation; laid in supine; head facing eastward; No. of skull one; bones mostly decayed condition.
- 46 Burial No. H. 820: Multiple burial; North South orientation; laid on left side; head facing eastward; No. of skull three.
- 47 Burial No. H. 821: Complete burial; North South orientation; laid in supine; head slightly tilted eastward; No. of skull one; bones in partly decayed and partly well preserved condition.

Out of 47 burials 15 were complete, 4 multiple and 15 fractional comprising collections of dismembered bones.

MATERIAL

The collection comprises materials varying from complete bones of individual skeleton to only a few bits representing one individual. Only thirty-six well-preserved crania and mandibulae of Cemetery R 37 have been utilised for osteological study and report. These thirty-six individuals represent 15 adult males, 19 adult females and 2 juveniles, but no children. While non-adults are not well-representated, as it ought to have been in a regular cemetery, children's skeletons are conspicuously absent. The age and sex distribution of the material, which is detailed below, reveals an interesting fact that, with rare exceptions, the majority of them died when they were at the prime of their life. Only one woman (H 805) seems to have enjoyed a long life.

TABLE 5
SEXWISE AGE DISTRIBUTION OF CEMETERY R 37 SKELETONS

AGE	JUVENILE	18 - 21 YEARS	21 - 25 YEARS	25 - 30 YEARS	30 - 40 YEARS	40 - 50 YEARS	50 - 60 YEARS
MALE	H 801 (B)	_	H 820 (iii)	H 779 (e), H 793, H 794, H 796 (B), H 798 (A), H 798 (B) H 798 (C), H 806, H 793 (B), H 811, H 818	Skl. 10, H 793(A)	Skl. 1	
FEMALE	Н 798/А2	Н 817	H 795/A	H 779 (a), H 804, H 788, H 806 A, H 812, H 816, H 820 (II), Skl. 2, H 791/A, H 798/A I	H 780, H 798 (a), H 779 (c), H 801/A	_	H 80 5

Age of two women H 779(d) and H 820(l) could not be estimated but they undoubtedly were adults.

A. CRANIA


Detailed descriptions of individual skull of Cemetery R 37 have been furnished below. The measurements of skulls and extremity bones are produced in the Collective Tables appended at the end. Juveniles have not been taken into account for the determination of physical types, but their measurements and observations have been incorporated in the Collective Tables (A—B and G—H).

The skull with mandible belongs to an adult female of about 30 years of age. The skull is very badly preserved and is warped. Part of frontal and parietal bones of both sides are missing. Mastoids are small and the muscular ridges are not developed. Contour in norma verticalis (Sergi 1899: 20) appears to be elliptical and the side walls are nearly flattened. Basilar suture is united and the vault sutures



show indications of beginning union. The skull is long (182.0 mm) and medium in height, the length-auricular height index being 60.71 (orthocranic). The orbits tend to be rectangular and are mesoconchic (right OI 84.44? and left OI 79.07). Nasal aperture is narrow and the nasal index lies at the border of leptorrhine and mesorrhine (NI 46.94). Palate is deep and paraboloid in shape. All the maxillary teeth are fairly intact, only crowns of upper left M₂ and M₃ are partially broken.





Mandible was broken and had to be repaired before measurements. The lower dental arcade is narrow. The width of the dental arcade, measured between the outer margin of the second molars, is 51 mm. The depth of the symphysis is 36 mm.

SKL. 779 (c) [Fig. 5]

Skull is that of an adult female approximately 35 years of age. It is in a very poor state of preservation. Right side of the cranium has disintegrated producing granular surface. Supraorbital ridges are not perceptible, mastoids small, and muscular ridges are only moderately developed. The basilar suture is closed, while the coronal and the sagittal have started closing. Forehead is narrow, high and vertical with almost parallel side-walls. Viewed from above the skull is elliptical in shape. From the side, the vault appears to be high with flattened temples. Occiput is protruding and shelved abruptly downwards and forwards from the superior nuchal line. The skull is dolichocranic (L-B Index 66.30?) and acrocranic (B-H Index 106.56?). The orbits are rectangular, being hypsiconchic at right (OI 85.00) and mesoconchic at left (OI 82.05). The calculated cranial capacity of the skull is 1198.35 c.c. The depression at nasion is medium and the nose falls in chamaerrhine class (NI 53.33). The palate is short, wide and parabolic. Maxillary teeth are all relatively small sized.

Lower jaw is not preserved.

skl. H 779 (d)

The skull is badly damaged; the calva, excepting the basicranii and part of the frontal bone, is missing. The skull is that of an adult female; the supraorbital ridges are not marked and the nasion

is only slightly depressed. The existing right orbit tends to be rectangular and mesoconchic (OI 80.49). A few measurements could be taken on the skull. Three molar teeth of each side of the maxilla are present.

No mandible.

SKL. H 779 (e) [Figs. 3, 4, 6]

The skull appears to be that of a male in the prime of life. The skull is complete without mandible, only the posterior portion of the skull vault is broken. Supraorbital ridges are prominent, having large mastoids and well-developed muscular impressions. Basicranial synchondrosis has completed its ossification, while ectocranial sutures are open. The forehead is receding and slopes back posteriorly in a low but regular curve upto the inion, and the nuchal region shelves abruptly downwards and forwards. The occiput is markedly protruding (occiput en chignon) and retreats sharply from inferior nuchal line. On the whole the skull is long, narrow and moderately low, having almost






Fig. 4

parallel side-walls. The skull is hyperdolichocranic (L-B Index 68.06) and on the border of orthocranic and chamaecranic (L-AH Index 58.90). Glabella is well developed and the nose is sunken at the root. The nose is concave, long and narrow, being leptorrhine (NI 41.51). The inferior border of the nasal aperture is amblykraspedotic. The nasal bones are wing-shaped with medium depression at the root. Vertical contour is somewhat ellipsoides in shape and phaenozygous. The subnasal is only moderately prognathous. The palate is short, deep and upsiloid in shape. All the maxillary teeth have cut but the incisors and left canine have fallen out.

SKL. H 780 [Figs. 7, 8, 11]

The skull is that of an adult female between 30 and 40 years of age at the time of death. The skull is in a fairly good state of preservation and complete excepting for the right zygomatic arch, tip of the left mastoid process, styloid process and squama of the right temporal. It is without mandible. Skull is smooth, gracile with medium supraorbital ridges and small mastoids. Muscular impressions are ill marked. Subnasal part of the maxilla is directed forward presenting a mild degree of subnasal




Fig. 5 Skl. H 779 (c)





Fig. 6 Skl. H 779 (e)

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

prognathism. Seen from above, the outline of the skull is *pentagonoides* in shape. The skull is submesocranic (L-B Index 75.69) and orthocranic (L-AH Index 61.88). The forehead is medium in height and is slightly receding. The vault appears to be well arched with smooth contour line and the occipital





region is moderately protruding. The orbits are low, rectangular with rounded angles and are meso-conchic (right OI 76.19 and left OI 79.52) in form. The nasal bones are wing-shaped with moderately depressed nasion and slightly concave nasal bridge. The nose is mesorrhine (NI 47.92), the inferior border of the pyriform aperture is oxykraspedotic. The teeth of the right maxilla, excepting PM₁ and PM₂, are intact, while left is devoid of lateral incisor and canine which have fallen out post mortem. The upper dental arcade is upsiloid and the palate is deep.

SKL. H 788

The skull which is in a very poor state of preservation belongs to an adult female. Unfortunately, the skull is compressed and very badly damaged; the entire vault of the left half and basicranii is missing, thus very few measurements could be taken on it. Some portion of face and palate are is missing. Mastoid is small, the supraorbital ridges are absent and superior orbital margins are sharp. Surviving. Mastoid is not prominent and there is only little depression at the nasal root. Excepting for the Glabella is not prominent and there is only little depression at the nasal root. Palate is deep and central incisors, which have fallen out, all the other maxillary teeth are intact. Palate is deep and paraboloid. Basilar suture is united, while the vault sutures are open.

Mandible is broken. The right half of the mandible bears all the teeth but the left one has the incisors and canine.

SKL. H 791/A

The skull belongs to that of an adult female. It is smooth-contoured and gracile, having small mastoids with weak supraorbital ridges and prominent frontal eminences. The facial portion is damaged and distorted, particularly the right side. Right maxilla and zygoma are pushed backwards and inwards, due probably to some heavy pressure, while the left side also received the impact of the pressure of earth. Basilar suture is closed and coronal and sagittal sutures have just started

1

closing, while lambdoid suture is open. Contour of the norma verticalis is ovoides with moderately projecting occiput. The forehead is medium and vertical. The vault is somewhat low. The skull is dolichocranic (L-B Index 72.99), orthocranic (L-AH Index 59.77) and metriocranic (B-AH Index 81.89). Only the right orbit could be utilised for measurement, it being hypsiconchic (OI 91.67?).

The mandible is found in pieces having slightly absorbed margin with some of the teeth intact.


SKL. H 793

The skull is that of an adult male. It is not in a good state of preservation and had to be restored. The chief defects are as follows: the greater part of the facial region, part of the basis cranii, little of the left parietal and places around both the pterions are missing. The supraorbital ridges are prominent and the skull shows strong musculature. Norma verticalis is byrsoides in outline, narrowed in front and expanded at the parietals. From the side, the forehead is retreating and passes into a faintly arched vertex: the occiput is moderately protruding with a sharp retreat at the nuchal region. The glabella is prominent and the nose is sunken at the root. The skull is hyperdolichocranic (L-B Index 68.91) with somewhat laterally compressed side-walls, chamaecranic (L-AH Index 57.51?) and metriocranic (B-AH Index 83.46?). All the right maxillary teeth are intact. At the left PM2 and M3 are present. The cubic capacity of the cranium is euencephalic i.e. 1375.33 c.c.

The mandible which was obtained in fragments is restored, whose left coronoid is missing. All the mandibular teeth numbering sixteen are found intact in their sockets and teeth are only moderately worn down. Chin is prominent, corpus massive and maximum width of the ascending ramus is 44.0 mm.

SKL. H 793 (A) [Figs. 9, 10, 12; Pl. XIX]

The skull is massive in structure and belongs to an adult male between 30 and 35 years of age. It had to be restored before measurement. In absolute dimensions of certain characters the skull







Fig. 11 Skl. H 780



MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

exceeds all the other skulls in the whole of the Harappān series. The cubic capacity of the cranium is very large (1605.68 c.c.). The skull is incomplete; a considerable portion of the basis cranii and all the maxillary incisors and right canine are missing. The norma verticalis is wide ovoid with marked parietal eminences. The inion is very prominent and the muscle attachments in the occipital region are medium. The norma occipitalis is broad, house-shaped with a faintly convex roof. The glabella is prominent and depression at the nasal root is medium. In profile, the forehead ascends obliquely, passes back into a low arched vertex, while the occiput is only slightly convex with no actual protrusion. The skull is on the upper border of mesocrany (L-B Index 79.79), hypsicranic (L-AH Index 63.52) and tapeinocranic (B-AH Index 79.61). The superior facial index is 47.87? (euryprosop) and the nose is mesorrhine (NI 47.79).


Corpus mandibulae is in a fairly good state of preservation, whereas greater portion of both the ascending rami is not preserved. All the teeth have erupted but incisors lost post mortem. Teeth are all in much worn state. Chin is prominent, lying 5 mm in front of the lower alveolar point. The vertical depth of the horizontal ramus at the region of second molar is 30 mm.

The skull is that of an adult male of about 30 years of age. Zygomatic arches, occipital condyles and styloid processes are broken and missing. Both the parietals are eroded. Supraorbital ridges are absent but the mastoids are strong, having moderate muscular impressions at the nuchal region. Vault sutures are open, but synchondrosis sphenobasilaris is ossified. Vertical contour is oval in shape which is narrow in front and slightly expanded at the parietals. Seen from the side, the forehead rises vertically and slopes back posteriorly and the occiput bulges well behind. The vertex is of medium height (vertical porion height being 116.0 mm) and is slightly flattened. The skull is dolichocranic (L-B Index 70.43), orthocranic (L-AH Index 62.37) and acrocranic (B-H Index 98.47). Nasal profile is concave and not sunken at the root. All the maxillary teeth have erupted but many of them have lost post mortem.

The lower jaw is incomplete. The left ascending ramus is broken off immediately behind M_3 . All the teeth have erupted but incisors, right canine and right PM_1 are missing. Chin is only moderately developed and mylohyoid ridge is not very distinct.


SKL. H 794 [Fig. 14; Pls. V, VI, XX]

The skull belongs to an adult male of about 30 years of age. The general state of preservation of the skull is not very good. The ectocranial surface, particularly the left lateral side, is eroded and small portion of both the parietals are missing. The glabella is marked, superciliary arches well developed, and mastoids large. When viewed from the top, the skull is byrsoides in shape and the zygomatic arches are bowed out laterally from the vertical contour. In profile, the forehead is little inclined and the slope passes backwards into the occipital squama, which projects beyond the inion. There is also evidence of some degree of subnasal prognathism, and pre-auricular length of the skull is slightly greater than post-auricular length. The skull is delichocranic (L-B Index 72.14) and orthocranic (L-AH Index 61.72). The nasion is depressed at the root and the nasal index lies on the border



Norma Frontalis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 793 (B) (Natural Size)



Norma Lateralis




Fig. 13 Skl. H 793 (B)




Fig. 14 Skl. H 794

 $\textbf{MID-SAGITTAL CURVE} \hspace{0.1cm} (\ ^{1}\!/_{\!\!2} \ \ \text{NATURAL SIZE} \hspace{0.1cm})$

between mesorrhine and chamaerrhine (NI 50.94). The calculated cranial capacity of the skull is 1481.60 c.c., i.e. aristencephal according to Sarasin's classification and megacephal according to Flower and Turner's classification.

The lower jaw is fragmentary in nature. All the teeth have erupted but the right lateral incisor is found to be missing from the socket. The crowns of the teeth have considerably worn down. The chin is well developed, lying 6 mm in front of the lower alveolar point, corpus massive and ramus broad. The depth of the symphysis is 40.0 mm and maximum breadth of the ascending ramus is 44.5 mm.

SKL. H 795/A

This is an adult skull probably of a female. The skull is in badly warped condition. All the sixteen maxillary teeth are intact and worn well through the dentine. Supraorbital ridges are not developed and the mastoids are medium in size. From the top of the skull contour is ovoid with some amount of bulging at the parietals. Forehead is receding and of medium breadth. The subnasal height is medium with some amount of prognathism. The skull is long and the vault of the skull is low. The orbits are rounded and hypsiconchic (right OI 89.47 and left OI 93.24). The nasal aperture is narrow, moderately high and mesorrhine (NI 48.42); and the inferior border of the pyriform aperture is oxykraspedotic. The palate is long, narrowed in front and parabolic in shape.

Mandible had to be restored to a considerable extent before measurement. The jaw is of moderate size and the muscles of mastication are not well developed. Chin is prominent but corpus is moderate in bulk. Surface of molar teeth shows considerable attrition. The mandibular measurements* are as follows:

Ht. of mandibular ramus	60.0 mm.		
Ht. at mandibular symphysis	27.0 mm.		
Mandibular length	77.0 mm.		
Max. breadth of mandibular ramus	43.0 ? mm		
Min. breadth of mandibular ramus	31.0 mm.		
Mandibular angle	126°		

SKL. H 796 (B) [Fig. 18; Pis. VII, VIII]


It is a complete locked-jaw skull belonging to that of an adult male between 25 and 30 years of age. The skull is warped and both the malar bones have eroded considerably. Right zygomatic arch and styloid processes are missing. All the sixteen pairs of maxillary and mandibular teeth are intact and well set in position in clenched condition, excepting the upper right central incisor. Almost all the teeth exhibit vertical cracks. Basilar suture is united while the coronal, sagittal and lambdoidal sutures are open. The supraorbital crests tend to be well developed with sharp muscular attachments. Other indications of muscularity are strong temporal lines. The skull is sub mesocranic (L-B Index 75.84), hypsicranic (L-AH Index 64.04) and metriocranic (B-AH Index 84.44). The

^{*} Detailed measurements are presented here since the specimen was not correctly mended when the Collective Tables were sent to press.

orbits are hypsiconchic (right OI 92.31 and left OI 95.00). The nose is chamaerrhine (NI 52.34) and the face is long oval. The intracranial capacity is 1336.67 c.c.

The skull is that of an adult male. It is compléte except a gap in front of the coronal suture on the left frontal bone. Supraorbital ridges are very highly developed and the muscular impressions at the nuchal region are remarkably prominent forming an occipital torus. Left mastoid is strong and big, right is missing and the nasion is sunken. Forehead is receding and the sagittal contour is not so well arched and the occiput while fairly prominent, retreats sharply downwards and forwards. Face is medium in height and moderately broad, the upper facial index being on the border line of lepten—mesen (lepten 55.60). The palate is narrow and upsiloid in shape. All the





maxillary teeth have erupted and are intact excepting the left M_3 ; the cusps show considerable wearing. The degree of wear of incisors definitely suggests an 'edge-to-edge' bite. There appears to have some degree of lambdoid flattening, probably due to the presence of occipital 'bun'. When seen from above the skull is byrsoides in outline and expanded at the parietals. Orbits are low and rectangular, inclined laterally downwards and are both chamaeconchic (OI 73.49). The masal aperture is wide and short, and consequently chamaerrhine in nose form (NI 53.00). The margo pyriformis apertura is amblykraspedotic. Subnasal prognathism is evident. Palate is paraboloid in shape, deep and somewhat narrow anteriorly. The skull is hyperdolichocranic (L-B Index 69.31) and orthocranic (L-AH Index 59.79). The cranial capacity of the skull is 1357.20 c.c.

Probably female skull between 30 and 40 years of age; sutures are unobliterated and teeth are all in a very much worn state. There is no lower jaw. The mastoid processes, muscular attachments and temporal curved lines are moderate. The vault of the skull is moderately low as shown by the





Fig. 17 Skl. H 798 (A)




 $\textbf{MID-SAGITTAL CURVE} ~(~^{1}\!\!\!/_{\!\!2}~\text{NATURAL SIZE}~)$


vertical porion height (111.5 mm). The orbits are low, rectangular and inclined laterally and downwards and the maxillo-frontal orbital index lies marginal to the middle class (OI 76.83 both). The skull is dolichocranic (L-B Index 72.47) and orthocranic (L-AH Index 62.64) and very near to acrocrany (B-AH Index 86.43). Outline of the norma verticalis is ovoides. The calculated cubic capacity of the skull is 1254.88 c.c. or distinctly euencephalic.

The glabella and supraorbital ridges are prominent and from the side, the forehead appears to be slightly receding. The sagittal contour is not so well arched and the occiput is rounded with a house-shaped outline of norma occipitalis. The nasion is moderately depressed at the root, having broad constricted nasal bone with concave bridge. The upper face (nasion-prosthion) is 66.0 mm long, bizygomatic width 127 mm, and there is also some amount of subnasal prognathism. The dental arcade is upsiloid, the palate being short and broad.

SKL. H 798/A1 [Figs. 19, 20, 22]

The skull is typically female; small, smooth, gracile with weak supraorbital ridges and muscular impressions are feminine features. It is complete and in an excellent state of preservation and approximately 30 years of age. When seen from above the skull is sphenoides in outline and phaenozygous. Forehead rises vertically and then sweeps back in a regular curve to meet the rounded occiput. The orbits are large and squarish. The skull is brachycranic (L-B Index 80.70) which is nearer to upper margin of mesocranic class and hypsicranic (L-AH Index 64.91). The nasion is only slightly depressed, with a concave bridge formed by wing-shaped nasal bones. The inferior border of the nasal aperture is oxykraspedotic. Dental arch is upsiloid in shape. There are thirteen teeth in the sockets of the maxillae, the crowns of which are deeply worn. The upper wisdom teeth are not erupted and right M₁ lost ante mortem.





SKL. H 798 (B)

The skull is incomplete. Mandible, right half of the facial region, right os temporale and whole of the basilar part is missing. The sex appears to be that of a male; the prominent supraorbital




Fig. 21 SKL. H 798 (a)




Fig. 22 SKL. 798/A 1

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

ridges, large left mastoid process and thick bones are masculine features. The age is definitely of the adult class, part of sagittal suture is ossified; the coronal suture is, however, open. Norma verticalis is rather ovoid in shape. In norma lateralis the forehead is low but retreating and the vertex is somewhat flat. The glabella is moderately prominent and nasion depressed at the root. The left orbit is somewhat rectangular in shape and mesoconchal in form, the right orbit is missing.

SKL. H 798 (C)

The skull is distorted and a large part is missing. The missing parts include whole of the frontal region with zygomatic arches and sphenoidal part of the basis cranii. The sex is undoubtedly male, having thick bones, prominent supraorbital ridges, well-developed muscular attachments and big mastoids. When seen from the top, the skull appears to be byrsoides in outline. The frontal bone is evenly sloping and passes back to the slightly domed vertex. The occiput is protruding with a moderately pronounced tuber occipitale. The occipital contour is house-shaped. The skull is hyperdolichocranic (L-B Index 69.68?), orthocranic (L-H Index 72.87) and acrocranic (B-H Index 104.58).

SKL. H 801/A [Fig. 23; Pl. XXVIII: 1-2]

The skull is complete and in a fairly good state of preservation. The ectocranium is granular and pitted. The sex is undoubtedly female. It is gracile, smooth contoured, having ill-developed supraorbital ridges and muscular impressions with small mastoids. Supraorbital notches are present on both sides of the sharp margin. There is a trace of persisting frontal suture. Basilar suture is closed. The skull is delichocranic (L-B Index 71.47) and hypsicranic (L-AH Index 64.67). The forehead is high and nearly vertical to about ophryon where it commences gradually to slope backwards; occiput being moderately protruding. When viewed from above, the skull is elliptical in outline. Parietal eminences are somewhat prominent. The orbits tend to be rectangular, both being hypsiconchic (right OI 86.84, left OI 87.18). Nose is prominent and leptorrhine (NI 45.28), having narrow constricted nasal bones. Nasal profile is only slightly concave with no depression at the root. The face is long and narrow. Alveolar margin is absorbed.

The mandible is moderate in size. The bicondylar breadth is 117.0 mm and the minimum breadth of the ramus shows a medium value (36.0 mm). The mandibular teeth are very badly worn (dentine exposed on cusps). Chin juts out anteriorly.

SKL H 804

The skull is incomplete being that of an adult female. It is survived by only the anterior third with right temporal, masteid processes and mandible. All the teeth of right maxilla, excepting the molars which were fallen out post mortem are intact; at left, excepting the left lateral incisor, PM, PM, and others are all in their position. Teeth show marked attrition (in case of incisors lingual attrition) and widening up of space between roots, particularly in the mandible. The supraorbital ridges are not developed and the masteids are small. The forehead is nearly vertical and the frontal eminences are prominent. The orbits are somewhat squarish, both being hypsiconchic (right OI 87.50)






Fig. 23 SKL. H 801/A




 $\label{eq:mid-sagittal} \textbf{MID-SAGITTAL} \quad \textbf{CURVE} \quad (\ ^{1}/_{2} \quad \text{NATURAL} \quad \text{SIZE} \)$



Norma Frontalis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 794 (Natural Size)



Norma Lateralis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 794 (Natural Size)

and left OI 89.47). The superior orbital foramen is present at the right, while at the left there is a supraorbital notch. The nose is mesorrhine (NI 48.91).

Mandible is complete, though several teeth had been lost during life, and sockets are absorbed. Mandibular teeth are also in much worn state, particularly molars are hollowed by wear. The chin is not prominent to a marked degree and depth of the symphysis is only 24.0 mm. The minimum breadth of the ascending ramus is moderate (30.0 mm).

The skull appears to belong to an adult female fairly old in age. Most of the vault sutures being ossified and molars show considerable wearing. Outer wall of the right orbit, right malar bone and right maxilla are missing. The skull is gracile and smooth contoured, having weak muscular impressions with no supraorbital ridges. Forehead is narrow; it rises vertically and sweeps back in an even curve. The skull is long, narrow with laterally compressed wall-like sides. The skull is markedly dolichocranic (L-B Index 70.78), orthocranic (L-AH Index 62.73) and acrocranic (B-AH Index 88.64). Nasal aperture is moderately short and broad, chamaerrhine (NI 52.94); nasal profile is concave without any depression at the root. The left orbit is hypsiconchic (OI 86.36).

When seen from above, the skull is *ellipsoides* in contour and cryptozygous. The occiput is moderately protuding and the occipital contour is house-shaped; subnasal prognathism is absent. All the maxillary teeth are missing. Mandible is fragmentary in nature and no useful measurements could be taken.

The skull is well preserved and belongs to that of an adult male. Most of the sagittal suture and greater part of the lambdoidal suture being ossified. Supraorbital ridges and mastoids are well and greater part of the lambdoidal suture being ossified. Supraorbital ridges and mastoids are well developed. Oblique cut mark presents at the vertex. In norma verticalis, the outline of the skull is developed. When seen from side, the forehead is receding and passes back posteriorly. Slight degree ellipsoides. When seen at the bregmatic region and the occiput is markedly en chignon. The face is of flattening is seen at the bregmatic region and the occiput is markedly en chignon.




Fig. 25




Fig. 26





Fig. 28 SKL. H 806/A



MID-SAGITTAL CURVE (1/2 NATURAL SIZE)


moderately high and moderately wide, superior facial index being 52.06 (mesen) with flattened temples, while the total face is middle and mesoeuryprosop (88.76). The skull is hyperdolichocranic (L-B Index 69.31), and on the border between orthocranic and chamaecranic (L-AH Index orthocranic 58.47). The nose is broad and relatively short, being chamaerrhine (NI 56.60). Nasal profile is concave and the nose is sunken at the root. The skull shows a considerable degree of alveolar prognathism, with the upper incisors projecting forwards. The palate is *parabolic* and the teeth are somewhat worn down.

The lower jaw is well formed with prominent and slightly everted chin. The corpus is massive and ramus broad, the minimum breadth of the ramus being 36.0 mm.

SKL. H 806/A [Figs. 28, 29, 30; Pl. XXIII]

The skull is in a very good state of preservation. The sex is difficult to determine with certainity; smooth contour, small mastoids and weak muscular impressions rather suggest a female. The vertical contour is pentagonoid in shape, narrow in front and bulging at the parietals. From the sides, the forehead is slightly receding and slopes back posteriorly with a slight flattening in the bregma





region. The occiput is moderately protruding and from the superior nuchal line it projects forward sharply. The skull is delichocranic (L-B Index 71.28) and acrocranic (B-AH Index 86.33). The masal aperture is short and broad, hence chamerrhine (NI 53.19). The submasal space is high with marked aperture is short and broad, hence chamerrhine (NI 53.19). The submasal space is high with marked aperture is short and broad, hence chamerrhine (NI 53.19). The submasal space is high with marked aperture is short and broad, hence chamerrhine (NI 53.19). All the teeth have cut and the prognathism. The palate is long and narrow and parabolic in shape. All the teeth have cut and the cusps show appreciable wearing. Mandible is well preserved, having prominent and slightly everted cusps show appreciable wearing. Mandible is well preserved, having prominent and slightly everted cusps show appreciable wearing. Mandible is well preserved, having prominent and slightly everted cusps show appreciable wearing. Mandible is well preserved, having prominent and slightly everted cusps show appreciable wearing. Mandible is well preserved, having prominent and slightly everted cusps show appreciable wearing. The degree of wear of the incisors definitely suggests that the teeth during life time met in an edge to edge bite (pl. XII: 1).

SKL. H 811 [Fig. 31; Pl. XXIV: 1-3]

The skull is nearly complete. The left lateral surface of the skull is eroded and part of the left orbit, zygomatic arch and left mastoid process are missing. The sex is difficult to determine but appears





Fig. 31 SKL. H 811




MID-SAGITTAL CURVE (1/2 NATURAL SIZE)




Norma Frontalis

DIOPTOGRAPH TRACING (WASH) : MALE SKL. H 796 (B) (Natural Size)




Norma Lateralis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 796 (B) (Natural Size)



Norma Frontalis



Norma Lateralis

to be an adult male. The contour of the norma verticalis is somewhat *ellipsoides*. In norma temporalis supraorbital ridges are slight, the frontal bone slightly inclined and ascends into a smooth curve upto the bregma. The vertex is flat and the occipital area is abruptly sloping with a *chignon*. The nose is moderately prominent (NI 48.15?), bridge concave and nasion only slightly depressed. The skull is dolichocranic (L-B Index 71.74?), hypsicranic (L-AH Index 64.67) and acrocranic (B-AH Index 90.15?). The calculated cranial capacity is 1389.77? c.c., i.e. euencephalic. Only the right orbit, which could be measured, is squarish in form and meosconchic (OI 80.00). The length of the upper face (nasion-prosthion) is 74.0 mm.

SKL. H 812 [Fig. 32]

The skull is not well preserved. Right lateral side and facial region have undergone considerable degree of warping. The skull is typically female. Gracile, smooth contour, ill-marked supraorbital ridges and small mastoids are all feminine features. The head shape, seen from above, is elliptical. Seen from the side, the forehead is slightly receding; it slopes back posteriorly and the occiput is well protruded behind. The roof of the skull is relatively low; the vertical porion height being only 109 mm. The skull is hyperdolichocranic (L-B Index 68.68?) and orthocranic (L-AH Index 59.89). The nasal index is leptorrhine (NI 43.40) and the nasal depression is medium with a concave nasal bridge. The orbits are large and tend to be squarish and are mesoconchic (right OI 84.78, left OI 82.95). The calculated cranial capacity of the skull is 1221.39? c.c., thus places it in the euencephalic class.

SKL. H 816 [Figs. 33, 34, 35; Pl. XXV]

The skull is that of an adult female approximately 30 years of age. Relatively small-sized skull, gracile and smooth contoured, with small mastoids but medium supraorbital development. The face is broad in relation to its height. Spheno-basilaris is closed while coronal and sagittal sutures have just started ossifying. There is a large elliptical gap on the posterior part of the left parietal. The skull is mesocranic (L-B Index 77.84) and hypsicranic (L-AH Index 65.27).





Fig. 33



Seen from above, the skull is byrsoides in outline, being expanded at the parietals: and is phaenozygous. The forehead is slightly receding, and slopes back posteriorly and merges with the rounded occiput which sharply retreats from the superior nuchal line. Orbits are relatively low





Fig. 35 SKL. H 816



MID-SAGITTAL CURVE (1/2 NATURAL SÍZE)

and broad and fall in the mesoconchic class (right OI 82.05 and left OI 76.83). The face is moderately high and moderately wide (the superior facial index 50.78) with a moderate subnasal prognathism. The nasal aperture is wide and low resulting in a chamaerrhine nose (NI 52.58). Nasal bridge is slightly concave and nasion depression is medium.

SKL. H 817

The skull is gracile belonging to a young adult female of about 20 years of age. Almost the whole of the occipital, basicranii and some portion of right parietal are missing. The skull possesses a complete set of maxillary teeth nicely preserved without any wearing of cusps. The third molars did not cut. All sutures are open. From the top, the skull is byrsoides, with some bulging at the parietal eminences, and it is dolichoid. Supraorbital ridges are absent and the nasion is without any depression. The forehead is slightly receding and its breadth is medium. The face is moderately low and of medium breadth, superior facial index being euryn (49.59?) with slight degree of subnasal prognathism. The nose is of medium breadth and height, index being mesorrhine (NI 50.52) and the pyriform aperture is oxykraspedotic. The palate is broad, shallow and parabolic.

Mandible is fragmentary in nature and no useful measurements could be taken.

SKL. H 818 [Fig. 36; Pl. XXIV . 4-6]

The skull belongs to that of an adult male about 25 years of age. The skull is broken and without the lower jaw. Missing parts include, part of the right frontal bone, portion of right parietal, left mastoid process and part of the basis cranii. The vault sutures are simple and show sign of commencing ossification. Supraorbital ridges are feeble, right mastoid process large and nuchal impressions in the occipital bone are well marked. The forehead is slightly inclined; the cranial vault is fairly arched in the fronto-parietal region and slopes backwards and downwards in the parieto-occipital region. The occiput is prominent and projectes behind the inion. The skull is hyperdolichocranic (L-B index 69.15) hypsicranic (L-AH index 62.23) and acrocranic (B-AH Index 90.00). The nose is chamaerrhine (NI 52.08). The nasion is not depressed and the bridge is concave. There are fourteen teeth in the upper jaw. The teeth are all sound and there is no evidence of dental caries. All the teeth (with the exception of 2nd molars) are very much worn and the dentine freely exposed.

The skull belongs to an adult female of about 45 years of age. It is in a fairly good state of preservation; the missing parts are the right zygomatic arch and left mastoid process. All the maxillary teeth are absent and only the stump of right M₁ could be seen. Vertex presents biparietal oval depressions, probably the result of craniotabes. Perforations present in the centre of each parietal depression appear to be due to thining out of the bone and the sharp margins suggest recent occurrence. Right mastoid is small and supraorbital ridges are only slightly developed. Sphenobasilaris is united. Coronal suture on the right side has completely synostosed and the other half is

closing while traces of commencing synostosis is present on the sagittal suture. Other sutures are open. Forehead is almost vertical to about ophryon and then passes backwards. When seen from above the skull is *ovoides*, narrow in front and expanded at the parietals. The occiput is moderately pro-








Fig. 38

truding, the contour being house-shaped. The skull is dolichocranic (L-B Index 71.27) and acrocranic (B-AH Index 87.60). Orbits are low and tend to be rectangular and mesoconchic (OI 79.49) on the left. Nasal aperture is very broad and low. The upper alveolar margin is broken.

No mandible is preserved.

SKL. H 820 (II)

The skull is that of an adult female, probably between 25 and 30 years of age. The supraorbital ridges are moderately developed and the nasion depression is shallow. The forehead is low, slightly receding and the slope is gradual with some degree of flattening at the bregma. From the top, the skull is byrsoides in shape with bulging at the parietals. The orbits are low and rectangular (OI 76.09). The face is low and broad with a marked subnasal prognathism. The nasion is not sunken at the root and the inter-orbital breadth is broad (18.00 mm). The nasal-bridge is concave and the shape of the nasal bone is broad constricted. The inferior border of the pyriform aperture is amblykraspedotic. Palate is long, narrow and paraboloid. All the maxillary teeth have erupted and the cusps are badly worn down.

The mandible is moderately built with moderately prominent symphysis menti. The mandibular teeth are badly worn. Marked attrition of the incisors clearly states that the teeth during life met in an edge to edge bite (Pl. XI: 3). The mandibular angle is 120° and the bigonial width 87 mm.

SKL. H 820 (iii) [Fig. 40]

The skull belongs to an adult male between 21 and 25 years of age at death. Portion of left frontal and left parietal are reconstructed, while left mastoid is broken at the tip. It is large with big mastoids, well-developed supraorbital ridges, strong muscular attachments and in an excellent state of preservation. Intracranial capacity is 1400.84 c.c. Nose is depressed at the root and the space



Fig. 1 Skl. H 806/A




Fig. 2 Skl. H 815



Fig. 3 Skl. H 820 (H)






Fig. 4 Skl. 2

MANDIBLES SHOWING ATTRITION OF INCISOR TEETH



Norma Frontalis

DIOPTOGRAPH TRACING (WASH) : MALE SKL. 10 (Natural Size)



g n

Fig. 39 SKL. H 820(1)

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

between the two orbits is 20 mm. Viewed from the top, the skull is byrsoides in outline, and the arcus zygomaticus is bulged out from the contour line. The forehead is receding and passes back into an uniform curve. Auditory meatus lies nearly midway between the nasion and inion. The vault of the skull presents a slight trace of sagittal keel (scaphocephaly). Occiput moderately projects out. Face is very broad in comparison to its height, the superior facial index is on the margin of hypereuryn—euryn (44.20). The orbits are low and rectangular and lie within the chamaeconchal class (right OI 66.67 and left OI 75.90). Nose is wide (chamaerrhine, NI being 54.17), and the pyriform aperture is oxykraspedotic. The skull is dolichocranic (L-B Index 72.83), hypsicranic (L-AH Index 64.40) and acrocranic (B-AH Index 88.43). Palate is short, broad and parabolic. The width of the palate at the second molar alveoli is 43 mm and the palatal length is 44 mm. All the teeth have erupted but the incisors, canine and left PM₁ have fallen out; the cusps are badly worn down. Subnasal prognathism is present.

The skull is without mandible.

SKL. 1 [Figs. 41, 42, 43]

The skull is well preserved. It is typically male, having well developed glabello-supercillary region, large mastoids and the nuchal impressions in the occipital bone are strongly marked. The face is broad and low, and the malar bones are strongly developed. In profile, the forehead is low, slightly retreating and ascends gradually into a somewhat low vault; the occiput recedes obliquely with a moderate projection. A portion of the roof of the skull shows keeling of the vault (scaphocephaly) sagittaly with a para sagittal depression. The skull is dolichocranic (L-B Index 74.47), orthocranic (L-AH Index 61.17) and metriocranic (B-H Index 93.57?). The nasal bridge is long, concave with slight aquilinity of nose and the nasion is deeply depressed at the root. There is also evidence of subnasal prognathism and the forward growth of the facial region from the meatus is marked. Posterior part of the alveolar margin of the maxilla is absorbed. The calculated cranial capacity of the skull is 1437.82 c.c. i.e. distinctly euencephalic.




Fig. 41




Fig. 42

Mandible which has survived in pieces had to be restored. Posterior part of the alveolar margin of the mandible shows absorption. The mandible is strongly built with promient and everted chin.

The corpus is massive and ramus broad. The muscular attachments for the masseter and internal pterygoid are distinct. The angle formed by the ascending ramus with the body of the mandible is 118 degree.

The extremity bones are powerful in structure with strong muscular attachments. The stature is about 182.90 cm, estimated according to Dupertuis and Hadden's 'general formulae'.




Fig. 43 SKL. 1

The skull is that of an adult female of about 30 years of age. The skull is complete excepting for the part of left parietal and left temporal bones. Left anterior wall of the frontal sinus is broken. Supraorbital ridges are feeble, mastoids small and the glabellar region is broken. The cusps of the teeth show considerable attrition, particularly the degree of wear of both mandibular and maxillary incisor teeth suggests definitely an 'edge to edge' bite. Seen from above, the contour of the norma verticalis is elongated oval in outline. Seen from the side, the forehead is low and the slope is gradual; is elongated oval in outline. Seen from the side, the forehead is low and the slope is gradual; occiput is moderately protruding. The skull is delichocranic (L-B Index 72.19) and orthocranic (L-AH occiput is moderately protruding in mesoprosopic (89.26), and superior facial index is meson (52.07); lndex 59.55?). Total facial index is mesoprosopic (89.26), and superior facial index is meson (52.07); thus the face is medium in height-breadth ratio. The right orbit is rectangular in shape sloping laterally and downwards (mesoconchic OI 83.12), while left is distorted. The nasal aperture is broad and medium and chamaerrhine (NI 54.35). The nasal bridge is high and prominent.

Mandible is narrow and compressed at the gonial angles, its bigonial width being 70.0 mm. Corpus is moderate in bulk and ramus moderate in width. All the teeth have erupted excepting left M_{\odot} , which appears to have been newly cut. The lower dental arcade is upsiloid in shape.




Fig. 44 SKL. 2




Fig. 45 SKL. 10

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

SKL. 10 [Fig. 45; Pls. II, XII, XXVII, XXVIII : 5-6]

The skeleton was found inside a mud brick enclosure, a practice observed by Hargreaves (1935: 26f) at Nal in South Baluchisthan also.

The skull is that of an adult male of about 40 years of age. It is complete with mandible and practically devoid of post mortem defect. The skull is massive and rugged. The glabella and supra-orbital ridges are well marked, muscular impressions at the nuchal plane are exuberently developed with a very prominent inion and the mastoids are quite large. When seen from top, the cranium is a longish oval, well-filled in the temporo-parietal region. From the side, the forehead is low and retreating. It passes back into a slightly arched vertex producing small pre-lambdoid flattening. The roof of the cranium also present a longitudiual sagittal keel. The occiput is well protruded behind and stepped abruptly downwards and forwards in its lower half. The skull is dolichocranic (L-B Index 71.81), orthocranic (L-AH Index 61.17) and B-AH Index lies at the border between metriocranic and acrocranic (85.19). Orbits are large, somewhat rectagular and mesoconchal in form (right OI 82.95, left OI 83.33). The nose is well arched, bridge being slightly concave and nasion is deeply sunken at the root. The face is moderately prognathic due to the subnasal projection.

The mandible is powerfully built with a prominent chin and broad ramus. All the mandibular teeth have erupted and show considerable attrition, but the third molar is absolutely unworn.

The stature is estimated about $190.00~\mathrm{cm}$ (Dupertuis and Hadden). The extremity bones are massive with strong muscular attachments.

ANALYSIS

Analysing in detail the metrical and morphological descriptions (Collective Tables A-B and G-H) of the cranial materials from Cemetery R 37, it is revealed that at least two physical types are found to have been present in the population. For the sake of convenience and in order to avoid confusion as to racial nomenclatures, these two have been termed by us as Type A and Type A. Both the types, A and A_1 are long-headed in character.

Of thirty-six skulls of Cemetery R 37, thirty-one adult skulls are capable of being classified. Of these, twenty-one adult skulls are attributable to Type A and ten adult skulls to Type A. Of the remaining five skulls, two non-adults and two defective adult female skulls [H 804 & H 779 (d)] have not been considered, while one adult male skull H 793 (A) could not be attributed to either of

the types because of its excessive massiveness, broad head and large dimensions of certain metric characters. Type-wise and sex-wise distribution of adults is tabulated below:

7	TYPE A	TYPE A			
MALE	FEMALE	MALE	FEMALE		
Н 779 (е), Н 793,	H 779 (a), H 780,	Н 793 (В), Н 811,	H 779 (c), H 791/A,		
Н 794, Н 796 (В),	Н 788, Н 795/А,	Н 818	Н 798/А1, Н 805,		
H 798 (A), H 798 (B),	H 798 (a), H 806/A,		Н 820 (I), Н 801/A,		
H 798 (C), H 806,	Н 812, Н 816,		Н 817		
H 820 (iii), Skl. 1,	H 820 (II), Skl. 2				
Skl. 10					
			OK A STANKER THE		
Total 11	10	3	7		

TYPE A

People of Type A are numerically stronger and may be considered as basic of Cemetery R 37. Mean values with standard errors, ranges and indices of Type A males are presented in Tables 10-15 and that of females in Tables 28-33, while photographs and dioptograph drawings of two skulls (one male and one female) representing Type A are given in Plates XIII, XIV and Figures 46-48, 49-51.

Before proceeding to a detailed discussion of each of the types indicated, it will be worthwhile to mention their salient morphological characters. Essential cranio-facial features of Type A males and females are noted below:

	MALE		FEMA	LE
Length-breadth index	Dolichocranic	(71.24 ± 0.82)	Dolichocranic	(73.02 ± 1.33)
Length-height index	Orthocranic	(71.34 ± 1.01)	Orthocranic	(70.27 ± 0.87)
Breadth-height index	Acrocranic	(100.19 ± 1.26)	Metriocranic	(97.69 ± 1.97)
Length-auricular height index	Orthocranic	(60.80 ± 0.79)	Orthocranic	(61.64 ± 0.73)
Total facial index	Leptoprosopic	(93.60 ± 4.35)	Mesoprosopic	(89.26)
Superior facial index	Mesen	(51.96 ± 2.26)	Mesen •	(51.61 ± 0.42)
Orbital index*	Mesoconchic	(80.80 ± 1.82)	Mesoconchic	(81.19 ± 1.26)
Nasal index	Chamaerrhine	(51.58 ± 1.92)	Chamaerrhine	(50.96 ± 1.32)
Palatal index	Leptostaphylin	(79.31 ± 3.73)	Brachystaphylin	(88.27 ± 2.11)
Trans, fronto-parietal index	Eurymetopic	(70.60 ± 0.80)	Eurymetopic	(72.17 ± 1.86)

It is difficult to find any significant sex difference when the above essential metric features of Type A skulls are compared. In cranial index, however, the females have a slightly higher average than the males, a feature more often found in delichocephalic races, a distinction first noted by Stockard (Kappers 1934: 3). In Panjabi male and female crania similar observation was made by Pearson (1897: 368).

^{*} Pooled average of right and left orbital index.

The average values of absolute dimensions of both the sexes are: Cranial length 188.00 mm for the males and 179.38 mm for the females; cranial breadth 133.95 mm for the males and 131.42 mm for the females; auricular height 114.22 mm and 111.14 mm for the males and the females respectively; nasal height 52.28 mm for the males and 48.30 mm for the females; nasal breadth 27.19 mm for the males and 24.55 mm for the females; anterior inter-orbital breadth 19.44 for the males and 18.62 for the females; orbital breadth (rt.) 42.38 mm for the males and 41.95 mm for the females; orbital height (rt.) 34.00 for the males and 33.95 mm for the females; minimum frontal diameter 94.60 mm for the males and 93.44 mm for the females; vertical transverse arc 300.56 mm for the males and 300.71 mm for the females.

In all the cranio-facial dimensions the males have higher averages than the females. But the sex difference is quite apparent in Type A skulls if the morphological traits only are taken into consideration. These may be laid down as follows:

	TYPE A			
CHARACTERS	MALE	FEMALE		
Development of Supraorbital ridges	Pronounced	Møderate		
Slope of the forehead	Receding	Slightly receding		
Depression of the nasal root	Depressed	Medium to Shallow		
Protuberence of the occipital region	Markedly protruding (occiput en chignon) to Moderately protruding	Moderately protruding to Rounded		
Mastoid process	Large	Medium		
Nuchal plane	Muscle ridges heavy	Muscle ridges medium		
Subnasal prognathism	Marked	Medium to Marked		

A careful assessment of all the skulls belonging to Type A, has revealed important facts about some of the morphological characters which merit mentioning. These are pronounced supraorbital ridges, low receding forehead, ill-arched sagittal contour, protruding occiput 'shelving' abruptly downwards and forwards, low rectangular orbital cavities inclined laterally and downwards, deep nasion depression, 'wing-shaped' and short nasal bones, broad face, and strong muscular impressions. In this connection specially one male skull No. H 779(e) deserves mentioning; although the sum total of the features including the neurocranium of the said skull exhibit Type A characters, the splanchnocranium reveals some features to be considered as Type A.

Now, special attention is drawn to five skulls (3 males Nos. H 794, Skl. 1, Skl. 10 and 2 females Nos. H 806/A, H 812) attributed as Type A, which exhibit more rugged features having long, slightly against nose and narrow nasal aperture with comparatively narrow facial width in relation to its height. These crania have well-arched vault and well-filled temporo-parietal region.




Fig. 46 Norma Frontalis




Fig. 47 Norma Verticalis





Fig. 48 Norma Lateralis

Figs. 46, 47 and 48. Dieptographic contours of Type A Male [Skl. H 820 (iii)] of Cemetery R 37



Norma Frontalis



Norma Verticalis



Norma Lateralis

Photographs of Type A Male $|\,Skl,\,\,H\,$ 820 $(iii)\,\,|\,\,$ of Cemetery R $\,37$

TYPE A

A little less than 30% (3 males and 7 females) of the collection under study of Cemetery R 37 has been placed under Type $A_{\rm L}$.

In Tables 16-21 the mean values with standard errors, ranges and indices of males of Type A_1 are given and in Tables 34-38 those of females given. Photographs and dioptograph drawings of two skulls (one male and the other female) representing Type A_1 are produced in Plates XV-XVI and in Figures 52-57.

Some essential features of Type A_1 male and female are detailed below:

	MALE		FEM	IALE
Length-breadth index	Dolichocranic	(70.44 ± 0.75)	Dolichocranic	(72.25 ± 1.92)
Length-height index	Orthocranic	(71.63)	Orthocranic	(70.43 ± 1.07)
Breadth-height index	Acrocranic	(100.75)	Acrocranic	(100.77 ± 2.36)
Length-auricular height index	Hypsicranic	(63.09 ± 0.79)	Orthocranic	(62.20 ± 1.03)
Superior facial index		* * *	Mesen	(50.81)
Orbital index	Mesoconchic	(79.83 ± 2.40)	Hypsiconchic	(85.37 ± 1.10)
Nasal index	Meserrhine	(49.75 ± 1.19)	Chamaerrhine	(51.48 ± 1.72)
Palatal index	Mesostaphyline	(83.56 ± 5.94)	Brachystaphylin	(93.72 ± 2.76)
Trans. fronto-parietal index	Eurymetopic	(74.52)	Eurymetopic	(70.65 ± 1.87)

It appears that sexual difference is present only in three indices, namely palatal, orbital and trans. fronto-parietal. Among females, auricular height, nasal height and orbital breadth are slightly lesser in dimension. Auricular height is 111.36 mm for females and 117.33 mm for males; nasal height 48.90 mm for females and 51.00 mm for males; orbital breadth 39.72 mm for females and 41.96 mm for males. All the other essential indices are more or less uniform between the sexes.

The measurements which warrantably prove the homogenity of the type are: Cranial length 186.00 mm for males and 180.08 mm for females; cranial breadth 131.00 mm for males and 129.92 mm for females; basion-bregma height 132.00 mm for males and 128.60 mm for females; nasion-inion length 173.33 mm for males and 167.17 mm for females; longitudinal arc 373.00 mm for males and 374.80 mm for females.

The cranial capacity of males has a higher average than females, which is 1378.67 cc. and 1281,12 cc. respectively.

Some of the morphological traits of males and females of Type A_1 are given below. They indicate that sexual difference is practically absent:

190000 10000 10000 10000 10000	TYPE A	
CHARACTERS	MALE	FEMALE
Development of supraorbital ridges	Slight	Slight to Absent
Slope of the forehead	Vertical	Vertical
Depression of the nasal root	Shallow	Shallow to Absent
Protuberence of the occipital region	Moderately protruding	Moderately protruding
Mastoid process	Medium .	Small
Nuchal plane	Muscular ridges Medium to Slight	Muscular ridges Slight
Subnasal prognathism	Slight to Absent	Absent

It appears that Type A₁ crania are notable for ill-defined muscular relief and general gracility. Both glabella and supraorbital ridges are less developed, forehead narrow vertical, occiput moderately protruding, upper face is of moderate breadth and height, nose narrow-constricted and high-pitched. On the whole, 'total morphological pattern' of Type A₁ skulls resembles Mohenjo-daro 'Mediterranean' type.

DISCUSSION

In summing up the physical characteristics of the population of Cemetery $\mathbb R$ 37, we may state that the oldest inhabitants of Harappa, so far discovered, consisted of mainly two physical types, namely, Type A and Type A_1 , both of which were long-headed.

Type A, which is more numerously represented (11 males and 10 females), were marked by large, strongly musculatured skulls, having low receding forehead, pronounced supraorbital ridges, relatively low rectangular orbits, broad nose sunken at the root, upper face height medium or low and occipital line marked by a prominent occipital protuberence (may be due to the upward and backward development of the brain-case) and rather broad cheek bones.

The average length-breadth index for males is 71.24, for females 73.02: the length-height index for males 71.34, for females 70.27 and breadth-height index for males 100.19, for females 97.69. As a whole, the Type A skulls are mostly delichecranic, chamaecranic or low orthogranic and acrocranic.

Among the Type A skulls, there are at least five adult specimens (3 males Nos. H 794, Skl. 1, Skl. 10 and 2 females Nos. H 806/A, H 812), which deserve special attention for having massive brow ridges, remarkably strong cranial relief endowed with long, slightly acquiline nose and narrow nasal




Fig. 52 Norma Frontais




Fig. 53 Norma Verticalis





Fig. 54 Norma Lateralis


Figs. 52, 53 and 54. Dioptographic contours of Type A, Male [Skl, H 793 (B)] of Cemetery R 37



Norma Frontalis




Norma Verticalis



Norma Lateralis

Photographs of Type A_1 Male [Skl. H 793 (B)] of Cemetery R 37



Norma Frontalis



Norma Verticalis



Photographs of Type A_1 Female [Skl. H $801/A\,\!\!\mid\,$ of Cemetery R 37




Fig. 55 Norma Frontalis




Fig. 56 Norma Verticalis




Fig. 57 Norma Lateralis

Figs. 55, 56 and 57. Dioptographic contours of Type A_i Female [Skl. H 801/A] of Cemetery R 37

aperture with comparatively narrow facial width (Pl. XVIII). Furthermore, the male skulls (Nos. Skl. 1 and Skl. 10) exhibit some tendency toward sagittal keeling of the vault. It is, however, difficult to state, whether these five crania are merely expected variants or they pose a separate entity. But we may well ignore the latter view following Krogman (1937: 216), as the sum total of craniofacial features are not differing significantly, while the outstanding features are fairly well in agreement. Sewell and Guha (1931: 641-642) tried to show the evidence of relationship between 'the Kish skulls with a nose that is long and narrow; in the Al'Ubaid skulls the nose is slightly shorter and distinctly broader, and these changes are more marked in the Adittanallur and Mohenjo-daro skulls, which have chamaerrhine noses, and finally the maximum alteration is found in the Veddas. Similarly, the orbit is very low in the Kish skulls and reaches its maximum height in the Veddas. Another alteration in the character in the face is the change from orthognathism to prognathism.... The bizygomatic breadth, again, shows a graded series from the Kish to the Adittanallur skulls'. And if such variation is possible within normal range, then it is not unjustifiable to lump together the present series with Type A. These five skulls with relatively rugose muscular attachments (particularly in males) and few distinctive features already mentioned can be attributed to Krogman's (1940:17) 'Proto-Nordic' type of Hissar or Dixon's 'Caspian', while rest of the skulls also belonging to Type A conforms with his (Krogman) Pseudo-Australoid type. The mean values of few important features of the male Type A skulls similar to Proto-Nordic and Pseudo-Australoid types are given in the table below. The chief difference between them lies mainly on the facial architecture. They, however,

Characters	Mean values of the male Type A crania similar to Proto-Nordic type	Mean values of the male Type A crania similar to Pseudo-Australoid type
Cranial capacity	1406.12 cc.	1369.03 cc.
Cranial module	155.06 mm	150.54 mm
Nasal index	51.67	54.02*
Stature	1830.63	1746.70

do not differ to any great degree, most of the measurements and indices showing good agreement. We do not find it—justifiable to sort out the five skulls as a separate type. In this connection Hooton's (1946: 616) view regarding the presence of Nordic strain among certain of the earlier skulls of Mohenjodaro may also be referred to. Taking into account the morphological and metrical traits, it is reasonable to conclude that the Type A of the present series stands closer to the norms of Mohenjodaro 'Group A' (Guha and Basu 1938: 630-631) skulls, which has been initially designated by Sewell and Guha (1931: 638-642) as 'Proto-Australoid' and subsequently redesignated by Sir Aurthur Keith (1931: 1002) as 'Caucasic'. Friederichs and Müller (1933: 383) preferred to call them as 'Veddoid'. Buxton and Rice (1931: 69) have referred to similar type of crania as 'Eurafrican'—a term coined by Sergi, Giuffrida-Ruggeri and Fleure. Coon (1939: 87) is of opinion that the large, heavy, and purely delichocephalic type belongs to the large- and longer-headed Mediterranean division, nearest in vault size and form to the earlier Galley Itill and Combe Capelle.

^{*} In averaging nasal index skull No. H 779 (e) had been excluded from reasons see page 51 infra)

Type A₁, represented by fewer skulls (3 males and 7 females) is, on the whole, more smooth and gracile, the entire physical structure being finer and weaker. The musculature is weak, having steep forehead with faintly developed glabello-superciliary region, medium nose with no depression at the root and protruding tuber occipitale. There is no evidence of distinct subnasal prognathism. The average length-breadth index for males is 70.44, for females 72.25; length-auricular height index for males is 63.09, for females 62.20; and superior facial index for females is 50.81 and for males there is no definite index value due to absence of craniometric landmarks but their face appears to be mesen. In other words, Type A₁ is a dolichocranic-orthocranic type having a medium face. This Type A1 of Harappa is similar to 'Group B' of Mohenjo-daro which has been designated by Sewell and Guha as 'Mediterranean' race. Sir Aurthur Keith (1931: 1002) did not, however, agree to the use of this term, while Kappers (1934: 118) is convinced that the name 'Indo-European' or 'Caspian' race used by Dixon (1923: 483) is more appropriate than the name 'Mediterranean' used by Sewell and Guha. In general, Type A skulls differ from Type A more in morphological features than in absolute measurements and cranio-facial proportions.

B. EXTREMITY BONES

Due to fragmentary condition very small number of extremity bones could have been utilized for study. Limb bones described here comprise those which are intact or nearly so. Individual measurements and indices of the long bones are given in detail in the Collective Table L, and average values for the principal measurements and indices are entered in Tables 6-9

In general, extremity bones representing Cemetery R 37 are long and powerful in structure. In this respect Skl. 1 and Skl. 10 deserve special mention for their unusual length and large muscle insertions. On the other hand Skl. H 818 and H 811 are slender with moderate to weak muscular markings.


HUMERUS

Measurements have been made on eight adult humeri (five right and three left) on which most measurements could be taken. Of the eight humeri, maximum length of Skl. 10 is remarkably high (389 mm). Mean values* of some important measurements and indices of humeri are given in Table 6. It shows that mean maximum length is 343 mm and 303 mm for males and females


TABLE 6 Humerus

***		MALES			FEMALES	
Characters	n	Range	Mean	n	Range	Mean
-	5	301.0 - 389.0	343.20	2	293.0 - 314.0	303.50
Maximum length	4	45.0? - 56.5	52.00	2	42.0 - 43.0	42.50
Breadth of proximal epiphysis	4	57.0 - 72.0	65.50	2	52.5 - 53.0	52.75
Breadth of distal epiphysis Breadth of distal epiphysis Circumference of the shaft at the middle	5	57.0 - 77.0	66.80	2	52.0 - 55.0	53.50
Circumference of the solution of diaphysis	5	55.0 - 65.0	58.80	3	50.0 - 52.5	50.83
Robusticity index	5	14.4 - 18.31?	17.24	2	15.92 - 17.06	16.49

^{*} Unless otherwise stated the right and left bones will be put together.



Figs. 1-4 Anterior aspect of Tibla & Femur : Cemetery R 37 (1/2 Natural size)



Frontal view




Vertical view




Left lateral view




Occipital view




Frontal view




Vertical view



Left lateral view



Occipital view



Frontal view



Vertical view



Left lateral view




Occipital view



Frontal view



Vertical view



Left lateral view




Occipital view




Frontal view



Vertical view



Left lateral view



Basal view




Frontal view



 $Vertical\ view$



Left lateral view



Occipital view








Fig. 2 Vertical view





Fig. 3 Occipital view



Fig. 4 Frontal view



Fig. 5 Left lateral view Fig. 6 Occipital view



Figs. 1-3, Skl. H 811; Figs. 4-6, Skl. H 818: R 37



Frontal view



Vertical view



Left lateral view



Occipital view



 $Frontal\ view$




Vertical view




Left lateral view




Occipital view






Vertical view

 $Occipital\ view$



Left lateral view

Skl. 10 : R 37



Ski. H 801/A

Skl. H 806



Fig. 2 Left lateral aspect



Fig. 3 Occlusal aspect



Fig. 4 Left lateral aspect



Fig. 5 Occlusal aspect




Fig. 6 Left lateral aspect

MANDIBLE: R 37

Skl. 10




Fig. 1 Anterior aspect of the Femora of Skl. 10 a, left; b, right




Fig. 3 Anterior view of the left Tibia and 1 bula in articulation (Skf. 10)



Fig. 2 Posterior aspect of the Femora of Skl. 10 a, left: b, right




Fig. 4 Lateral aspect of Fibula: a, Skl. 10; b, Skl. 2; c, H 818

respectively. Circumference at the middle of the shaft also shows distinct difference; it is 66.80 mm for males and 53.50 mm for females. In robusticity index females have slightly lower index value. In none of the humeri perforated olecranon fossa or medial supracondylar spar is found.

RADIUS AND ULNA

 N_0 complete or well-preserved radii and ulna were available. Very few measurements taken on incomplete bones are given in Collective Table L.

FEMUR

Measurements have been made on only eight femora (five right and three left) which are complete or partially complete, showing no pathological changes. The average maximum length of the male is 510.60 mm and that of the female is 437.0 mm. The platymeric index for the Cemetery R 37 femora are 82.24 and 90.91, for males and females respectively. Robusticity index, which expresses the relation of the dimensions at the diaphysis middle and the length in natural position, is 11.73 for males and 11.83 for females. The circumference at the middle of the shaft is 90.71 mm for males and 80.0 mm for females. Pilastric index is somewhat higher in females (104.00) than males (97.81).

The gluteal ridges and linea aspera of the femora of Skl. 1, Skl. 10, H 794 and H 793 are strongly marked. Trace of hypotrochanteric fossa is exhibited on Skl. H 815. It may be pointed out that the measurement of length in natural position of one femur (Skl. 10) exceeds even the world record noted by Martin (1928: 1133).

TABLE 7
Femur

Characters		MALES		FEMALES			
	n	Range	Mean	n	Range	Mean	
e de la							
Maximum length	5	467.0 - 545.0	510.60*	1		437.0	
Length in natural position	5	462.0 - 543.0	507.40	1		431.0	
Sagittal diameter at middle of diaphysis	7	22.0 - 34.5	28.71	1		26.0	
Transversal diameter at middle of diaphysis	7	27.0 - 32.5	29.36	1	Married .	25.0	
Circumference of the shaft at the middle	7	79.0 - 100.0	90.71	1	-	80.0	
Length-thickness index	5	17.10 - 19.96	18.12	1) memory	18.56	
Robusticity index	5	10.68 - 13.04	11.73	1		11.83	
Pilastric index	7	78.57 - 117.54	97.81	1		104.00	
Platymeric index	7	77.42 - 90.91	82.24	1 .	_	90.91	

^{*} In calculating mean values, slightly defective maximum length of Skl. H 811 has been excluded.

In all eleven tibia (four right and seven left) of Cemetery R 37 could be measured. Mean maximum length (spino-malleolar) for males is 409.60 mm and for females 379.00 mm. Index of cross-section at the middle of the diaphysis shows higher value for females (76.43) than males (64.81).

Table 8
Tibia

Characters		MALES			FEMALES	
	n	Range	Mean	n	Range	Mean
Maximum length (spino-malleolar)	5	374.0 - 460.0	409.60	2	372.0 - 386.0	379.00
Physiological length	7	348.0 - 433.0	377.29	2	352.5 - 364.0	358.25
Sagittal diameter at middle	8	27.0 - 36.0	32.56	3	23.5 - 30.0	25.83
Transversal diameter at middle	8	19.5 - 24.0	21.00	3	19.0 - 20.0	19.50
Minimum circumference of diaphysis	8	64.0 - 88.0	75.69	3	62.0 - 71.0	65.50
Index of cross-section in the middle	8	57.35 - 75. 9 3	64.81	3	65.00 - 85.11	76.43
Robusticity index	5	17.16 - 20.21	18.51	1		16.71

Only two male fibulae (both left) are intact. Their mean measurements are as follows:

Table 9

Channel			MALES		
Characters			Range	Mean	
Maximum length		2	368.0 - 435.0	401.50	
Minimum circumference		2	34.0 - 39.0	36.5 0	
Robusticity index		2	8.97 - 9.24	9.10	

STATURE AND PROPORTION

For the purposes of comparison of the deviations of the results given by different methods, the living stature was estimated by applying Pearson's, Manouvrier's and Dupertuis & Hadden's formulae basing on measurements of dry long bones. Karl Pearson was a pioneer in applying the statistical method of regression analysis to the prediction of stature from the length of long bones. It is difficult to ascertain as to whose formulae are most suitable for our purpose. Then again, the formulae are based upon researches on materials of countries other than India and, therefore, the estimations made by applying those formulae may not be very accurate with respect to Harappan skeletons. Notwithstanding the criticisms made on the fundamental researches of Manouvrier and Pearson, we have utilized their tables and formulae respectively in addition to a set of 'general formulae' of Dupertuis and Hadden, which is superior to any other designed for a similar purpose and is expected to yield

good approximations to statures of many non-European people (Boyd and Trevor 1953: 147). Most of the formulae yielding best results are based either on the combination of corresponding bones of upper and lower limbs or two bones of either extremity, which unfortunately were not available in the skeletons of the individuals from Harappa. We had, therefore, mainly the recourse of predicting stature from the length of a single dry long bone or bones of an individual.

It is evident from the Collective Table O, that on the whole Pearson's values are lowest and Dupertuis and Hadden's are highest, those of Manouvrier's standing intermediate. The only exception are with Skl. 10 (male) and H 820 (female), where results obtained using Manouvrier's method give higher figures than Dupertuis and Hadden's.

Of the 108 individuals of Cemetery R 37, stature of only thirteen (eight males and five females) could be estimated from dry long bones which are intact, and display no pathological changes. It is seen that Skl. 1 (male) was the tallest individual not only of Cemetery R 37, but of the whole Harappan population.

Among the males representing Type A the tendency towards tallness is quite apparent. Skl. 1 and Skl. 10 were very tall in stature. Skl. H 794 and H 796 (B) were tall and Skl. H 793 was either above medium or tall. Skl. H 798, whose skull was not available, and hence could not be classified in either of the types, was medium statured. Among the females Skl. H 820 and H 821 were very tall in stature. They too were not diagnosed with respect to the physical features of the two groups. Similar is the case with Skl. H 805 (a), who was either above medium or tall. Stature of two females, viz., Skl. H 780 and H 806/A belonging to Type A could be estimated, the former one being either above medium or tall and the latter medium.

Of the Type A_{I} skeletons, Skl. H 811 was tall and H 818 was either below medium or medium in stature.

It is observed that there is a distinct difference in mean stature between Type A and Type A_1 males although sample sizes are too small to comment.

100 miles (100 miles (STATURE ACCORDING TO	
	MANOUVRIER	PEARSON	DUPERTUIS AND HADDEN
Type A (M)	1780.25 (5)	1743.84 (5)	1797.06 (5)
Type $A_1(M)$	1684.46 (2)	1665.65 (2)	1711.65 (2)
Type A (F)	1574.12 (2)	1578. 4 (1)	1626.6 (1)

The estimated average stature for the males is 1757.1 mm for the males and 1650.6 mm for the females.

For the determination of limb proportions only two skeletons (Nos. Skl. 1 and Skl. 10) could be utilized from Cemetery R 37. The tibiae of Skl.10 are relatively long with respect to its femora, the tibio-femoral index being dolichocnemic (right 84.01? and left 84.40) according to Turner's (1899) classification. Humero-femoral index could be worked out of Skl. 1 and Skl. 10. In both the cases humeri are proportionately shorter when compared with femora. In case of Skl. 1 humerus is 69.69% of the femur and it is 71.51% in case of Skl. 10.

STATISTICAL CONSTANT : CEMETERY R 37

Table 10 ADULT MALE TYPE A Mean Measurements* and Indices of the Neurocranium

Characters	n	Mean	±	S. E.	Min.	Max.
Max. cranial length	10	188.00	±	1.37	178.00	193.00
Max. cranial breadth	11	133.95	\pm	0.97	130.00	140.00
Nasion-inion length	9	173.89	\pm	2.73	157.00	182.50
Basion-bregma height	10	134.05	\pm	1.59	128.00	145.00
Min. frontal breadth	10	94.60	\pm	0.89	90.00	97.00
Vertical porion height	9	114.22	\pm	0.96	110.50	118.50
Median sagittal arc	8	376.12	\pm	4.37	353.00	395.00
Vertical transversal arc	9	300.56	\pm	1.66	294.00	309.00
Herizontal circumference	9	521.67	\pm	3.35	500.00	532.00
Cranial module	10	151.97	\pm	0.80	148.50	156.17
Calculated cranial capacity	10	1390.24	<u>+</u>	15.00	1335.42	1481.60
Length-breadth index	10	71.24	\pm	0.82	68.06	75.84
Length-height index	10	71.34	<u>+</u>	1.01	66.84	77.13
Breadth-height index	10	100.19	\pm	1.26	93.57?	107.41
Length-auricular height index	9	60.80	±	0.79	57.51?	64.40
Breadth-auricular height index	9	85.15	土	0.61	82.14	88.43
Tr. fronto-parietal index	10	70.60	\pm	0.80	64.98	74.05

TABLE 11 ADULT MALE TYPE A Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	±	S. E.	Mirr.	Max.
Prosthion-basion line Nasion-prosthion line Nasion-gnathion line Bizygomatic breadth Nasal height Nasal breadth Ant. inter-orbital breadth Orbital breadth (right) Orbital breadth (left)	9 9 5 5 9 8 8 9	101.28 70.78 125.70 133.10 52.28 27.19 19.44 42.38 42.50 34.00	* * * * * * * * * * * * * * * * * * * *	1.08 1.52 3.29 2.30 1.09 0.96 0.58 0.63 0.70 1.12	 96.0? 61.0 118.5 125.0 48.0 22.0 18.0 39.0 40.0	106.0 77.0 135.0 138.0 58.5 31.0 23.0 44.5 45.5 37.0

^{*} Measurements fellowed by query i c sless oper to the true values, except stated specifically.

TABLE 11 - Continued

Characters	n	Mean	土	S.E.	Min.	Max
Orbital height (left)	9	34.44	±	0.86	30.5	38.0
Maxillo-alveolar length	8	59.31	±	0.74	56.5	63.0
Maxillo-alveolar breadth	7	65.64	土	1.17	61.5	69.0
Palatal length	7	49.57	±	1.43	44.0	55.0
Palatal breadth	7	40.00	±	1.29	35.0	45.0
Total facial index	3	93.60	土	4.35	88.76?	102.27
Superior facial index	5	51.96	\pm	2.26	44.20	57.2 0
Orbital index (right)	8	80.33	\pm	2.82	66.67	92.31
Orbital index (left)	9	81.23	\pm	2.48	71.43	95.00
Nasal index	8	51.58	\pm	1.92	41.51	57.94
Maxillo-alveolar index	7	111.67	\pm	1.67	106.03	118.58
Palatal index	6	79.31	\pm	3.73	74.00	97.73

TABLE 12

ADULT MALE TYPE A

Mean Indices of the whole Skull

and the same of th						
Characters	n	Mean	土	S. E.	Min.	Max.
**** · · · · · · · · · · · · · · · · ·				CORC CONTRACTOR		
Tr. cranio-facial index	5	98.70	\pm	1.60	9 5.31	102.99
Vert. cranio-facial index	9	52.97	\pm	1.20	45.19	58.11
Long. cranio-facial index	9	53.90	\pm	0.68	50.26?	56.74
Jugo-frontal index	5	71.31	\pm	1.63	68.18	77.60

TABLE 13

ADULT MALE TYPE A

Mean Measurements & Indices of the Mandible

Characters	, n	Mean	\pm	S. E.	Min.	Max,	
Bigonial breadth	7	90.79	土	2.92	81.0	101.0	
Bicondylar breadth	6	121.75	\pm	5.22	99.0	135.0	
		66.33	土	2.00	60.0	74.0	
Ht. of mandibular ramus	6						
Max. breadth of mandibular ramus	6	45.42	±	1.73	41.0	52.0	
Min. breadth of mandibular ramus	7	33.07	\pm	0.88	30.0	36.0	

TABLE 13 - Continued

Characters	n	Mean	土	S.E.	Min.	Max.
Ht. at mandibular symphysis	7	34.93	±		29.0	40.0
Mandibular length	7	81.00	±	2.09	74.0	90.0
Mandibular angle	6	122.17°	±	2.18	116.5°	128°
Mandibular index	6	68.47	±	4.83	60.38	90.91
Breadth index of mandible	6	75.64	±	2.19	68.94	83.84
Jugo-mandibular index	4	68.78	土	2.98	60.67?	73.72

Table 14 ADULT MALE TYPE A Mean Measurements & Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean	\pm	S. E.	Min.	Max.		
Mesiodistal crown diam. of M_1	6	10.38	\pm	0.22	9.50	11.00		
Labiolingual crown diam, of M_1	6	11.38	\pm	0.17	10.75	12.00		
Mesiodistal crown diam, of M_2	7	9.75	\pm	0.30	8.50	10.50		
Labiolingual crown diam. of M_2	7	11.36	土	0.26	10.50	12.50		
Mesiodistal crow diam. of M ₃	1.15	9.50	\pm	0.42	8.50	10.50		
Labiolingual crown diam. of M ₃	5	10.80	\pm	0.41	9.50	12.00		
Crown index of M_1	6	109.93	\pm	2.40	104.76	121.39		
Crown index of M ₂	7	117.62	\pm	4.08	104.88	136.50		
Crown index of M_3	5	114.07	\pm	2.92	104.76	122.60		

TABLE 15 ADULT MALE TYPE A Mean Measurements & Indices of Permanent Mandibular Molar Teeth

Characters	n	Mean	\pm	S. E.	Min.	Max.
2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2						
Mesiodistal crown diam. of M ₁	5	10.15	\pm	0.31	9.00?	10.75
Labiolingual crown diam, of M1	5	10.90	\pm	0.19	10.50	11.50
Mesiodistal crown diam, of M ₂	4	10.25	+	0.44	9.00	11.00
Labiolingual crown diam, of M2	4	10.31	\pm	0.28	9.50	10.75
Mesiodistal crown diam, of M.	5	9.80	<u>+</u>	0.66	8.00	11.75
Labielingual crown diams of M3	5	9.75	+	0.32	9.00	10.50
Clown index of M.	5	107.98	\pm	5.02	100.00	127.78?
or and index of Ma	14	,101,28	+	*5.70	90.48	116.67
Crown index of M ₃	18	100.92	±	6.00	85.71	118.75

TABLE 16

ADULT MALE TYPE A

Mean Measurements & Indices of the Neurocranium

				man the		
Characters	n	Mean	\pm	S. E.	Min.	Max.
Max. cranial length	3	186.00	\pm	1.16	184.0	188.0
Max. cranial breadth	3	131.00	土	0.58	130.0	132.0?
Nasion-inion length	3	173.33	\pm	1.77	170.0	176.0
Basion-bregma height	2	132.50			129.0	136.0
Min. frontal breadth	2	98.00			97.0	. 99.0
Vertical porion height	3	117.33	±	0.88	116.0	119.0
Median sagittal arc	3	373.00	\pm	8.75	356.0	3 8 5. 0
Vertical transversal arc	1	301.00			-	
Horizontal circumference	2	512.50			508.0	517.0
Cranial module	2	149.67			148.67	150.67?
Calculated cranial capacity	3	1378.67	土	6.50	1367.30	1389.77?
Length-breadth index	3	70.44	±	0.75	69.15	71.74?
Length-height index	2	71.63			69.35	73.91
Ereadth-height index	2	100.75			98.47	103.03?
Length-auricular height index	3	63.09	±	0.79	62.23	64.67
Breadth-auricular height index	3	89.57	\pm	0.51	88.55	90.15?
Tr. fronto-parietal index	2	74.52			73.48?	75.57

Table 17 ${\tt ADULT\ MALE\ TYPE\ A_1}$ Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	土	S. E.	Min.	Max.
Prosthion-basion line	2	105.50			98.0	113.0
Nasion-prosthion line	3	70.17	土	1.92	68.0	74.0
Nasion-gnathion line	-				· ·	
Bizygomatic breadth	1	122.00			-	
Nasal height	3	51.00	土	1.73	48.0	54.0
Nasal breadth	3	25.33	\pm	0.34	25.0	26.0?
Ant. inter-orbital breadth	3	19.33	土	0.88	18.0	21.0
Orbital breadth (right)	3	42.17	土	0.88	40.5	43.5
Orbital breadth (left)	2	41.75			41.5	42.0
	3	34.67	土	0.67	34.0	36.0
Orbital height (right)		32.33	土	0.60	31.5	33.5
Orbital height (left)	3	63.00				
Maxillo-alveolar length	1	63.00				enement.
Maxillo-alveolar breadth	1	63.00				

TABLE 17 - Continued

		STATE OF THE PARTY					
Characters	n	Mean	土	S.E.	Min.	Max.	
9-30							
Palatal length	3	45.67	土	1.34	43.0	47.0	
Palatal breadth	3	38.00	\pm	1.53	36.0	41.0	
Total facial index	_	_			_	_	
Superior facial index	1	56.15			_	_	
Orbital index (right)	3	82 .35	土	3.32	78.16	88.89	
Orbital index (left)	2	76.06			75.00	77.11	
Nasal index	3	49.75	\pm	1.19	48.15?	52.08	
Maxillo-alveolar index		_			_	_	
Palatal index	3	83.56	\pm	5.94	76.60	95.35	

TABLE 18 ADULT MALE TYPE A

Mean Indices of the whole Skull

Characters	n	Mean \pm S.E.	Min.	Max.
Tr. cranio-facial index	1	93.85		
Vert. cranio-facial index	2	53.56	52.71	54.41
Long, cranio-facial index	2	57.05	52.69	61.41
Jugo-frontal index	_	-	_	

TABLE 19 ADULT MALE TYPE A

Mean Measurements & Indices of the Mandible

			•	
Characters	n	Mean ± S.E.	Min.	Max.
Bigonial breadth	1	97.0	and the state of t	
Bicondylar breadth	1	114.0?		
Ht. of mandibular ramus	1	65.0		
Max. breadth of mandibular ramus	1	39.0?		
Min. breadth of mandibular ramus	2	34.0	31.0	37. 0
Ht. at mandibular symphysis	1	26.0		0.10
Mandibular length	2	76.0	74.0	78.0
Mandibular angle	2	119°	118°	1200
Mandibular index ,	1	64.91	_	120°
Breadth index of mandible	1	85.09		
Jugo-mandibular index	1	79.51		

Characters	n	Mean	土	S. E.	Min.	Max.
Mesiodistal crown diam. of M ₁	3	10.00	±	0.14	9.75	10.25
Labiolingual crown diam. of M1	3	10.67	\pm	0.44	10.00	11.50
Mesiodistal crown diam. of M_2	3	10.08	\pm	0.21	9.75	10.50
Labiolingual crown diam. of M2	3	10.67	\pm	0.44	10.00	11.50
Mesiodistal crow diam. of M ₃	1	7.50				_
Labiolingual crown diam. of M ₃	1	10.00				. —
Crown index of M ₁	3	106.80	\pm	6.03	97.37	118.02
Crown index of M2	3	105.63	±	2.09	102.38	109.52
Crown index of M_3	1	133.33			-	

TABLE 21

ADULT MALE TYPE A,

Mean Measurements & Indices of Permanent Mandibular Molar Teeth

Characters	n	n Mean ± S.E.		Max.	
Mesiodistal crown diam. of M_1	2	10.25	10.00	10.50	
Labiolingual crown diam. of M_1	2	9.62	9.00	10.25	
Mesiodistal crown diam. of M ₂	2	9.75	9.50	10.00	
Labiolingual crown diam. of M2	2	8.25	7.50	9.00	
Mesiodistal crown diam. of M ₃	2	9.62	8.75	10.50	
Labiolingual crown diam. of M ₃	2	8.25	7.50	9.00	
Crown index of M ₁	2	94.20	85.91	102.50	
Crown index of M ₂	2	84.58	79.16	90.00	
Crown index of M ₃	2	85.74	85.71	85.78	

TABLE 22

ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean ±	S. E.	S. D.	c.v.	Min.	Max.
Max. cranial length	13	187.54 ±	1.09	3. 9 3	2.10	178.0	193.0
Max. cranial breadth	14	$133.32\ \pm$	0.83	3.11	2.33	130.0	140.0
Nasion-inion length	12	$173.75 \pm$	2.05	7.11	4.09	157.0	182.5
Basion-bregma height	12	$133.79 \pm$	1.39	4.81	3.60	128.0	145.0

TABLE 22: Continue d

		• 1							
Characters	n	Mean <u>-</u>	<u>+</u>	S. E.	S. D.	c. v.	Min.	Max.	
Min. frontal breadth	12	95.17 ±	±	0.84	2.89	3.04	90.0	99.0	
Vertical porion height	12	115.00	±	0.84	2.91	2.53	110.5	119.0	
Median sagittal arc	11	375.27 ±	<u>+</u>	3.75	12.44	3.31	353.0	395.0	
Vertical transversal arc	10	300.60 ±	±	1.48	4.69	1.56	294.0	309.0	
Horizontal circumference	11	520.00	±	2.99	9.93	1.91	500.0	532.0	
Cranial module	12	151.58	±	0.72	2.49	1.64	148.50	156.17	
Calculated cranial capacity	13	1387.57 ±	± :	11.53	41.61	3.00	1335.42	1481.60	
Length-breadth index	13	71.05	±	0.65	2.34	3.29	68.06	75.84	
Length-height index	12	71.39	±	88.0	3.04	4.26	66.84	77.13	
Breadth-height index	12	100.28	±	1.08	3.74	3.73	93.57	107.41	
Length-auricular height index	12	61.37	±	0.68	2.36	3.85	57.51?	64.67	
Breadth-auricular height index	12	86.26	±	0.74	2.57	2.98	82.14	90.15	
Tr. fronto-parietal index	12	71.26	±	0.80	2.78	3.90	64.98	75.57	

TABLE 23 ADULT MALE COMBINED Statistical Constants of the Measurements and Indices of the Splanchnocranium

Characters	n	Mean ±	S. E.	S. D.	c.v.	Min.	Max.
Prosthlon-basion line	11	102.05 ±	1.43	4.76	4.66	96.0?	113.0
Nasion-prosthion line	12	$70.62 \pm$	1.20	4.15	5.88	61.0	77.0
Nasion-gnathion line	5	$125.70~\pm$	3.29	7.38	5.87	118.5	135.0
Bizygomatic breadth	6	$131.25 \pm$	2.64	6.46	4.92	122.0	138.0
Nasal height	12	51.96 ±	0.90	3.11	5.99	48.0	58.5
Nasal breadth	11	26.68 ±	0.74	2.45	9.18	22.0	31.0
Ant, inter-orbital breadth	11	19.41 ±	0.46	1.53	7.88	18.0	23.0
Orbital breadth (right)	11	42.32 ±	0.50	1.65	3.90	39.0	44.5
Orbital breadth (left)	11	42.36 ±	0.57	1.90	4.49	40.0	45.5
Orbital height (right)	11	34.18 ±	0.82	2.73	7.99	28.0	37.0
Orbital height (left)	12	33. 9 2 ±	0.70	2.43	7.16	30.5	38.0
Maxillo-alveolar length	9	$59.72 \pm$	0.77	2.31	3.87	56.5	63.0
Maxillo-alveolar breadth	8	6 5.31 ±	1.06	3.01	4.61	61.5	69.0
Palatal length	10	$48.4 \pm$	1.20	3.78	7.81	43.0	55. 0
Palatal breadth	10	39.4 ±	1.01	3.20	8.12	35.0	45.0
Total facial index	3	93.60 ±	4.35	7.52	8.03	88.76?	102.27
Superior facial index	6	$52.66 \pm$	1.98	4.84	9.19	44.20	57.20
Orbital index (right)	11	80.88 ±	2.18	7.23	8.94	66.67	92.31
Orbital index (left)	11	80.29 ±	2.11	7.00	8.72	71.43	95.00
Nasal Index	11	51.08 ±	1.42	4.71	9.22	41.51	57.94
Maxillo-alveolar index	7	111.67 ±	1.67	4.43	3.97	106.03	118.58
Palata] index	9	80.72 ±	3.04	9.11	11.29	74.00	97.73

 $\begin{array}{c} {\bf TABLE~24} \\ \\ {\bf ADULT~MALE~COMBINED} \end{array}$ Statistical Constants of the Indices of the Whole Skull

Characters	n	Mean <u>+</u>	S. E.	S. D.	c. v.	Min.	Max.
Tr. cranio-facial index	6	97.89 ±		3.77	3.85	93.85	102.99
Vert. cranio-facial index	11	$53.08 \pm$	0.98	3.24	6.10	45.19	58.11
Long. cranio-facial index	11	$54.47 \pm$	0.89	2.96	5.43	50.26?	61.41
Jugo-frontal index	5	$71.31 \pm$	1.63	3.65	5.12	68.18	77.60

TABLE 25

ADULT MALE COMBINED

Statistical Constants of the Mandibular Measurements and Indices

Characters	n	Mean <u>+</u>	S. E.	S. D.	c. v.	Min.	Max.
Bigonial breadth	11	89.14 ±	3.52	11.70	13.13	60.0	101.0
Bicondylar breadth	7	$103.93 \pm$	3.14	8.31	8.00	99.0	135.0
Ht. of mandibular ramus	8	$68.38 \pm$	2.67	7.56	11.06	60.0	84.0
Max. breadth of mandibular ramus	8	$45.31 \pm$	1.70	4.82	10.64	39.0	52. 0
Min. breadth of mandibular ramus	13	$34.27 \pm$	0.89	3.21	9.37	30.0	39.0
Ht. at mandibular symphysis	11	$33.50 \pm$	1.40	4.66	13.91	26.0	40.0
Mandibular length	12	$81.92 \pm$	1.78	6.16	7.52	74.0	91.0
Mandibular angle	9	$120.22\ \pm$	1.89	5.66	4.71	111.0	128.0
Mandibular index	7	$67.96 \pm$	4.11	10.88	16.01	60.38	90.91
Breadth index of mandible	7	$76.99 \pm$	2.28	6.05	7.86	68.94	8 5. 09
Jugo-mandibular index	6	$69.74 ~\pm$	2.83	6.93	9.94	60.67?	79.51

TABLE 26

ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean +	S. E.	S. D.	C. V.	Min.	Max.
Characters		mean <u>.</u>	0	2.0			
Mesiodistal crown diam. of M_1	9	10.25 ±	0.16	0.49	4.78	9.50	11.00
Labiolingual crown diam, of M1	9	$11.14 \pm$	0.21	0.62	5.57	10.00	12.00
Mesiodistal crown diam. of M2	10	9.85 ±	0.22	0.69	7.01	8.50	10.50
Labiolingual crown diam. of M_2	10	$11.15 \pm$	0.23	0.73	6.55	10.00	12.50
Mesiodistal crown diam. of M ₃	6	9.17 ±	0.48	1.17	12.76	7.50	10.50
Labiolingual crown diam. of M3	6	10.67 ±	0.36	0.88	8.25	9.50	12.00
Crown index of M ₁	9	108.88 ±	2.39	7.17	6 .5 9	97.37	121.39
Crown index of M2	10	114.02 ±	3.39	10.70	9.38	102.38	136.50
Crown index of M ₃	6	117.28 ±	4.00	9.81	8.36	104.76	133.33

Table 27 ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of Permanent Mandibular Molar Teeth

Characters	n	Mean <u>+</u>	S. E.	S. D.	c. v.	Min.	Max.
Mesiodistal crown diam. of M_1	8	10.28 ±	0.22	0.62	6.03	9.00	11.00
Labiolingual crown diam. of M ₁	8	$10.50 \pm$	0.26	0.74	7.05	9.00	11.50
Mesiodistal crown diam. of M ₂	8	$10.12 \pm$	0.23	0.64	6.32	9.00	11.00
Labiolingual crown diam. of M ₂	. 8	9.59 ±	0.39	1.09	11.37	9.00	10.75
Mesiodistal crown diam. of M ₃	9	9.78 ±	0.38	1.14	11.66	8.00	11.75
Labiolingual crown diam. of M ₃	9	9.33 ±	0.30	0.89	9.54	7.50	10.50
Crown index of M _I	8	$102.69 \pm$	4.26	12.06	11.74	85.91	127.78?
Clown index of M ₂	8	95.00 ±	4.08	11.55	12.16	79.16	116.67
Crown index of M ₃	9	95.41 ±	3.90	11.71	12.27	85.71	118.75

TABLE 28 ADULT FEMALE TYPE A Mean Measurements and Indices of the Neurocranium

Characters	n	Mean	\pm	S. E.	Min.	Max.
Max. cranial length	8	179.38	\pm	2.90	167.0	195.0
Max. cranial breadth	6	131.42	土	2.21	125.0?	139.0
Nasion-inion length	7	170.36	\pm	2.68	157.0	181.0
Basion-bregma height	5	126.60	土	1.21	124.0	131.0
Min. frontal breadth	8	94.75	\pm	1.06	91.0	100.0
Vertical porion height	7	111.14	\pm	1.66	106.0?	120.0
Median sagittal arc	6	35 9.50	\pm	3.69	346.0	372.0
Vertical transversal arc	7	300.71	土	5.08	289.0	325.0
Horizontal circumference	7	502.00	\pm	5.02	483.0	528.0
Cranial module	4	145.46	\pm	1.10	143.50	148.33
Calculated cranial capacity	6	1289.49	\pm	56.18	1174.23	1542.89
Length-breadth index	6	73.02	\pm	1.33	68.68?	77.84
Length-height index	5	70.27	\pm	0.87	68.68	73.60
Breadth-height index	4	97.69	\pm	1.97	92.70	101.55
Length-auricular height index	* 7	61.64	<u>+</u> :	0.73	59.55?	65.27
Breadth-auricular height index	6	84.68	\pm	0.93	81.75	87.20?
Tr. fronto-parietal index	6	72.17	\pm	1.86	66.91	80.00

TABLE 29

ADULT FEMALE TYPE A

Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	土	S. E.	Min.	Max.
Prosthion-basion line	4	93.62	±	3.32	85.0	101.0
Nasion-prosthion line	10	66.45	土	0.97	63.0	72.0
Nasion-gnathion line	3	111.67	土	4.18	107.0	120.0
Bizygomatic breadth	3	125.33	\pm	2.19	121.0	128.0
Nasal height	10	48.30	土	0.73	45.0	53.0
Nasal breadth	10	24.55	土	0.46	23.0	27.0
Ant. inter-orbital breadth	8	18.62	±	0.83	15.0?	22.5
Orbital breadth (right)	9	42.06	±	1.06	38.0	46.0
Orbital breadth (left)	7	41.36	土	0.83	37.0	44.0
Orbital height (right)	9	34.06	土	0.92	31.5	39.0
Orbital height (left)	8	33.75	\pm	0.60	31.5	36.5
Maxillo-alveolar length	8	56.06	\pm	0.70	53.0	59.0
Maxillo-alveolar breadth	8	61.56	\pm	0.57	59.0	63.5
Palatal length	8	45.50	土	1.13	41.0	48.0
Palatal breadth	8	`40.00	\pm	0.03	39.0	41.0
Total facial index	1	89.26			_	_
Superior facial index	3	51.61	\pm	0.42	50.78	52.07
Orbital index (right)	9	81.08	\pm	1.61	76.09	89.47
Orbital index (left)	7	81.34	\pm	2.14	76.83	93.24
Nasal index	10	50.96	\pm	1.32	43.40	56.25
Maxillo-alveolar index	8	109.91	土	1.51	105.36	117.92
Palatal index	8	88.27	土	2.11	8 3.33	100.00

TABLE 30

ADULT FEMALE TYPE A

Mean Indices of the Whole Skull

				c E	Min.	
Characters	n			S. E.		Max.
Tr. cranio-facial index	3	97.02			94.16	98.46
Vert. cranjo-facial index	5	53.11	\pm	1.46	50.38	57.60
Long. cranio-facial index	3	50.48	土	2.03	46.70	53.65
Jugo-frontal index	3	74.24	土	1.58	71.09	76 .03

TABLE 31 ADULT FEMALE TYPE A Mean Measurements and Indices of the Mandible

						Date to severe
Characters	n	Mean	土	S. E.	Min.	Max.
Bigonial breadth	3	78.67	±	4.92	70.0	87.0
Bicondylar breadth	2	104.00			104.0	104.0?
Ht. of mandibular ramus	3	58.50	土	0.76	57.5	60.0
Max. breadth of mandibular ramus	3	39.17	\pm	1.17	38.0	41.5
Min. breadth of mandibular ramus	3	31.33	土	1.09	30.0	33.5
Ht. at mandibular symphysis	4	31.25	\pm	2.50	26.0	35.0
Mandibular length	4 .	75.75	\pm	2.28	73.0	81.0
Mandibular angle	3	117.83	土	1.17	116°	120°
Mandibular index	2	76.44			75.00	77.88?
Breadth index of mandible	2	71.64			67.31	75.96?
Jugo-mandibular index	1	57.85			_	_

Table 32 ADULT FEMALE TYPE A Mean Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean	土	S. E.	Min.	Max.	
Mesiodistal crown diam. of M ₁	7	10.00			10.00	10.00	
Labiolingual crown diam. of M ₁	6	10.42	土	0.46	9.00	12.00	
Mesiodistal crown diam. of M_2	7	9.46	土	0.29	8.00	10.50	
Labiolingual crown diam. of M ₂	7	9.82	土	0.42	8.00	11.25	
Mesiodistal crown diam. of $ m M_3$	6	8.62	土	0.35	7.50	10.00	
Labiolingual crown diam. of $ m M_3$	6	8.79	土	0.45	7.25	10.00	
Crown index of M ₁	6	104.17	土	4.59	90.00	120.00	
Crown index of M2	7	104.09	土	2.13	94.74	113.16	
Crown index of M_3	6	101.98	\pm	2.56	94.12	111.76	

TABLE 33 ADULT FEMALE TYPE A Mean Measurements and Indices of Permanent Mandibular Molar Teeth

Characters	n	Mean	\pm	S. E.	Min.	Max.
Mesiodistal crown diam. of M	2	10.75			10.50	11.00
Labielingual crewn diam, of M1	2	10.12			10.00	10.25
Mesiodistal crown diam. of M ₂	3 -	9.58	土	0.12	9.00	10.25
Labiolingual crown diam. of M2	3	9.00	\pm	0.87	7.50	10.50

TABLE 33 - Continued

Characters	n	Mean	\pm	S.E.	Min.	Max.		
Mesiodistal crown diam. of M_3	3	8.97	\pm	0.30	8.50	9.50		
Labiolingual crown diam. of M_3	3	8.33	\pm	0.88	7.00	10.00		
Crown index of M ₁	2	94.44			90.91	97.96		
·Crown index of M ₂	3	94.01	\pm	8 .53	8 3.33	110.84		
·Crown index of M ₃	3	93.24	\pm	7.38	80.06	105.56		

TABLE 34 ${\hbox{\tt ADULT FEMALE TYPE A}_1}$ Mean Measurements and Indices of the Neurocranium

Characters	n	Mean	土	S.E.	Min.	Max.
Max. cranial length	6	180.08	±	2.53	171.0	186.5
Max. cranial breadth	6	129.92	\pm	2.19	122.0?	138.0
Nasion-inion length	6	167.17	\pm	1.66	162.0	174.0
Basion-bregma height	5	128.60	\pm	1.86	125.0	135.0
Min. frontal breadth	6	91.58	\pm	0.76	89.0	94.0
Vertical porion height	7	111.36	\pm	2.03	104.0	119.0
Median sagittal arc	5	374.80	\pm	5.57	358.0	387.0
Vertical transversal arc	5	308.40	\pm	3.12	297.0	315.0
Horizontal circumference	5	503.00	±	3.97	491.0	514.0
Cranial module	4	147.33	土	1.28	145.00	150.17
Calculated cranial capacity	6	1281.12		39.97	1145.86	1388.02
Length-breadth index	6	72.25	土	1.92	66.30?	80.70
Length-height index	4	70.43	土	1.07	68.6 3	73.37
Breadth-height index	4	100.77	土	2.36	96.90	106.56?
Length-auricular height index	6	62.20	\pm	1.03	58.70	64.91
Length-auriculai height index	6	86.26	土	1.67	80.43	90.49
Breadth-auricular height index [Tr. fronto-parietal index]	5	70.65	土	1.87	64.49	75.00?

TABLE 35

ADULT FEMALE TYPE A₁

Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	±	S.E.	Min,	Max.
Prosthion-basion line Nasion-prosthion line Bizygomatic breadth Nasal height	2 3 4 5	96.00 62.33 124.00 48.90	± ±	1.20 1.29 1.42	94.0 60.0 121.0? 42.0	98.0 64.0 127.0 53.0

TABLE 35 - Continued

Characters	n	Mean	土	S.E.	Min.	Max.
Nasal breadth	5	25.10	±	0.60	24.0	27.0
Ant. inter-orbital breadth	6	18.33	\pm	1.15	14.0	21.0
Orbital breadth (right)	6	39.25	\pm	0.89	36.0?	42.0
Orbital breadth (left)	6	40.50	土	0.84	39.0	44.0
Orbital height (right)	6	33.50	\pm	0.80	31.0	37.0
Orbital height (left)	6	34.25	±	1.12	31.0	38.0
Maxillo-alveolar length	3	51.00	\pm	2.09	48.0	55.0
Maxillo-alveolar_breadth	3	62.50	\pm	2.29	58.0	65.5
Palatal length	3	42.00	\pm	0.58	41.0	43.0
Palatal breadth	3	39.33	\pm	0.67	38.0	40.0
Superior facial index	2	50.81			49.59?	52.03
Orbital index (right)	5	86.42	\pm	1.84	80.49	91.67?
Orbital index (left)	6	84.49	\pm	1.36	79.49	88.10
Nasal index	5	51.48*	±	1.72	45.28	55.32
Maxillo-alveolar index	3	122.64	\pm	2.73	119.09	128.00
Palatal index	3	93.72	±	2.76	88.37	9 7.56

TABLE 36 ADULT FEMALE TYPE A Mean Indices of the Whole Skull

		- 00					
Characters	n	Mean	±	S.E.	Min.		Max.
Tr. cranio-facial index	3	94.71	\pm	2.84	89.13		98.43
Vert. cranio-facial index	2	48.23			48.00		48.46
Long. cranio-facial index	1	51.09				rie.	
Jugo-frontal index	4	73.42	\pm	1.28	70.87		76.86?

TABLE 37 ADULT FEMALE TYPE A Mean Measurements and Indices of the Mandible

Characters		144	n	Mean
	- 1	10		
Bigonial breadth			1	89.0
Bicondylar breadth		Á	1.	117.0
Ht. of mandibular ramus	*		1	62.0
Max. breadth of mandibular	ramus		1	44.0

^{*} Due to depletion of calcium in the alveolar process of maxilla, nash index of H 820 (i) gives an inflated result which has been omitted for consideration of statistical constants.

TABLE 37 - Continue d

Characters ,	n	Mean
Min. breadth of mandibular ramus	1	36.0
Ht. at mandibular symphysis	1	31.0
Mandibular length	1	80.0
Mandibular angle	1	120.5°
Mandibular index	1	68.38
Breadth index of mandible	1	76.07
Jugo-mandibular index	1	70.08

Table 38 ${\bf ADULT\ \ FEMALE\ \ \ TYPE\ A}_1$ Mean Measurements and Indices of Permanent Maxillary Molar Teeth

			*			
n	Mean	Min.	Max			
1	10.50	_				
1	11.00	_				
2	9.88	9.00	10.75			
2	9.62	8.00	11.25			
1	7.00					
1	6.50					
1	104.76		**********			
2	97.10	88.89	105.32			
1	92.86					
	1 1 2 2 2 1 1 1	1 10.50 1 11.00 2 9.88 2 9.62 1 7.00 1 6.50 1 104.76 2 97.10	1 10.50 — 1 11.00 — 2 9.88 9.00 2 9.62 8.00 1 7.00 — 1 6.50 — 1 104.76 — 2 97.10 88.89			

Table 39

Adult Female Combined

Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean	土	S.E.	S.D.	C.V.	Min.	Max.
** C. 11 De 100 DE 1 COMMON DE 100 DE								
Max. cranial length	14	179.68	土	1.91	7.16	3.98	167.0	195.0
	12	130.67	土	1.50	5.20	3.98	122.0?	139.0
Max. cranial breadth		168.88	土	1.63	5.90	3.49	157.0	181.0
Nasion-inion length	13		+	1.10	3.47	2.72	124.0	135.0
Basion-bregma height	10	127.60	<u>+</u>	0.80	3.08	3.31	89.0	100.0
Min. frontal breadth	15	93.10		1.26	4.71	4.23	104.0	
Vertical porion height	14	111.25	土		12.92	3.53	346.0	120.0
Median sagittal arc	11	366.45	土	3.89	14.54	0,00	040.0	387.0

TABLE 39-Continued

Characters	n	Mean	土	S. E.	S. D.	c.v.	Min.	Max.
Vertical transversal arc	12	303.92	±	3.33	11.51	3.79	289.0	325.0
Horizontal circumference	12	502.42	\pm	3.33	11.52	2.29	483.0	528.0
-Cranial module	8	146.40	\pm	0.86	2.43	1.66	143.50	150.17
Calculated cranial capacity	12	1285.31	\pm	32.94	113.97	8.87	1145.86	1542.89
Length-breadth index	12	72.64	\pm	1.12	3.89	5.36	66.30?	80.70
Length-height index	9	70.34	\pm	0.63	1.90	2.70	68.63	73.60
Breadth-height index	8	99.23	\pm	1.53	4.34	4.37	92.70	105.56?
Length-auricular height index	13	61.90	\pm	0.60	2.15	3.47	58.70	65.27
Breadth-auricular height index	12	85.47	\pm	0.94	3.26	3.81	80.43	90.49
Tr. fronto-parietal index	11	71.48	土	1.28	4.24	5.93	64.49	80.007

Table 40 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Splanchnocranium

n	Mean	+	S.E.	S.D.	c.v.	Min.	Max.
6	94.42	\pm	2.22	5.44	5.76	0.88	101.0
14	65.11	\pm	0.94	3.52	5.41	60.0	72.0
3	111.67	土	4.18	7.23	6.47	107.0	120.0
7	124.57	土	1.11	2.94	2.36	121.0	128.0
17	47.97	±		2.89	6.02	42.0	53.0
17				1.70	6.85	22.5	28.5
				2.37	12.83	14.0	22.5
					7.29	36.0?	46.0
					5:33	37.0	44.0
8"					6.93	31.0	39.0
					6.06	31.0	38.0
					6.04		59.0
					3.67		65.5
					7.11		48.0
					2.18		41.0
			0.20				
			0.40				
							52.07
							91.67?
		土					93.24
		土	1.34		10.66	43.40	67.86
	113.38	上	2.18		6 .3 9	10 5.3 6	128.00
11	89.76	土	1.81	6.00	6.68	8 3.33	100.00
	6 14 3 7	6 94.42 14 65.11 3 111.67 7 124.57 17 47.97 17 24.82 15 18.47 16 40.88 14 40.75 16 33.91 15 33.97 11 54.68 11 61.82 11 44.56 11 39.82 1 89.26 5 51.29 15 83.29 14 83.27 17 51.99 11 113.38	6 94.42 ± 14 65.11 ± 3 111.67 ± 7 124.57 ± 17 47.97 ± 17 24.82 ± 15 18.47 ± 16 40.88 ± 14 40.75 ± 16 33.91 ± 15 33.97 ± 11 54.68 ± 11 61.82 ± 11 44.56 ± 11 39.82 ± 1 89.26 5 51.29 ± 15 83.29 ± 14 83.27 ± 17 51.99 ± 11 113.38 ‡	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$

TABLE 41

ADULT FEMALE COMBINED

Statistical Constants of the Indices of the Whole Skull

Characters	n	Mean	±	S. E.	S. D.	c.v.	Min.	Max.
Tr. cranio-facial index	6	95.87		1.51	3.71	3.87	89.13	98.46
Vert. cranio-facial index	7	51.71	\pm	1.35	3.57	6.90	48.00	57.60
Long. cranio-facial index	4	50.64	±	1.44	2.89	5.71	46.70	53.65
Jugo-frontal index	7	73.77	\pm	0.92	2.44	3.31	70.87	76.86?

Characters	n	Mean	±	S. E.	S. D.	c. v.	Min.	Max.
Bigonial breadth	6	81.00	土	2.82	6.90	8.52	70.0	89.0
Bicondylar breadth	3	108.33	\pm	4.34	7.51	6.93	104.0	117.0
Ht. of mandibular ramus	5	59.00	\pm	0.88	1.97	3.34	57.5	62.0
Max. breadth of mandibular ramus	5	39.90	\pm	1.23	2.75	6.89	38.0	44.0
Min. breadth of mandibular ramus	6	31.50	\pm	1.09	2.68	8.51	29.0	36.0
Ht. at mandibular symphysis	7	30.00	\pm	1.68	4.44	14.80	24.0	36.0
Mandibular length	7	75.00	\pm	2.02	5 .35	7.13	66.0	81.0
Mandibular angle	5	118.70	\pm	0.84	1.89	1.59.	116°	120.5°
Mandibular index	3	73.75	\pm	2.82	4.87	6.60	68.38	77.88?
Breadth index of mandible	3	73.11	\pm	2.91	5.03	6.88	67.31	76.07
Jugo-mandibular index	2	63.96					57.85	70.08

TABLE 43

ADULT FEMALE COMBINED

Statistical Constants of the Measurements and Indices
of Permanent Maxillary Molar Teeth

Characters	n	Mean		S. E.	-	c. v.	Min.	Max.
Mesiodistal crown diam. of $\mathrm{M_{f}}$	8	10.06	土	0.06	0.17	1.69	10.00	10.50
Labiolingual crown diam, of M _I	7	10.50	\pm	0.40	1.05	10.00	9.00	12.00
Mesiodistal crown diam. of M ₂	9	9.56	\pm	0.27	0.82	8.58	8.00	10.75
Labiolingual crown diam. of M_2	9	9.78	土	0.42	1.25	12.78	8.00	11.25
Mesiodistal crown diam. of M ₃	7	8.39	\pm	0.38	1.00	11.92	7.00	10.00
Labiolingual crown diam. of M ₃	7	8.46	土	0.50	1.33	15.72	6.50	10.00
Crown index of M ₁	7	104.25	土	3.88	10.28	9.86	90.00	120.00
Crown index of M ₂	9	102.54	\pm	2.36	7.09	6.91	88.89	113.16
Crown index of M ₃	7	100.68	土	2.52	6.69	6.64	92.86	111.76
							· · · · · · · · · · · · · · · · · · ·	

TABLE 44 ADULT FEMALE COMBINED

Statistical Constants of the Measurements and Indices of Permanent Mandibular Molar Teeth

								_
Characters	n	Mean	土	S.E.	S.D.	c.v.	Min.	Max.
Mesiodistal crown diam. of M_1	3	10.67	±	0.16	0.28	2.62	10.50	11.00
Labiolingual crown diam, of M_1	3	10.08	土	0.08	0.14	1.39	10.00	10.25
Mesiodistal crown diam. of M2	3	9.58	土	0.36	0.63	6.58	9.00	10.25
Labiolingual crown diam. of M_2	4	9.25	\pm	0.66	1.32	14.27	7.50	10.50
Mesiodistal crown diam. of M_3	4	9.19	土	0.34	0.69	7.51	8.50	10.00
Labiolingual crown diam, of $ m M_3$	4	8.50	土	0.64	1.29	15.18	7.00	10.00
Crown index of M_1	, 3	94.70	土	2.06	3.56	3.76	90.91	97.96
Crown index of M2	3	94.01	\pm	8.53	14.75	15.69	83.33	110.84
Crown index of M_3	4	92.43	±	5.28	10.55	11.41	80.06	105.56

AREA G 289

From Area G 289 in Division I, II and III altogether twenty-three skulls and some post-cranial bones were recovered, which were found in a tightly packed condition lying in a narrow trench. The bones lay in a confused mass intermingled with animal bones. These crania represent nine adult males, four adult females, four adults (unsexed) and six children. Not all these skeletons are well preserved to permit study, only a small series of ten adults (seven males and three females) and five children's crania are found in restorable and measureable condition. Adult skulls only have been considered for evaluating the physical characteristics. The adult crania featured in this report may be classified by age and sex as follows.

TABLE 45
SEXWISE AGE DISTRIBUTION OF AREA G SKELETON

AGE	LESS THAN 25 YEARS	25-30 YEARS	30-40 YEARS
MALE	_	II S 18, II S 42, III S 2, III S 22, III S 23, III S 47	I S 11
FEMALE	I S 13, III S 21	II S 5	


The proportion of males is more than double than that of females. The markedly unusual sex-ratio is apparently fortuitous. The age distribution indicates that all the male individuals were in their prime of life within the age-group of twenty-five to thirty excepting one individual, who was thirty to forty years of age. Of the three females, two were young adults less than twenty-five years, while the remaining one was twenty-five to thirty years of age. Combined sex represents a total of 70% individual within twenty-five to thirty years age-group class. On a close scrutiny of the available skulls it was found that many of them have cut marks, injuries and abrasions (Pl. XXXI Figs. 1-3). Children's crania numbering five only, whose ages range from four to ten years have not been taken into account for determining their physical types. However, their measurements have been incorporated in the Collective Table C. Besides, measurements of a well-preserved mandible, which could not be associated with any of the adult cranium considered here, are recorded in the Collective Table F under the heading 'Remaining Mandibles'. The age of the children's crania are:

III S 1	ca.	4-5 years
III S 4	ca.	5-6 years
I S 1	ca.	6 years
III S 8	ca.	6-7 years
I S 15	ca.	9-10 years.

A. CRANIA

SKL. I S 11 [Figs. 58, 59; Pl. XXXVIII]

The skull is that of an adult male. The age of the individual is about 35 to 40 years, part of the coronal and sagittal suture being ossified. The skull is in a good state of preservation with its mandible. A distinct inclined and transverse cut at the nasal root by some sharp weapon is evidenced. The skull is big with moderate supraorbital ridges, strong mastoids with prominent supramastoid crests and well-developed muscular attachments. Post-auricular distance is nearly equal to pre-auricular. The skull is dolichocranic (L-B Index 71.62) and orthocranic (L-AH Index 61.01). The cranial contour in norma verticalis clearly corresponds with pentagonoides outline and is expanded at the parietals. Seen from the side, the forehead is slightly receding and turns to a convex vertex which slopes evenly down to moderately prominent tuber occipitale. In norma facialis the face is of medium height and breadth. Orbits appear to be rather low, chamaeconchic (OI 74.42) at right and mesoconchic (OI 79.01) at left. Right zygoma is evenly arched, while left is missing. The nose is of moderate length and falls in the messorrhine class (NI 50.98?). The palate is deep and short, the dental arcade being paraboloid in shape. All the maxillary teeth excepting left Ma have erupted and show second-degree wear.



The mandible is large and complete with all the sixteen teeth intact, and crowns of the teeth show considerable attrition. The bigonial diameter is not very wide (85 mm) and the mandibular angle seems to be evenly rounded. The height of the mandibular ramus is 67 mm and maximum breadth of the ramus measures 47 mm in width—giving a square appearance of the ascending ramus, emphasized by moderately deep incisura or sigmoid notch. Chin is not prominent but rounded, and height at mandibular symphysis is 35 mm.




Fig. 59 Skl. I S 11




Fig. 60 Skl. II S 18

 $\textbf{MID-SAGITTAL} \quad \textbf{CURVE} \quad (^{1}/_{2} \quad \text{NATURAL SIZE})$

MEMOIR: No. 9-

SKL. II S 18 [Fig. 60; Pl. XXX]

The skull belongs to an adult male. It has a horizontal crack below the infraorbitals. The skull was recovered in an excellent condition. Supraorbital ridges are not marked but big mastoids and medium muscular impressions are exhibited. Left lacrimal groove is deep and wide. None of the cranial sutures is ossified, but the *synchondrosis* spheno-basilaris is united. The skull is subbrachycranic (L-B Index 79.33) and hypsicranic (L-AH Index 64.25). The head shape seen from above is *sphenoides*, being broad in front and expanded at the parietals. Seen from the side, the forehead rises vertically and sweeps backwards into the general contour and merges with the rounded occiput. Post-orbital constriction is absent, the temporal region being well filled. The forehead is wide and the face is medium in height, the superior facial index being mesen (51.56). The orbits are rectangular which slope laterally and downward and fall under chamaeconchic class (right OI 70.24 left OI 66.29). The nose is mesorrhine (NI 49.02). Dentition is well preserved, shows second-degree wear, the arch being evenly *paraboloid* with a deep palate. All the maxillary teeth are in their sockets, excepting two left incisors, the crowns of which are broken. There are signs of caries in the first and second molars on right side as well as second molar on the left side. The calculated cranial capacity is 1401.31 cubic centimeter.

SKL. II S 42 [Figs. 61, 64; Pl. XXXI: 3]

The skull of an adult male. The skull is complete and survived in good condition excepting for a depressed horizontal fracture on its left temporal region possibly due to injury. Superciliary ridges are moderately developed, mastoid processes of moderate size and the muscular attachments are well





Fig. 61

developed. Vault sutures are open and spheno-basilaris is ossified. The skull is dolichocranic (L-B Index 72.47), and high-vaulted, length-height index hypsicranic (78.37). The vertical porion height of the skull is approximately 116 mm; the measurement could not be considered for indexing as it was taken on the reconstructed vertex region. Norma verticalis is ovoides in outline with some degree of bulging at the parietals. Seen from the side, the forehead is vertical and from ophryon it passes backwards into a full curve and merges with the moderately protruding occiput. Forehead is narrow and face moderately narrow. Right zygomatic arch is slender and evenly curved while the left one is missing. Nasal root is narrow, the interior interorbital space being 17 mm and the nasal depression is shallow. In profile, the bridge of the nose is somewhat straight and prominent, nose



Norma Lateralis

DIOPTOGRAPH TRACING (WASH) : MALE SKL. II S 18 (Natural Size)



Fig. 1 Skl. I S 1




Fig. 2 Skl. III S 4




Fig. 3 Skl. II S 42

Figs. 1-3 Showing cranial injuries: Area G.

being mesorrhine (NI 48.54). Subnasal prognthism is present to a slight degree and the meatus lying nearly midway between nasion and lambda. Palate is short, wide and moderately deep; dental arcade is paraboloid. All the maxillary teeth are erupted and intact but very much eroded.

SKL. III S 2 [Pl. XL]

The skull is that of an adult male. Supraorbital ridges are slightly developed. Post-mastoid portion of the occiput and contiguous part of the basi cranii are missing. The skull appears to have been subjected to the destructive action of saltpetre. Some scratches are noticed on the wide forehead. In the lambdoid suture two wormian bones are present. Basilar suture is united but the coronal and sagittal sutures are clearly open. The skull is brachycranic (L-B Index 81.29), hypsicranic (L-AH Index 67.25) and metriocranic (B-AH Index 82.73). From the top, contour in norma verticalis resembles with sphenoides type, being expanded at the parietals. Zygomatic arch is slightly jutting out from the vertical contour. From the side, the forehead is nearly straight. The orbits tend to be rectangular and are both mesoconchic (right OI 80.00, left OI 77.11). The palate is short, wide, deep and paraboloid. The anterior alveolar process of the maxilla is broken.

SKL. III S 22 [Figs. 62, 63; Pl. XXXIX]

The skull is undoubtedly male. Supraorbital ridges are prominent mesially, masteids being strong with well-developed muscular impressions. None of the cranial sutures is closed but sphenobasilaris is united. These, together with the state of dentition suggest that the skull belongs to an adult individual perhaps 25 to 30 years of age. Abrasions are noticed on the posterior part of both the parietals. Trace of metopic suture is noticed which extends from the nasion to glabella. Greater








Fig. 63

part of the squama of the occipital bone is missing. Forehead is wide and exhibits some degree of sloping. The skull is subbrachycranic (L-B Index 79.12) and orthocranic (L-AH 59.89). The shape of the head from the top is somewhat sphenoides in outline produced by swollen parietals. Face is moderate in height and width falling under mesen (superior facial index 50.00). Prenasal sulcus is present and slight degree of subnasal prognathism is observed when viewed from the side. Well-developed incisor fossae are present. The orbits are rectangular, low and medium and chamaeconchic




Fig. 64 Skl. II S 42




Fig. 65 Skl. III S 47

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

(OI 75.00) at the right, mesoconchic (OI 82.93) at the left. The depression of the nasal root is shallow with a relatively broad interorbital space; in profile the nose is concave. The dental arcade is broad and *paraboloid*. Excepting the incisors all the teeth are in their sockets, which are in very much worn state.

SKL. III S 23 [Pl. IXL: 1 & 2]

The skull is that of an adult male having moderately developed supraorbital ridges above the medial angle of each orbit, large right mastoid with strong supramastoid crests (left mastoid missing) and well-marked muscular impressions. Skull is complete with its mandible. Slight abrasion is noticed on the right frontal. The upper portion of the right half of the squama of the frontal bone is missing. The cusps of the teeth are partially worn. Trace of commencing synostosis on the sagittal suture is observed while coronal is open. Post-auricular development is nearly equal to pre-auricular. The skull is dolichocranic (L-B Index 72.77), orthocranic (L-AH Index 61.26 and L-H Index 72.25). Seen from above, the skull presents a pentagonoides shape, the greatest breadth being at the bi-parietals. Seen from the side, the narrow forehead is slightly receding and the occiput moderately projecting. Sagittal curve is full. Depression at the nasal root is shallow; concave-convex nasal bridge and slight subnasal prognathism are in evidence. The face is longish, the superior facial index being lepten (56.91). The nose is mesorrhine (NI 49.52), the bridge is high pitched, the tip slightly twisted to right and narrow at the root, anterior interorbital space being 17 mm. Orbits are more or less squarish and both are mesoconchic (right OI 82.05 and left OI 77.22). Maxillary teeth are mostly intact. Molars highly abraded (dentine exposed on cusps).

The mandible is of moderate size. The massesteric and pterygoidal areas are not strongly marked. The chin is prominent. The minimum breadth of the ascending ramus is 32 mm. The width of the dental arcade measured between the outer margin of the second molar is 62 mm and bigonial breadth is 93 mm. All the mandibular teeth excepting left central incisor are present in sound condition.

SKL. III S 47 [Fig. 65; Pls. XXXII, XXXIII]

Skull is that of an adult male. The age at death appears to have been around 25-30 years. Supraorbital ridges are perceptible above the medial angle of each orbit; mastoids and paramastoids are moderately developed, and the muscular attachments are well marked. Abrasions on the frontal and left parietal region are noticed. Left canine fossa is markedly deep. Closing of sagittal suture at obelion is observed. The skull is subbrachycranic (L-B Index 78.41) and hypsicranic (L-AH Index 65.34). The cranial capacity is 1355.57 cc. Viewed from above, the skull is byrsoides in outline with zygomatic arches bowed out laterally to the vertical contour. The valut is well filled at the temporal region. Seen from the side, the forehead is vertical and goes back in an even curve to meet the occiput which is only moderately protruding. The height of the face is less than half of its breadth, the superior facial index is euryen (48.24). The nasal profile is seen to be concave without any depression at the root. Moderate subnasal prognathism is evidenced. The incisors are protruding forwards. The orbits are somewhat squarish in form. Dental arcade is paraboloid, narrowed in front.




Fig. 66 Skl. II S 5





Fig. 67 Skl. III S 21

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)



Norma Lateralis

pioptograph tracing (Wash) : Male skl. III s 47 (Natural Size)



Norma Frontalis

DIOPTOGRAPH TRACING (WASH) : MALE SKL. III S 47 (Natural Size)

SKL. I S 13

The skull is that of female adult, probably at the margin of juvenile and adult age. Size is small having ill-marked supraorbital ridges and small mastoids. Superior orbital margins are sharp. Major portion of the cranial vault, specially the entire left side is badly damaged and missing; only the facial portion and the base are present. Incisors protruding forwards and the lateral incisors split vertically. Important measurements, such as head length and head breadth could not be taken. Hence, type-diagnosis of this skull is made on morphological and not on metric grounds. Observed from the top, the fragmentary remnants of the bones present a delichoid and ellipsoides shape. The forehead is vertical while the occiput is rounded. The orbits are somewhat squarish in form and are mesoconchic (right OI 80.00 and left OI 81.69).

SKL. II S 5 [Fig. 66]

The skull of an adult female. Supraorbital ridges are ill developed mesially and both the mastoids are small. Contour of the face and calva is smooth and gracile. Crack mark is observed on both the right and left parietals. Basilar suture is united. The skull is mesocranic (L-B Index 77.14) and orthocranic (L-AH Index 62.86). Cranial contour in norma verticalis is byrsoides, narrow in front and expanded at the parietals. Forehead is vertical, well arched and narrow. Occiput is rounded. Face is long and narrow in mid-facial width having slender zygomatic arches. The face is high belonging to the lepten class. The bridge of the nose is straight having a narrow nasal root. Nose is mesorrhine (NI 48.94). The orbits are more or less rectangular and mesoconchic. The palate is moderately deep and paraboloid. The cusps of the teeth are not much worn down.

SKL. III S 21 [Fig. 67]

The skull is that of young female. Basilar, sagittal and coronal sutures are open and there is no evidence of beginning union. The supraorbital ridges tended to be poorly developed mesially, mastoids small and muscular attachements undeveloped giving a gracile appearance to the skull. Viewed from top, the skull is ellipsoides in shape. The skull is dolichocranic (L-B Index 71.84) and orthocranic (L-AH Index 60.34). Viewed from norma lateralis sinistra, forehead appears to be slightly receding, vertex moderately arched and the union of the parietal and occipital bones is marked by a lambdoid depression. There is also evidence of some amount of subnasal prognatium. In norma facialis the face is seen to be moderate in height and breadth and superior facial index is middle (mesen 54.55). Orbits are horizontal and squarish in form. The nasal root is only slightly depressed, ridge short and concave; and nasal aperture is moderately high and broad, mesorrhine (NI 49.48). Norma occipitalis is house-shaped with converging side walls. The shape of the dental arcade is paraboloid and somewhat narrow anteriorly. Fourteen maxillary teeth have erupted and there is no indication of the eruption of upper 3rd molars. Tooth wear is not very marked.

ANALYSIS

The skulls with marks of injury, signs of wounds and abrasions in some, found in a tightly packed condition inside a trench at Area & 289, indicate a deliberate and hasty nature of inhumation

unlike any other regular burial. Out of twenty-three crania only 10 well-preserved adult skulls could be considered for attempting diagnosis of physical type. The children's crania numbering five only have been excluded from the present consideration.

The foregoing observations on the adult skulls and their metric data (Collective Table B-C) are quite distinctive for identifying their physical types. Accordingly, two morphological types have been derived, namely, long-headed Type $A_{\rm l}$ and round-headed Type $B_{\rm l}$. Type-wise and sex-wise distribution of the adult skulls are as follows:

Тур	De A,	Туре	B_{i}
Male	Female	Male	Female
I S 11	IS 13	II S 18	II S 5
II S 42	III S 21	III S 2	
III S 23		III S 22	
111 5 25		III S 47	

TYPE A

In Tables 46-51 the mean values with standard errors, ranges and indices of males of Type A_1 are given; and in Tables 62-65 those of females are given. Photographs and dioptograph drawings of two skulls (one male and the other female) representing Type A_1 of Area G are produced in Plates XXXIV. XXXV and in figures 68-70, 71-73.

The essential cranio-facial features of the long-headed Type A: are shown below:

	MALE		FEMAL	.E
Length-breadth index Length-height index Breadth-height index Length-auricular height index Superior facial index Orbital index Nasal index Palatal index Trans, fronto-parietal index	Dolichocranic Orthocranic Acrocranic Orthocranic Lepten Mesoconchic Mesorrhine Brachystaphylin Eurymetopic	(72.29 ± 0.35) (72.58 ± 3.26) (100.37 ± 4.21) (61.14) (56.91) (79.91 ± 1.59) (49.68 ± 0.71) (87.60 ± 6.39) (69.02 ± 1.47)	Dolichocranic Orthocranic Acrocranic Orthocranic Mesen Mesoconchic Mesorrhine Leptostaphylin Eurymetopic	(71.84) (70.69) (98.40) (60.34) (54.55) (83.46 ± 1.55) (49.48) (79.17) (72.00)

It is worthy to note that despite the sex differences in absolute dimensions, there is a close harmony in cranio-facial proportions of Type A₁ skulls. The mean value shows that both male and female are delichocranic (the skulls representing Type A₁ lie on the cranial index peak of 71-72). The average values for length-height index is 72.33 for males and 70.69 for females. By far the most common occurrence in the Type A₂ skulls is delicho-ortho-acrocrany.

The mean absolute measurements of both the sexes are as follows: Cranial length 185.83~mm for males and 174.00~mm for females; cranial breadth 134.33~mm for males and 125.00~mm for females; auricular height 116.00~mm for males and 107.00~for females; nasal height 51.67~mm for males and 45.25~mm for females; nasal breadth 25.67~mm for males and 23.75~mm for females. Some of the important morphological characters of both the sexes of Type A_1 are furnished below:

Andreas de management (all al la company)	TYPE A	
CHARACTERS	MALE	FEMALE
Development of supraorbital ridges	Medium	Slight
Slope of the forehead	Slightly receding to Vertical	Slightly receding to Vertical
Depression of the nase! root	Shallow	Shallow
Protuberence of the occipital region	Moderately protruding	Rounded
Mastoid processes	Medium	Small
Nuchal plane	Muscular ridges Medium to Slight	Muscular ridges Slight

The above observations in addition to other traits exhibit some common morphological features of the skulls of both sexes of Type A. The skulls of this type are marked by general gracility, medium to slight supraorbital ridges, slightly receding to vertical forehead, medium to narrow face with square and horizontal orbits, medium to narrow nose, shallow nasion depression and projecting occiput.

TYPE B,

Five skulls, comprising four adult males and one female are attributable to this physical type.

In Tables 52-55 the mean values with standard errors, ranges and indices of males of Type B₁ are given and those of females in Tables 66-69. Photographs and dioptograph drawings of two skulls (one male and the other female) representing Type B₁ of Area G are produced in Plates XXXVI, XXXVII and in Figures 74-76, 77-79.

The essential cranio-facial features of the round-headed Type B: are given below:

MALE			FEMALE	
Length-breath index Length-height index Breadili-height index Length-auricular height index Superior facial index Orbital index Nasal index Palatal index Trans. fronto-parietal index	Mesocranic Hypsicranic Metriocranic Hypsicranic Euryn Mesoconchic Chamaerrhine Brachystaphylin Eurymetopic	(79.54 ± 0.62) (75.01 ± 2.08) (94.28 ± 2.10) (64.18 ± 1.56) (49.93 ± 0.96) (77.50 ± 2.49) (51.46 ± 1.18) (95.70 ± 6.15) (73.10 ± 1.30)	Mesocranic Orthocranic Metriocranic Orthocranic Lepten Mesoconchic Mesorrhine Mesostaphylin Metriometopic	(77.14) (72.00) (93.33) (62.86) (56.54) (77.60) (48.94) (81.63) (68.15)




Fig. 68 Norma Frontalis







Fig. 69 Norma Verticalis



Figs. 68, 69 and 70. Dioptographic contours of Type A: Male (Skl. III S 23) of Area G







Norma Verticalis



Norma Lateralis

Photographs of Type A_1 Male $[\,Skl.\,\,III\,\,S\,\,23\,\,]$ of Area G



Norma Frontalis



Norma Verticalis



Norma Lateralis

Photographs of Type A. Female [Skl. III S 21] of Area G




Fig. 71 Norma Frontalis




Fig. 72 Norma Verticalis




Fig. 73 Norma Lateralis

Figs. 71, 72 and 73. Dioptographic contours of Type A1 Female [Skl. III S 21] of Area G 13

Both males and females are mesocranic, the cranial index of the females being slightly higher than the males. The vault is relatively high for males (hypsicranic). In major cranial proportions the Type B₁ skulls are mesocranic or subbrachycranic, ortho- to hypsicranic and metriocranic. With the exception of one skull rest of this series lie on the border of meso- and chamaerrhine with an index around 49-51.

The average values of absolute dimensions of both the sexes belonging to Type B_1 are: Cranial length 177.00 mm for males and 175 mm for females; cranial breadth 140.75 mm for males and 135.00 mm for females; nasal height 50.00 mm for males and 47.00 for female; nasal breadth 25.75 for males and 23.00 for female.

On a close examination of the skulls belonging to Type B_{I} , the following important morphological features of both the sexes may be tabulated as below:

	TYPE B				
CHARACTERS	MALE	FEMALE			
Development of supraorbital ridges	Medium to Slight	Slight			
Slope of the forehead	Slightly receding to Vertical	Vertical			
Depression of the nasal root	Shallow to Absent	Absent			
Protuberence of the occipital region	Moderately protruding to Rounded	Rounded			
Contour of the skull from top	Sphenoides	Byrsoides			
Zygomatic arch	Phaenozygous and Orthozygous	Orthozygous			

Taking into consideration the morphological features of the round-headed skulls certain traits are well delineated. The crania are characterized by having relatively high vault, broad and moderately protruded to rounded occiput, well-filled temporal region, slightly receding to vertical and wide fore-head, byrsoides and sphenoides cranial contour, medium orbits and medium to heavy muscular ridges.

Morphology and cranial proportions of Type B₁ of Area G are in close agreement with the round-headed Alpine type of Hissar III (Krogman 1940 : 22-23).

DISCUSSION

Thus, on the evidence of figures and form two distinct physical types are observed in Area G

Area G may well be compared with the Type A_1 of Cemetery R 37. The calvarial part of the Type A_1 skull from Area G is long, narrow and moderately high, so that the three cardinal cranial indices (length-breadth, length-height and breadth-height) almost coincide with the mean values of the Type A_1 of Cemetery R 37. At the same time absolute dimensions and indices of the facial part also show close resemblance. Both are long-headed, having smooth and gracile contours, faintly developed glabello-superciliary region; the entire physical structure being finer and weaker, with medium orbits, mesorrhine nose, medium face with moderate to narrow mid-facial width and protruding occiput. This gracile long-headed Type A_1 population seems to have continued to survive at Area G from the earlier population represented in Cemetery R 37. Thus, Type A_1 of Area G may probably be regarded as the basic type and the survival of the gracile long-headed type (Type A_1) of Cemetery R 37.

Round-headed Type B: of Area G is distinct in physical features. The skulls show some well delinated features, having moderately high and full vault, well-filled temporal region, vertical contour varying between byrsoid and sphenoid with slightly receding to vertical wide forehead and mildly curved occipital bone.

Metrically the Type B₁ skulls shows the following characteristics: On an average it is mesocranic or low brachycranic, orthoto hypsicranic, mesoconchic orbit, high mesorrhine to low chamaerrhine. This round-headed population appears to be a new-comer in the scene who had so far, shown no evidence of its presence in the earlier period. The fact, naturally, leads us to postulate that an extraneous group is represented by this round-headed population. Haddon (1919: 27) held that in ancient times certain unrecorded brachycephalic people migrated in India.

The whole collection of the Area G was discovered in a confused mass in a tightly packed heap unlike any other regular burial, and most of the skulls bear evidences of positive injury. These lead to assume the possibility of a sudden attack by a round-headed invading people on the long-headed autochthones.

B. EXTREMITY BONES

The skeletal materials of Area G, as already noted, were not recovered from regular burials and as such the bones were disarticulated and lay in a disorderly condition. The bones therefore could not be associated with respective skull. All the extremity bones are fragmentary in nature but for one radius (Skl. II S 51) belonging to an adult individual. The scarcity of long bones in good condition baffled all attempts of measurement.




Fig. 74 Norma Frontalis




Fig. 75 Norma Verticalis





Fig. 76 Norma Lateralis

Figs. 74, 75 and 76. Dioptographic contours of Type B. Male [Skl. III S 47] of Area G



Norma Frontalis

Norma Verticalis



Norma Lateralis


Photographs of Type B_1 Male [Skl. III S 47] of Area G



Norma Frontalis



Norma Verticalis



Photographs of Type B Female [Skl. II S 5] of Area G




Fig. 77 Norma Frontalis




Fig. 78 Norma Verticalis




Fig. 79 Norma Lateralis

Figs. 77, 78 and 79. Dioptographic contours of Type B₁ Female (Skl. II S 5) of Area @

MEMOIR: No. 9

Basic measurements and index of one radius derived from them are shown below:

Skl. II S 51

Maximum length 243.0 mm
Physiological length 233.0 mm
Minimum circumference of
diaphysis 38.0 mm
Robusticity index 16.31

As the sex of the individual could not be identified from a single bone, stature has been calculated assuming it to be a male one and a female one separately. The results are shown below:

		ESTIMATED STATURE ACCORDING	то
	MANOUVRIER	PEARSON	DUPERTUIS AND HADDEN FORMULAE
			/
If a male radius	1666.00 mm	1654.10 mm	1726.92 mm
If a female radius	1675.50 mm	1624.59 mm	1717.07 mm

Had the individual happened to be a male one he was tall- or medium-statured, if a female she was very tall or tall.

STATISTICAL CONSTANT : AREA G

TABLE 46

ADULT MALE TYPE A₁

Mean Measurements and Indices of the Neurocranium

MARKET CONTROL OF STATE OF STA					
Characters	n	Mean	S.E.	Min.	Max.
Manager 1 1 and 1	0	107.00	0.00	150.0	101.0
Max. cranial length	3	185.83 ±	3.99	178.0	191.0
Max. cranial breadth	3	$134.33 \pm$	2.91	129.0	139.0
Nasion-inion length	3	$171.67 \pm$	2.85	166.0	175.0
Basion-bregma height	3	$134.67 ~\pm$	4.11	126.5	139.5
Min. frontal breadth	3	$92.67 \pm$	1.77	90.0?	96.0
Vertical porion height	2	116.00		115.0	117.0
Median sagittal arc	3	$382.33 \pm$	10.10	3 6 3. 0	397.0
Vert. transversal arc	3	$307.00 \pm$	2.31	303.0	311.0
Horizontal circumference	3	$512.67\ \pm$	8.88	495.0	523.0
Cranial module	3	$151.61 ~\pm$	2.23	148.83	156.0
Calculated cranial capacity	3	1298.95 ± 1	17.53	1265.39	1324.35
Length-breadth index	3	$72.29 \pm$	0.35	71.62	72.77
Length-height index	3	$72.58 \pm$	3.26	67.11	78.37
Breadth-height index	3	$100.37 \pm$	4.21	93.70	108.14
Length-auricular height index	2	61.14		61.01	61.26
Breadth-auricular height index	2	84.68		84.17	85.19
Tr. fronto-parietal index	3	$69.02 \pm$	1.47	66.19	71.11

TABLE 47 ${\tt ADULT\ MALE\ TYPE\ A}_1$ Mean Measurements and Indices of the Splanchnocranium

The state of the s					
Characters	n	Mean ±	S.E.	Min.	Max.
For manager and the second sec					
Prosthion-basion line	3	94.50 ±	0.76	9 3.5?	96.0
Nasion-prosthion line	3	68.50 ±	1.04	66 .5	70.0
Nasion-gnathion line	1	119.00			
Max. bizygomatic breadth	1	123.00			Stamone
Nasal height	3	51.67 ±	0.44	51.0?	52.5
Nasal breadth	3	$25.67 \pm$	0.34	25.0	26.0
Ant. inter-orbital breadth	3	17.33 ±	0.34	17.0	18.0
Orbital breadth (right)	3	40.67 ±	1.20	39.0	43.0
Orbital breadth (left)	3	39.33 ±	0.73	38.0	40.5
Orbital by A. (tabt)	3	32.17 ±	0.16	32.0	32.5
Orbital height (right)	3	31.67 ±	0.60	30 .5	32.5
Orbital height (left)	_	53.50 ±	2.47	49.0	57.5
Maxillo-alveoler length	3	64.67 ±	4.42	56.0	70.5
Maxillo-alveolar breadth	3	04.07 -	7.72	00.0	

TABLE 47 - Continued


				90 N N N N N N N N N N N N N N N N N N N	
Characters	n	Mean \pm	S.E.	Min.	Max.
water to the second of the sec					
Palatal length	3	$49.67 \ \pm$	1.86	46.0	52.0
Palatal breadth	3	$43.33 \pm$	2.34	39.0	47.0
Superior facial index	1	56.91		_	_
Orbital index (right)	3	$79.24 \pm$	2.42	74.42	82.05
Orbital index (left)	3	$80.59 \pm$	2.53	77.22	85.53
Nasal index	3	$49.68 \pm$	0.71	48.54	50.98?
Maxillo-alveolar index	3	$121.36 \ \pm$	9.86	103.70	137.76
Palatal index	3	$87.60 \pm$	6.39	75.00	95.65

TABLE 48 Mean Indices of the Whole Skull ADULT MALE TYPE A

Characters	n	Mean ±	S.E.	Min.	Max.
Tr. cranio-facial index	1	88.49		_	_
Vert. cranio-facial index	3	$50.98 \pm$	1.99	47.67	54.55?
Long. cranio-facial index	3	$50.89 \pm$	0.98	49.60?	52.81
Jugo-frontal index	1	74.80		-	_

TABLE 49 ADULT MALE TYPE A Mean Measurements and Indices of the Mandible


		•		and the same
Characters	n	Mean \pm S.E.	Min,	Max.
Bigonial breadth	2	89.00	85.0	93.0
Bicondylar breadth	2	117.50	115.0?	120.0
Ht. of mandibular ramus	2	61.75	56.5	67.0
Max. breadth of mandibular ramus	2	45.25	43.5	47.0
Min. breadth of mandibular ramus	2	34.50	32.0	37.0
Ht. at mandibular symphysis	2	35.00	35.0	35.0
Mandibular length	2	83.25	83.0	83.5
Mandibular angle	2	116.75°	113.5°	120°
Mandibular index	2	70.88	69.58	72.17?
Breadth index of mandible	2	75.85	70.83	80.87?
Jugo-mandibular index	1	75.61		




Frontal view



Vertical view



Left lateral view



Occipital view

SKL. I S 11 : AREA G



Fig. 1 Occlusal aspect



Fig. 2 Left lateral aspect



Fig. 3 Occlusal aspect




Fig. 4 Left lateral aspect

MANDIBLE : AREA G

TABLE 50 ${\bf ADULT\ MALE\ TYPE\ A}_1$ Mean Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean	土	S.E.	Min.	Max.
Mesiodistal crown diam. of M ₁	3	10.42	±	0.21	10.00	10.75
Labiolingual crown diam. of M_1	3	11.25	\pm	0.14	11.00	11.50
Mesiodistal crown diam, of M ₂	3	10.08	\pm	0.30	9.50	10.50
Labiolingual crown diam. of M ₂	3	11.67	\pm	0.51	10.75	12.50
Mesiodistal crown diam. of M ₃	3	9.25	\pm	0.24	8.75	9.50
Labiolingual crown diam. of M ₃	3	10.92	土	0.65	9.75	12.00
Crown index of M ₁	3	108.26	±	0.87	107.28	110.00
Crown index of M ₂	3	117.79	±	3.62	113.61	125.00
Crown index of M ₃	3	117.85	±	4.42	111.44	126.32

TABLE 51 ${\bf ADULT\ MALE\ TYPE\ A}_1$ Mean Measurements and Indices of Permanent Mandibular Molar Teeth

Characters	n	Mean	Min.	Max.
Mesiodistal crown diam. of M ₁	2	10.38	10:00	10.75
Labiolingual crown diam. of M ₁	2	11.00	10.00	12.00
Mesiodistal crown diam, of M ₂	2	10.12	9.25	11.00
Labiolingual crown diam. of $ m M_2$	2	10.62	9.25	12.00
Mesiodistal crown diam. of M ₃	2	9.88	9.50	10.25
Labiolingual crown diam. of M ₃	2	9.88	8.50	11.25
Crown index of M ₁	2	106.09	100.00	112.18
Crown index of M ₂	2	104.62	100.15	109.09
Crown index of M ₃	2	99.62	89:47	109.76

Table 52 ${\rm Adult\ male\ Type\ B}_1$ Mean Measurements and Indices of the Neuroeranium

The state of the s					the term of the second of the
Characters	n	Mean ±	S.E.	Min.	Max.
Max. cranial length	4	$177.00 \pm$	2.34	171.0	182.0
Max. cranial breadth	4	$140.75\ \pm$	1.38	138.0	144.0
Nasion-inion length	4	$166.50 \pm$	1.71	162.0	170.0
Basion-bregma height	4	$132.62\ \pm$	1.95	128.0	137.5

TABLE 52-Continued

Characters	n	Mean <u>⊣</u>	S.E.	Min.	Max.
Min. frontal breadth	4	102.88 ±	1.74	99.0	106.5
Vertical porion height	4	113.50	1.50	109.0	115.0
Median sagittal arc	3	363.00 ±	4.73	354.0	370.0
Vert. transversal arc	4	307.00 ±	3.39	300.0	315.0
Horizontal circumference	4	512.00 ±	1.00	505.0	523.0
Cranial module	4	150.12 ±	0.10	148.67	151.33
Calculated cranial capacity	4	1367.34 ±	14.38	1334.56	1401.31
Length-breadth index	4	79.54 ±	0.62	78.41	81.29
Length-height index	4	$75.01 \pm$	2.08	70.33	80.41
Breadth-height index	4	94.28 ±	2.10	88.89	98.92
Length-auricular height index	4	64.18 ±	1.56	59.89	67.25
Breadth-auricular height index	4	80.68 ±	1.74	75.69	
Tr. fronto-parietal index	4	$73.10 \pm$			83.33
			2.50	70.14	75.54

TABLE 53 ADULT MALE TYPE B Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	土	S.E.	Min.	Max.
Prosthion-basion line	3	95.00	+	2.09	92.0	99.0
Nasion-prosthion line	3	64.67		1.59	61.5	66.5
Max. bizygomatic breadth	3	129.50	±	1.76	127.5	133.0
Nasal height	4	50.00	土	1.29	47.0	53.0
Nasal breadth	4	25.75	土	1.11	24.0	29.0
Ant inter-orbital breadth	4	22.50	\pm	0.36	21.5	23.0
Orbital breadth (right)	4	41.50		0.96	40.0	44.0
Orbital breadth (left)	4	41.50		1.14	39.0	44.5
Orbital height (right)	4	31.62		0.74	29.5	33.0
Orbital height (left)	3	32.50		1.14	29.5	34.5
Maxillo-alveoler length	3	53.33		1.36	51.5	56.0
Maxillo-alveolar breadth	3	61. 8 3 44.33		2.62 2.34	57. 0 40 .0	66.0 48.0
Palatal length	4	42.25		2.34	38.0	48.0
Palatal breadth	3	49.93		0.96	48.24	51.56
Superior facial index Orbital index (right)	4	76.31		2.34	70.24	80.00
Orbital index (left)	Ä	78.70		4.74	66.29	88.46
Nasal index	4	51.46		1.18	49.02	54.72
Maxillo-alveolar index	2	111.14			110.68	111.61
Palatal index	3	95.70	\pm	6.15	85.42	106.67

TABLE 54

ADULT MALE TYPE B

Mean Indices of the Whole Shall

Mican	murces	OI	me	w noie	SKUII	

Characters	n	Mean ±	S.E.	Min.	Max.
Tr. cranio-facial index	3	9 1.62 ±	0.75	90.14	92.39
Vert. cranio-facial index	3	49.39 ±	1.55	46.59	51.95
Long. cranio-facial index	3	$53.08 \pm$	1.13	51.65	55.31
Jugo-frontal index	3	$78.93 \pm$	2.20	75.94	83.20

Table 55 ${\bf ADULT\ MALE\ TYPE\ B}_1$ Mean Measurements and Indices of the Permanent Maxillary Molar Teeth

Characters	n	Mean ±	S.E.	Min.	Max.
Mesiodistal crown diam. of M_{I}	4	10.69 ±	0.34	10.00	11.50
Labiolingual crown diam. of M ₁	4	11.75 ±	0.10	11.50	12.00
Mesiodistal crown diam. of M ₂	4	10.00 ±	0.18	9.50	10.25
Labiolingual crown diam. of M_2	4	$11.06 \pm$	0.33	10.50	11.75
Mesiodistal crown diam. of M ₃	4	8.62 ±	0.24	8.00	9.00
Labiolingual crown diam. of M_3	4	9.75 ±	0.36	9.00	10.50
Crown index of M ₁	4	$110.36 \pm$	4.39	100.00	120.00
Crown index of M ₂	4	$111.00 \pm$	2.76	103.35	115.07
Crown index of M ₃	4	113. 8 2 ±	4.08	103.56	123.58

TABLE 56

ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean ±	S. E.	S. D.	C. V.	Min.	Max.
Max. cranial length	7	180.79 ±	2.65	7.01	3.88	171.0	191.0
Max. cranial breadth	7	138.00 ±	1.85	4.90	3.55	129.0	1-14.0
Nasion-inion length	7	168.71 ±	1.75	4.64	2.75	162.0	175.0
Basion-bregma height	7	133.50 ±	1.91	5:07	3.80	126.5	139.5
Min. frontal breadth	7	98.50 ±	2.35	6.24	6.34	90.0?	106.5
Vertical perion height	6	114.33 ±	1.11	2.73	2.39	109.0	117.0
Longitudinal arc	6	$372.67 \pm$	6.60	16.16	4.34	354.0	397.0

TABLE 56-Continued

Characters	n	Mean <u>+</u> S	s. E. S. D.	c. v.	Min.	Max.
	= (100-11), (2011)			as the fire		
Vertical transversal arc	7	307.00 ± 3	2.01 5.32	1.73	300.0	315.0
Horizontal circumference	7	512.29 ± 4	4.08 10.81	2.11	495.0	523.0
Cranial module	7	150.76 ± 0	0.97 2.56	1.70	148.67	156.00
Calculated cranial capacity	7	1387.76 ± 13	5.84 41.98	3.03	1334.56	1465.78
Length-breadth index	7	76.43 ± 1	1.51 3.99	5.22	71.62	81.29
Length-height index	7	73.97 ± 1	1.57 4.57	6.18	67.11	80.41
Breadth-height index	7	96.89 ± 2	2.30 6.09	6.29	88.89	108.14
Length-auricular height index	6	63.17 ± 3	1.18 2.88	4.56	59.89	67.25
Breadth-auricular height index	6	82.02 ± 1	1.39 3.41	4.16	75.69	85.19
Tr. fronto-parietal index	7	71.36 ± 1	1.21 3.21	4.50	66.19	75.54

TABLE 57 ADULT MALE COMBINED Statistical Constants of the Measurements and Indices of the Splanchnocranium

Characters	n	Mean ±	S. E.	S. D.	c. v.	Min.	Max.
Prosthion-basion line	6	94.75 ±	1.00	2.44	2.58	92.0	99.0
Nasion-prosthion line	6	66.58 ±		2.96	4.45	61.5	70.0
Nasion-gnathion line	1	119.00	1,21	_			-
Bizygomatic breadth	4	127.88 ±	2.04	4.09	3.20	123.0	133.0
Nasal height	7	50.71 ±		2.08	4.10	47.0	53.0
Nasal breadth	7	25.71 ±	0	1.60	6.22	24.0	29.0
Ant. inter-orbital breadth	7	20.29 ±		2.83	13.95	17.0	23.0
Orbital breadth (right)	7	41.14 ±	2.01	1.87	4.55	39.0	44.0
Orbital breadth (left)	7	40.57 ±		2.11	5.20	38.0	44.5
Orbital height (right)	7	31. 86 ±		1.11	3.48	29.5	33.0
Orbital height (left)	7	$32.14 \pm$		1.77	5.51	29.5	34.5
Maxillo-alveolar length	6	$53.42 \pm$	1.26	3.09	5.78	49.0	57.5
Maxillo-alveolar breadth	6	63.25 ±	2.38	5.84	9.23	56.0	70.5
Palatal length	6	$47.00 \pm$	1.79	4.38	9.32	40.0	52.0
Palatal breadth	7	$42.71 \pm$	1.44	3.82	8.94	38.0	48.0
Superior facial index	4	51. 68 ±	1.87	3.74	7.24	48.24	56.91
Orbital index (right)	7	$77.57 \pm$	1.66	4.39	5.66	70.24	82.05
Orbital index (left)	7	79 .51 ±	2.73	7.23	9.09	66.29	88.46
Nasal index	7	50.69 ±	0.78	2.06	4.06	48.54	54.72
Maxillo-alveolar index	5	117.27 ‡	5.94	13.30	11.34	103.70	137.76
Palatal index	. 6	91.65 ±	3.98	9.74	10.63	75.00	106.67

TABLE 58

ADULT MALE COMBINED

Statistical Constants of the Indices of Whole Skull

Characters		n	Mean ±	S. E.	S. D.	c. v.	Min.	Max.
	a = 1							
Tr. cranio-facial index		4	90.84 ±	0.94	1.89	2.08	88.49	92.39
Vert. cranio-facial index		6	50.18 ±	1.18	2.90	5.78	46.59	54.55?
Long. cranio-facial index		6	$51.98 \pm$	0.83	2.03	3.91	49.60?	55.31
Jugo-frontal index		4	$77.90\ \pm$	1.86	3.72	4.78	74.80	83.20

TABLE 59

ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of the Mandible

Characters	n	Mean <u>+</u>	S. E.	S. D.	c. v.	Min.	Max.
Bigonial breadth	3	$91.00 \pm$	3.06	5.29	5.81	85.0	95.0
Bicondylar breadth	3	$119.50 ~\pm$	2.47	4.27	3.57	115.0?	123.5
Ht. of mandibular ramus	4	$57.38 \pm$	3.89	7.78	13.56	48.0	67.0
Max. breadth of mandibular ramus	4	$42.38 \pm$	1.82	3.64	8.59	39.0	47.0
Min. breadth of mandibular ramus	6	$32.92 \pm$	1.20	2.94	8.93	29 .5	37.0
Ht. at mandibular symphysis	4	$30.62 \pm$	3.06	6.13	20.02	22.0	35.0
Mandibular length	5	$78.50 \pm$	2.73	6.12	7.80	70.0?	83.5
Mandibular angle	3	$118.83^{\circ} \pm$	2.81	4.86	4.09	113.5°	123°
Mandibular index	2	70.88				69.58	72.17?
Breadth index of mandible	2	75.85				70.83	80.87?
	1	75.61			_	_	
Jugo-mandibular index							

TABLE 60
ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of Permanent Maxillary Molar Teeth

Characters **	n	Mean <u>+</u>	S. E.	S. D.	c.V.	Min.	Max,
	7 7 7 7 7 7 7	10.57 ± 11.54 ± 10.04 ± 11.32 ± 8.89 ± 10.25 ± 109.46 ± 113.91 ±	0.21 0.12 0.15 0.29 0.20 0.39 2.40 2.43 2.86	0.55 0.33 0.40 0.76 0.54 1.04 6.37 6.44 7.58	5.20 2.86 3.98 6.71 6.07 10.15 5.82 5.65 6.56	10.00 11.00 9.50 10.50 8.00 9.00 100.00 103.35	11.50 12.00 10.50 12.50 9.50 12.00 120.00
Crown index of M ₃	7	115.55 ±	2.00		9.56	103.56	126.32

TABLE 61

ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of Permanent Mandibular Molar Teeth

Characters	n	Mean ±	S. E.	S. D.	c. v.	Min.	Max.
and the second s							
Mesiodistal crown diam. of M_1	5	$10.30 \pm$	0.12	0.28	2.72	10.00	10.75
Labiolingual crown diam. of M_1	5	$10.05 \pm$	0.50	1.12	11.14	9.25	12.00
Mesiodistal crown diam. of M2	6	$9.67 \pm$	0.38	0.94	9.72	8.50	11.00
Labiolingual crown diam. of M2	6	$9.62 \pm$	0.58	1.41	14.66	8.00	12.00
Mesiodistal crown diam. of M ₃	5	$9.75 \pm$	0.31	0.69	7.08	8.75	10.50
Labiolingual crown diam. of M ₃	5	$9.25 \pm$	0.67	1.51	16.32	7.25	11.25
Crown index of M_1	5	$97.60 \pm$	3.98	8.92	9.14	90.36	112.18
Clown index of M ₂	6	99.83 ±	2.92	7.16	7.17	88.89	109.09
Crown index of M_3	5	94.47 ±	4.42	9.89	10.47	83.01	109.76

Characters	, n	Mean \pm S.E.	Min.	Max.
The second length		1774.0		
Max. cranial length	1	174.0		
Max. cranial breadth	1	125.0		_
Nasion-inion length	1	157.0		
Basion-bregma height	1	123.0	- •	
Min. frontal breadth	2	88.00	86.0	90.0
Vertical porion height	2	107.00	105.0	109.0?
Median sagittal arc	1	354.0		
Vertical transversal arc	1	277.0	40-mar/80	
Horizontal circumference	1	486.0	and the same of	
Cranial module	1	140.67		
Calculated cranial capacity	1	1139.85		
Length-breadth index	1	71.84		
Length-height index	1	70.69		
Breadth-height index	1	98.40	*****	
Length-auricular height index	1	60.34		-
Breadth-auricular height index	1	84.00		
Tr. fronto-parietal index	1	72.00	_	guarent.

Characters	n	Mean ± S.E.	Min.	Max.
Prosthion-basion line	1	90.50	_	
Nasion-prosthion line	1	66.00	_	
Bizygomatic breadth	2	115.50	110.0?	121.0
Nasal height	2	45.25	42.0?	48.5
Nasal breadth	2	23.75	23.5	24.0
Ant. inter-orbital breadth	2	18.50	17.0?	20.0
Orbital breadth (right)	2	37.25	35.0	3 9 .5
Orbital breadth (left)	2	37.5 0	35.5?	3 9 .5
Orbital height (right)	2	31.00	28.0	34.0
Orbital height (left)	2	31.50	29.0	34.0
Maxillo-alveolar length	1	55.00	_	
Maxillo-alveolar breadth	2	57.75	57.0	58.5
Palatal length	1	48.00	_	-
Palatal breadth	2	35.00	32.0	34.0
Superior facial index	1	54.55	_	
Orbital index (right)	2	83.04	80.00	86.08
Orbital index (left)	2	83.88	81.69?	86.08
Nasal index*	1	49.48	-	
Maxillo-alveolar index	1	106.36	_	
Palatal index	1	79.17	_	

TABLE 64

ADULT FEMALE TYPE A

Mean Indices of the Whole Skull

Characters	n	Mean ± S.E.	Min.	Max.
Tr. cranio-facial index	1 1	96.80 53.66		_
Vert. cranio-facial index Long. cranio-facial index Jugo-frontal ind ^{ex}	1 2	52. 01 76.2 8	74.38	78.18?

^{*} Due to defective nose, slightly exaggerated value of masal index of Skl. No. I S 13 has been excluded from consideration.

TABLE 65 ADULT FEMALE TYPE A Mean Measurements and Indices of Permanent Maxillary Molar Teeth

				and the latest terms and
Characters	n	Mean	Min.	Max.
Mesiodistal crown diam. of M_1	2	9.62	9.25	10.00
Labiolingual crown diam, of M _I	2	11.00	11.00	11.00
Mesiodistal crown diam. of M2	2	9.50	9.00	10.00
Labiolingual crown diam. of M2	2	11.25	11.00	11.50
Mesiodistal crow diam. of M3	1	8.25		_
Labiolingual crown diam. of M3	1	12.00		
Crown index of M ₁	2	115.06	111.11	119.00
Crown index of M2	2	119.64	111.11	128.17
Crown index of M3	1	145.40		

TABLE 66 ADULT FEMALE TYPE B Mean Measurements and Indices of the Neurocranium

Characters	n	Mean
Max. cranial length	1	175.00
Max. cranial breadth	1	135.00
Nasion-inion length	1	160.00
Basion-bregma height	1	126.00
Min. frontal breadth	1	92.00
Vertical porion height	1	110.00
Median sagittal arc	1	372.00
Vertical transversal arc	1	292.00
Horizontal circumference	1	490.00
Cranial module	1	145.33
Calculated cranial capacity	1	1270.89
Length-breadth index	1	77.14
Length-height index	1	72.00
Breadth-height index	1	93.33
Length-auricular height index	1	62.86
Breadth-auricular height index	1	81.48
Fr. from parietal index	1	68.15

Characters	n	Mean
Prosthion-basion line	1	93.00
Nasion-prosthion line	1	67.00
Bizygomatic breadth	1 .	118.50
Nasal height	1	47.00
Nasal breadth	1	23.00
Ant. inter-orbital breadth	1	15.00
Orbital breadth (right)	1	42.00
Orbital breadth (left)	1	40.50
Orbital height (right)	1	32.00
Orbital height (left)	1	32.00
Maxillo-alveolar length	1	58.00
Maxillo-alveolar breadth	1	63.50
Palatal length	1	49.00
Palatal breadth	1	40.00
Superior facial index	1	56.54
Orbital index (right)	1	76.19
Orbital index (left)	1	79.01
Nasal index	1	48.94
Maxillo-alveolar index	1	109.48
Palatal index	1	81.63

TABLE 68

ADULT FEMALE TYPE B.

Mean Indices of the Whole Skull

Characters	n	Mean
Tr. cranio-facial index	1	87.78
Vert. cranio-facial index	1	53.17
Long. cranio-facial index	1	53.14
Jugo-frontal index	1	77.64

TABLE 69 ADULT FEMALE TYPE B Mean Measurements and Indices of the Permanent Maxillary Molar Teeth

Characters	n	Mean
Mesiodistal crown diam. of M ₁	1	11.00
Labiolingual crown diam. of M_1	1	11.75
Mesiodistal crown diam. of M_{2}	1	9.75
Labiolingual crown diam. of M_2	1	11.50
Mesiodistal crown diam. of M_3	1	10.00
Labiolingual crown diam. of M_3	1	11.00
Crown index of M ₁	1	106.82
Crown index of M2	1	118.16
Crown index of M ₃	1	110.00

TABLE 70 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean ±	S. E.	S. D.	c.v.	Min.	Me ,
Max. cranial length	2	174.50				174.0	175.0
Max. cranial breadth	2	130.00				125.0	135.0
Nasion-inion length	2	158.50		_		157.0	160.0
Basion-bregma height	2	124.50			-	123.0	126.0
Min. frontal breadth	3	89 .33 ±	1.77	3.06	3.43	86.0	92.0
Vertical porion height	3	108.00 ±	1.53	2.65	2.45	105.0	110.0
Median sagittal arc	2	363.00		_		354.0	372.0
Vertical transversal arc	2	284.50				277.0	292.0
Horizontal circumference	2	488.00			******	486.0	490.0
Cranial module	2	143.00			**************************************	140.67	145.33
Calculated cranial capacity	2	1205.37			-	1139.85	1270.89
Length-breadth index	2	74.49			-	71.84	77.14
Length-height index	2	71.34		Performance	Professional	70.69	72.00
	2	95.86			***************************************	93.33	98.40
Breadth-height index	2	61:60				60.34	62.86
Length-auricular height index	2	82.74		******		81.48	84.00
Breadth-auricular height index Tr. fronto-parietal index	2	70.08				68.15	72.00

Table 71 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Splanchnocranium


Characters	n	Mean +	S. E.	S. D.	c. v.	Min.	Max.
		mean <u>_</u>	0				
Prosthion-basion line	2	91.75		_		90.5	93.0
Nasion-prosthion line	2	66.50		_		66.0	67.0
Bizygomatic breadth	3	116.50 ±	3.34	5.77	4.95	110.0?	121.0
Nasal height	3	45.83 ±	1.97	3.40	7.42	42.0?	48.5
Nasal breadth	3	$23.50 \pm$	0.29	0.50	2.13	23.0	24.0
Ant. inter-orbital breadth	3	$17.33 \pm$	1.46	2.52	14.54	.15.0	20.0
Orbital breadth (right)	3	38.83 ±	2.05	3.55	9.14	35.0	42.0
Orbital breadth (left)	3	$38.50 \pm$	1.53	2.65	6.88	35.5?	40.5
Orbital height (right)	3	$31.33 \pm$	1.77	3.06	9.77	28.0	34.0
Orbital height (left)	3	$31.67 \pm$	1.46	2.52	7.96	29.0	34.0
Maxillo-alveolar length	2	56.50		-	-,	55.0	58.0
Maxillo-alveolar breadth	3	$59.67 \pm$	1.97	3.40	5.70	57.0	63.5
Palatal length	2	48.50				48.0	49.0
Palatal breadth	3	$36.67 \pm$	2.40	4.16	11.34	32.0	40.0
Superior facial index	2	55.54			-	54.55	56.54
Orbital index (right)	3	$80.76 \pm$	2.88	4.99	6.18	76.19	86.08
Orbital index (left)	3	$82.26 \pm$	2.06	3.57	4.34	79.01	86.08
Nasal index	3	$51.46 \pm$	2.25	3.90	7.58	48.94	55.95?
Maxillo-alveolar index	2	107.92		-		106.36	109.48
Palatal index	2	80.40				79.17	81.63

TABLE 72 ADULT FEMALE COMBINED Statistical Constants of the Indices of Whole Skull


Characters	n	Mean \pm S.E.	S. D.	C.V.	Min.	Max.
Tr. cranio-facial index Vert. cranio-facial index Long. cranio-facial index	2 2 2	92.29 53.42 52.58		* Andreaded	8 7.7 8 53.17 52.01	96.80 53.66 53.14
Jugo-frontal index	3	76.73 ± 1.19	2.06	2.68	74.38	78.18?

Table 73 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Permanent Maxillary Molar Teeth

The second of th								
Characters	r	n	Mean <u>+</u>	S.E.	S.D.	c. V.	Min.	Max
And the Market and the Control of th								
Mesiodistal crown diam. of M_{1}	;	3	$10.08 \pm$	0.51	0.88	8.73	9.25	11.00
Labiolingual crown diam, of M_1		3	$11.25 \ \pm$	0.24	0.42	3.73	11.00	11.75
Mesiodistal crown diam. of M_2		3	$9.58 \pm$	0.30	0.52	5.43	9.00	10.00
Labiolingual crown diam. of M_2		3	$11.33 \pm$	0.16	0.28	2.47	11.00	11.50
Mesiodistal crown diam. of M_3	2	2	9.12				8.25	10.00
Labiolingual crown diam. of M_3	2	2	11.50		_		11.00	12.00
Crown index of M_1	*	3	$112.31 ~\pm$	3.57	6.18	5.50	106.82	119.00
Crown index of M_2	· (3	$119.15~\pm$	4.95	8.57	7.19	111.11	128.17
Crown index of M_3	2	2	127.70				110.00	145.40



Norma Frontalis



Norma Lateralis

AREA (MOUND) AB

Mound AB, F and Area J have yielded altogether 25 skeletons mostly represented by incomplete and stray bones. The skeletons comprise of 17 adults, 5 children and 2 infants below 3 years of age. As regards the burials of Mound AB and F, Guha (Vats 1940: 162) doubts their authenticity as real Harappan and regards as modern graves, on the ground that one skull possessess higher specific gravity as compared to bones of Cemetery H and Area G. Vats (1940: 162), however, refutes Guha on archaeological evidence and insists that Mound burials are genuinely of Harappan culture.

Out of the adult skeletons represented in the Mound Area, the sex of the 12 adults has been ascertained which shows equal number of males and females (6 males and 6 females).

In order to evaluate the physical types of the Mound Area, only three adult crania from Mound AB comprising one male and two females could be utilized, while the other adults are too fragmentary to be of use in this respect. The measurements of one child's cranium is recorded in Collective Table C, but the determination of its physical type has at present not been attempted.

CRANIA

The measurements and observations of the skulls are incorporated in Collective Table C and I. Detailed descriptive notes on the skulls are given below:

The skull belongs to an adult male between 25 and 30 years of age at death. It is complete with mandible and in an excellent state of preservation. Supraorbital ridges are heavy amounting almost to a torus orbitalis. Mastoid processes are large and the muscular impressions at the nuchal region are well marked. Basilar suture is united while coronal, sagittal and lambdoidal sutures are open. Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides. The skull is dolichocranic (L-B Index 73.08), high Cranial contour in norma verticalis is byrsoides.




Fig. 80

Mandible is moderately developed with central triangular chin eminence and thick short symphysis (27.5 mm). Corpus is moderate in bulk and ramus moderate in width with high coronoid process. The muscular attachements for the massesteric and *pterygoidei interni* are sharply defined. The lower jaw is wide at its angle, the bigonial diameter being 104 mm.

SKL. 5440 D

Between the IIIrd and IVth strata two fragmentary human skulls (Skl. 5440 B and Skl. 5440 C), one lower jaw (No. 5440 A) and some other stray bones were brought to light but no mention was made of the fragmentary human Skl. 5440 D in Vat's Report. One skull bearing number 5440 D has, however, been entered in the field catalogue and available in the collection at our disposal while there is no skull numbering 5440 B.

The skull (No. 5440 D) belongs to an adult female, and appears to be in slightly mineralized condition. The skull is without face and mandible. Mastoid processes are thin and slightly long. The skull is hyperdolichocranic (L-B Index 69.95?). Only few measurements could be taken due to the incompleteness of the skull.

Brain cast*: Left half of original brain, transformed into earth retaining its natural shape size and morphology, weighing 505.4 gms (including the shellac coat applied for preservation). The left cerebral hemisphere comprises frontal, temporal, parietal and occipital lobes on superolateral aspect of surface. Medial aspect shows smooth impression of falx crebrii. Below this is the hollowed impression of callosal sulcus, inferior of which the continuity of corpus callosum joining the two cerebral hemispheres is cut off. On the base deep hollow of colateral sulcus is noticed. Gyrii smoothed

^{*} Thomks are due to Dr 6. S. Medy for his observation on




Fig. 81 Skl. 11635





Fig. 82 Skl. Mound AB (Rec.)

MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

out except at the base of the frontal lobe. Among sulcii fissures of Sylvius and Rolando marked by deep surface. Similar replica of left cerebral hemisphere is artificially and inaccurately joined at the superomedial border with the corresponding right half (Pl. LXXVIII: 3).

SKL. Mound AB (Rec.) [Figs. 82, 83, 84]

The skull is that of an adult female. Age at death was probably 25 or under. It is comparatively smaller in size and the contours are gracile in nature. The supraorbital ridges are absent and the glabella is only slightly marked. Mastoid processes are moderately small and the muscular impressions at the nuchal region are ill developed. Some abrasion marks are noticed on the forehead. Forehead is very slightly receding and passes upwards and backwards in a uniform high curve. Frontal and parietal eminences are slightly marked. The skull is hyperdolichocranic (L-B Index 69.83) and orthocranic (L-AH Index 60.89). Nose being high pitched, long and narrow is leptorrhine (NI 45.83). The nose is not depressed at the root. The maxillary teeth are represented by a well set of 16 intact teeth of which only the left M3 is carious.







Fig. 84

The lower jaw is well preserved. The ascending ramus in moderately high (60 mm) and wide (35 mm), having shallow wide sigmoid notch and low blunt coronoid process. Chin is bilateral and protruding. The gonial angles are turned slightly inwards, the bigoniac diameter being 84 mm. The depth at the mandibular symphysis is 27 mm. Mandibular teeth are represented by 15 intact teeth and the left central incisor is missing. Both left and right Ms show signs of caries.

ANALYSIS


Three adult skulls comprising one male and two females from Mound Area have been taken into account for the purpose of present study. It is noteworthy that the adult male found at Mound AB stands apart from the other skulls of Harappa being exceptional in non metric and morphometric features. In norma frontalis the adult male skull Skl. 11635 shows pronounced and almost continuous



Frontal view



Vertical view



Left lateral view



Occipital view

SKL. H 11635 : MOUND AREA

browridges, receding forehead, chamaerrhine nose sunken at the root with no subnasal prognathism. The other cardinal features are (i) high vault of the skull rising to an apex at the crown, (ii) reduced post-porionic length and (iii) flattened occiput (abgehackt), which undoubtedly conform its affinity with Armenoid strain, hitherto unknown at Harappa.

Of the two adult female skulls represented in this area, one [Skl. Mound AB (Rec.)] is excellently preserved and complete, and the other (Skl. 5440 D) is incomplete. The complete one is long-headed, gracile and smooth-contoured having medium vault, smooth forehead, narrow prominent nose, protruding occiput and ill-developed muscular attachements. Although this female possesses a slightly lower value of cranial index (69.83), judging the essential cranio-facial features it could be well assigned to a type which have been designated as Type A₁. The other female skull (Skl. 5440 D) could not be definitely diagnosed due to its incompleteness; but it is enough to say that this one is also a gracile and smooth-contoured long-headed skull similar to Type A₁.

CEMETERY H STRATUM II (OPEN BURIALS)

Area (Cemetery) H lay south of Mounds AB and J, comprised two distinct strata, namely stratum I and stratum II, which yielded human skeletons. were recovered. The human remains found at Cemetery H st II (open burials) are fragile and have suffered a considerable damage; but the amount of disintegration that has taken place in the contents of the burial jars Cemetery H st I (jar burials) is mostly due to the pressure of the superincumbent earth and percolation of rain water into the jars.

The human skeletal remains found in Cemetery H st II (open burials) represent 26 individuals, 11 of whom are adult males, 8 adult females, a female youth aged about 14 years, two children aged between 6 to 12 years and four adults whose sex could not be determined. Of the total number of 19 adult skulls found, 13 adults were considered for the present study. The rest are either crushed or so distorted that no reliable observation about their shape and dimension can be done. In appearance, the bones are of pale fawn colour, not unlike those found at Nal and Mohenjo-daro. The sex and age distribution of the adult skulls studied are tabulated below. It appears from the following table that the percentages of aged individuals are strikingly low among the people from Cemetery H st II (open burials).

TABLE 74
SEXWISE AGE DISTRIBUTION

SEX	LESS THAN 25 YEARS	25-30 YEARS 30-40 YEARS	40-50 YEARS
MALE		H 184 (K) H 484 (a) H 487 (a) H 502 (G) H 695	H 307 (а) Н 698
FEMALE	H 306 (a) H 699	H 88 H 488 H 501 (a)	Н 710

A. CRANIA

SKL. H 307 (a) [Fig. 85]

The sex is undoubtedly male, aged about 40-50 years. Damaged in the left temporal and facial parts. The skull is rather large and thick-walled. Coronal and sagittal sutures are partly obliterated and the lambdoid is open. Both mastoids are large and muscle attachements well developed. Glabella is prominent and supraorbital ridges moderately developed. The forehead is broad (99.5 mm): left orbit is of medium-height, rectangular, slightly oblique and right orbit falls into a low mesoconchal division of index (75.56). Norma verticalis is pentagonoid with prominent parietal bosses. When viewed from the side the skull appears to be well arched with a gradual and uniform curve. The skull is mesocranic (L-B Index 75.13), orthogranic (L-H Index 70.37) and metriogranic (B-H Index 93.66). In norma occipitalis the cranial contour represents the so called 'house' shape. The estimated cranial capacity is 1474.07 cc.

SKL. H 487 (a)

Adult male, aged about 25-30 years. The right and left parietal surfaces are partially absent, leaving only the central portion of the cranial vault. The greater part of the forehead and the right half of the face including the nose and the palate are intact but the outer rim of the left orbit and part of the left maxilla are missing. The supraorbital ridges are not prominent and muscular attachments are moderately developed. As viewed in norma lateralis the skull appears very long (cranial length 191.0 mm). The forehead is slightly inclined and goes back in an even curve, and the occiput is fairly prominent. The left orbit is squarish and mesoconchal in form (left OI 79.49), inclined downwards and laterally. The nasal aperture is narrow and leptorrhine (NI 45.45). The palate is broad and parabolic in shape. All the teeth have erupted and the molars moderately abraded (dentine exposed on cusps).

SKL. H 502 (G)

The skull is incomplete and appears to be that of an adult male. The entire skull vault is missing and right side of the face is considerably distorted. The cranial contour is ovoides in shape and the occiput fairly rounded. The glabella is prominent, with narrow concave nose and relatively broad midfacial region. The face is of moderate length: from nasion to gnathion it measures 116 mm, the nasion-prosthion length is 66 mm.

The lower jaw is almost complete (with all the teeth intact), except for the coronoid and condyloid processes of the right ascending ramus. The teeth are moderately worn down. The lower molars on both sides have four cusps each.

SKL. H 695 [PL LXIII]

Adult male, aged about 30 years. Damaged on the left parietal side, right temporal region and part of the face. The colour of the skull is greyish brown, the head is large with thick-walled cranial bones. The right masteid is broken, the left quite large. Supraorbital ridges are medially strong but






Fig. 85 Skl. H 307 a)



MID-SACITTAL CURVE (1/2 NATURAL SIZE)



Norma Frontalis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 698 (Natural Size)



Norma Lateralis

laterally absent. It is a high skull, the basi-bregmatic height being 137 mm. The forehead is of medium height and slightly receding. The vertex is slightly arched and passes into a non-protuberent occiput. Orbits are squarish and the nose is rather wide, nasal index being 51.38? (chamaerrhine). Norma verticalis is pentagonoid in shape, narrowed in front and expanded at the parietals. The skull is subbrachycranic (L-B Index 79.27?) and orthocranic (L-H Index 70.98). The palate is broad, deep and upsiloid in shape. The calculated capacity of the cranium gives aristencephalic value (1600.70 cc?).

SKL. H 698 [Fig. 87; Pls. LIII, LIV, LXIV: 1-2]

The skull is complete with mandible. There is a long horizontal depressed fracture in the right parietal (plate LXXVIII fig. 2), probably produced by some metallic edged weapon. The sex is undoubtedly male; the prominent glabello-superciliary region, big right mastoid process and well-developed muscular attachments are masculine features. None of the cranial sutures is closed, but the basilar suture is united. In norma verticalis, the theca cranii is ovoides in shape. In norma temporalis the forehead is inclined and passes back into a convex vertex which slopes evenly down to a rounded occiput. The skull is mesocranic (L-B Index 75.75), hypsicranic (L-AH Index 64.85) and lies in the border between metriocranic and acrocranic (B-AH Index 85.61). Norma facialis shows fairly medium face (the total facial index being mesoprosopic 85.66) with quadratic orbits. The nose is leptorrhine with an index of 47.62. All the maxillary teeth have erupted and cusps are moderately abraded.




Fig. 87 Skl. H 698

Mandible is well preserved with all the teeth in their sockets. The corpus is moderately large and ramus broad. The mylohyoid ridge of the mandible is pronounced, as are the pterygoid attachments. Chin is bilateral and projecting.


SKL. H 88

Adult female, aged about 30 years. The skull is incomplete. Whole of the basis cranii, greater part of the nuchal plane of the occiput, lateral margin of the right orbit, os sphenoidale and left os temporale are missing. In norma verticalis the skull is somewhat ellipsoidal, the parietal eminences not being very prominent. The maximum length of the cranium is 194 mm. In norma temporalis the forehead rises with a somewhat backward inclination and passes into relatively low arched vertex which slopes down obliquely into a fairly prominent tuber occipitale. The glabello-superciliary region is not prominent and nasion is not sunken at the root. The nasal bone is short but prominent with the ridge faintly concave in outline. In norma facialis the face is of medium height, nose fairly narrow- and left orbit hypsiconchal in form (OI 94.67). Palate is moderately deep and parabolic in shape. All the maxillary and mandibular teeth have erupted and appear to be free of caries.

The mandible is small and gracile in its general appearence. The muscular markings in the diagastric, massesteric and pterygoidal insertional areas are only mildly developed. Chin is prominent and the mandibular angle is straight and not everted. The lower dental arcade is *upsiloid* in shape and teeth show considerable signs of attrition.

SKL. H 488 [Figs. 86, 88]

Probably adult female, about 30 years. The skull is incomplete, greater portion of the left temporal bone and small part of the left occipital bone are missing. Mastoids are medium in size. Supraorbital ridges are laterally absent but present moderately medially. The face is rather low and medium in breadth. Orbits are low, rectangular and chamaeconchal in form (right OI 73.81 and



left OI 75.00). The nasal aperture is fairly wide and short, and the index falls within chamerrhine class. Norma verticalis is long ovoid with slight parietal eminences. In norma temporalis the nasion is not deep, the frontal bone is slightly inclined and passes smoothly into the slightly domed vertex with a pre-lambdoid flatening. Occiput is house shaped. It is of interest to note the presence of os inca.

SKL H 699 [Fig. 89]

The skull is of an adult female in the prime of life. The glabella is not prominent, mastoids are small and superior margin of the orbits are sharp. The lower jaw was found articulated in the skull. The upper part of the forehead extending up to the bregma was chopped off presumably




Fig. 89

accidentally during excavation. From the top, the skull is ovoid without parietal eminences. The cranium is brachycranic (L-B Index 80.24). The nasal ridge is short, slightly concave and nasion is not depressed at the root. The nasal aperture is short and of medium breadth with an index of 52.87 (chamaerrhine). There is also evidence of slight mental protuberance.

The mandibular body length is 70.0 and bigoniac diameter is 79.0~mm.

SKL H 710

Adult female, about 45 years of age. The skull is incomplete, lacks the entire facial region and a large portion of the vertex. Size is rather small, supraorbital ridges are ill developed and the




Fig. 90

mastoids small. Viewed from the top, the skull is an elongated oval. From the side, the forehead is vertical with a gradual bend at the metopian. The occipital line is marked by a prominent occipital protuberance. The mandibular fossae are of medium depth.

ANALYSIS

There are 13 adult skulls belonging to Cemetery H st II (open burials). On the basis of morphometric data and cranioscopic observation (Collective Tables D, I & J) 5 of these have been typed as round-headed B2 type (four males, one female), 3 have been typed as long-headed gracile A1 type (one male, two females) while one long-headed female skull with medium supraorbital ridges, moderately deep subglabellar notch, low face and wide nasal aperture is attributed to Type A. The remaining four skulls [H 184 (K), H 306 (a), H 484 (a) & H 501 (a) | cannot be considered for type diagnosis. The sexwise distribution of the *typical* adult skulls are given below:


SEX	TYPE A	TYPE A	TYPE \mathbf{B}_2
MALE	-	H 487 (a)	Н 695 Н 502 (G)
Ø			H 307 (a) H 698
FEMALE	H 488	H 710 H 88	Н 699

The round-headed Type B_2 skulls are not identical with the round-headed skulls from Area G, called as Type B_1 . Three skulls, however, classified here as Type A_1 are comparable to Type A_1 skulls from Cemetery R 37 and Area G.

TYPE B.

In tables 85 to 90 are presented in detail the mean neurocranial and splanchnocranial measurements and indices for four males, and in tables 107 to 110 the neurocranial and splanchnocranial measurements and indices of a single female, classified as belonging to the round-headed Type B_2 . The cardinal cranio-facial proportions of the Type B_2 male and female from Cemetery H st II (open burials) are as follows:

	MALE	FEMALE
Length-breadth imdex	Mesocranic (76.72 ± 1.16)	8) Brachycranic (80.24)
Length-height index	Orthocranic (71.82 ± 1.10)	6) Hypsicranic (80.84)
Breadth-height index	Metriocranic (93.68 \pm 2.40	O) Acrocranic (100.75)
Length-auricular height index	Hypsicranic (64.85)	Hypsicranic (65.27)
Total facial index	Mesoprosopic (85.66)	Leptoprosopic (92.17)
Superior facial index	Mesen (50.00)	Mesen (50.43)
Orbital index	Mesoconchic (83.74 ± 3.0)	1) Hypsiconchic (91.57)
Nasal index	Mesorrhine (49.50)	Chamaerrhine (52.87)
Palatal index	Leptostaphylin (77.12)	_
Trans. fronto-parietal index	Metriometopic (67.13 \pm 3.33	5) Metriometopic (67.16)



Norma Frontalis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 699 (Natural Size)



Norma Lateralis

DIOPTOGRAPH TRACING (WASH): MALE SKL. H 699 (Natural Size)

Disparity in cranio-facial proportions in the male and the female B_2 type is distinct. But discussion on this point, on the basis of a single female cranium, appears not justifiable. The round-headed male B_2 type skulls have obtained their cranial form mainly by increased breadth rather than decreased length. The average cranial length, however, is nearly same as that of the basic long-headed type of Cemetery R 37.

In Plate LVII and in Figures 91-93 photographs and dioptographic drawings of a round-headed Type B_2 male skull are presented. In the front view the face is broad and of medium height, the nasal aperture is of average height and width to give a mesorrhine index. In the lateral view the forehead is high and little inclined, the vertex appears to be well arched and the occiput is curved, rather than flattened. From the top, the cranial contour is *ovoides* in shape with marked parietal bosses.

TYPE A

The important cranio-facial characters of the single female Type A skull from Cemetery H st II (open burials) is as follows:

Dolichocranic	(71.93)
Orthocranic	(73.02)
Acrocranic	(101.52)
Orthocranic	(59.95)
Metriocranic	(83.33)
Chamaeconchic	(74.40)
Hyperchamaerrhine	(66.67)
Leptostaphyline	(78.41)
Eurymetopic	(71.21)
	Orthocranic Acrocranic Orthocranic Metriocranic Chamaeconchic Hyperchamaerrhine Leptostaphyline

The cranio-facial proportions of the Type A female skull of the open burials are closely similar to the female Type A crania from Cemetery R 37. The discrepancy with regard to masal and orbital index is apparently fortuitious or may be individual variation.

In Plate LVIII and Figures 94-96 are presented photographs and dioptographic drawings of Type A female skull from stratum II (open burials). The face is rather low, medium in breadth and supraorbital ridges are present moderately medially. The vertical contour is seen to be long oviod.

TYPE A

In tables 102 to 106 the average cranil and facial measurements and indices of open burials Type A_1 are presented in detail for two females and in tables 81 to 84 for one male. In view of the fact all the three skulls representing Type A_1 are broken, typology was done primarily on the basis of morphological observations. The skulls are on the whole gracile; the supraorbital ridges are not prominent and muscular impressions weak. In lateral view the skulls are very long, the forehead slightly inclined and the occiput fairly prominent with a *chignon*.

DISCUSSION

Cemetery H st II (open burials) is represented by three cranial types, namely, Type A, Type Aı and Type B_2 . Type A and Type A_1 are both long-headed and they appear to be similar to the people representing Cemetery R 37. There is no hesitation in identifying the solitary female skull No. H 488with the dolichocephalic Type A crania, already noted in Cemetery R 37. There is also appreciable resemblance of Type A_{I} crania of Cemetery H open burials with those of Cemetery R 37 and Area G, both in morphometric and in non-metric characters. On the other hand, round-headed $B_{\scriptscriptstyle 2}$ type from open burial differs from the round-headed $B_{\scriptscriptstyle \rm I}$ type of Area G in being larger in size and having more retreating frontal bones. The cranial capacity of the Type B_2 skulls is fairly large and the absolute value for the capacity differs markedly from the earlier B_1 type skulls. B_2 type is characterized by large cranial capacity (1505.19 cc.) and mesocephalic head. The face is medium in breadth with an upper face index of 50 (males). The vertex is well arched with curvo-occiput and the skull is hypsicephalic. However, in cranio-facial dimensions the Type B1 skulls from Area G are distinctly shorter when compared with Type B_2 skulls. The most marked difference is found between the calvarial lengths. The average cranial length of the Type \mathbf{B}_2 male is 188.62 mm, while the mean cranial length for Type B_{I} male is 177.00 mm. In Type B_{I} the cranial vault seems to be shorter and lower than in the Type B_2 ; appreciable difference between the two types is also noticed in the facial region of the skull.

In the following table the average value for some important cranial and facial dimensions as well as indices of Type B_1 and Type B_2 have been compared with the round-headed Alpine male from Hissar III.

	: Maximum Length	M aximum breath	Basion Bregma height	Upper face height	Dizygomatic breath	Cephalic index	Length height index
Round-headed Alpine Hissar III	177.4	136.2	134.4	66.8	129.6	76.77	75.77
Round-headed Type B ₁ Area G	177.0	140.8	132.6	64.7	129.5	79.54	75.01
Round-headed Type B ₂ Cemetery (open burials)	H 188.6	144.7	135.3	69.7	136.0	76.72	71.82

It is at this point again worth mentioning that the Type B₁ skulls from Area G are closely similar in major calvarial diameters, in facial height and width to the round-headed type of Hissar III designated by Krogman (1940: 22) as Alpine.

In cranial and facial dimensions Type B₂ skulls are distinctly larger than the Type B₁ and round-headed Alpine type of Hissar III, but these differences in absolute measurements are not reflected in the salient cranio-facial proportions, of which such characters are components. This round-headed B₂ type may be an Alpine variant or a mixed variety of the Alpine. This type is predominant at Cemetery H st II.

B. EXTREMITY BONES

Great handicap was felt for a definitive morphological study of the extremity bones due to incompleteness. Measurements which have, however, been taken on them, are assembled in Collective Tables M-N.

HUMERUS

Fourteen adult male humeri (seven of each side) and twelve adult (five right and seven left) P have been measured.

TABLE 75

Characters		M A	L E		F E M	A L E
	n	Mean	Range	n	Mean	Range
Maximum length	7	326.86	320.0 - 341.0	7	285.71	267.0 - 298.0
Breadth of proximal epiphysis	6	51.17	50.0 - 52.0	6	43.75	40.0 - 46.5
Breadth of distal epiphysis	6	62.83	5 9 .5 - 64 .5	6	56.25	54.5 - 58.0
Circumference of the shaft at the middle	8	65.88	62.0 - 69.0	9	61.06	55.0 - 65.0
Minimum circumference of diaphysis	9	5 9 .33	55. 0 - 6 5.5	10	53.95	49.0 - 56.0
Robusticity index	7	17.95	16.72- 19.00	7	19.14	18.46- 20.22

The humeri of Skl. H 502 (G) rt & lt, H 695 (rt) and H 698 (lt), all of which belong to adult male, are well developed and sturdy built. Mean values pertaining to some of the important measurements and indices of humeri are produced in Table 75. The maximum length of humerus lies within a total range of 267 mm to 341 mm, irrespective of sex, which is nearer to the lower limit of the world range of 260-380 mm (Martin 1928: 1100). Males exceed the females in mean length measurement; the average for males is 326.86 mm and for females 285.71 mm.

Contrary to general tendency of different human groups (Martin 1928: 1134) in robusticity index the male average (17.95) falls short of the female average (19.14), the index for the males ranging between 16.72-19.00 and for the females between 18.46-20.22.

The perforation of fossae electrani was noticed in five cases, viz., Skl. H 488 (lt), H 502(G) (rt), H 694 (lt), H 696 (rt) and H 699 (lt). Mean length of male humeri of Cemetery H St II (open burials) falls much short of Cemetery R 37. In circumference also the present series gives a lower value than that of Cemetery R 37.




Fig. 91 Norma Frontalis




Fig. 92 Norma Verticalis





Fig. 93 Norma Lateralis


Figs. 91, 92 & 93. Dioptographic contours of Type B2 Male (Skl. H 698) of Cemetery H St. II (Open burials)



Norma Frontalis





Norma Verticalis



Norma Lateralis


Photographs of Type B_2 Male [Skl. H 698] of Cemetery H St. II (Open burials)







Norma Verticalis



Norma Lateralis

Photographs of Type A Female (Skl. H 488) of Cemetery H St. II (Open burials)




Fig. 94 Norma Frontalis




Fig. 95 Norma Verticalis




Fig. 96 Norma Lateralis

Figs. 94, 95 & 96. Dioptographic contours of Type A Female (Skl. H 488) of Cemetery H St. II (Open burials)

RADIUS

Only bones of three males (two right and one left) and two females (one right and two left) could be measured, the rest being unsuitable for metrical use. Some of the important mean values are shown in Table 76.

TABLE 76

Character		M A	L E		F E M	ALE	
	n	Mean	Range	n	Mean	Range	
Maximum length	2	253.50	251.0 - 256.0	3	225.00	221.0 - 233.0	
Physiological length	3	243.17	232.0 - 257.0	3	210.83	206.0 - 218.0	
Minimum circumference of diaphysis	, 3	41.33	41.0 - 42.0	3	36.67	35.0 - 38.0	
Robusticity index	. 3	17.03	15.95- 17.67	3	17.42	16.06- 18.2	

The mean maximum length is 253.5 mm for males and 225 mm for females. In robusticity index females (17.42) appear to have slightly higher average than males (17.03). The difference between the maximum length and the physiological length is 10.33 mm for males and 14.17 mm for females.

ULNA

Five bones (two right and three left) three males and seven bones (three right and four left) of four females have been considered useful for measurement. In Table 77, their mean values are given.

TABLE 77

							× 10 00 00 00 00
Character			M A	L E *		F E M	A L E
		n	Mean	Range	n	Mean	Range
							The control of the co
Maximum length		4	274.50	268.0 - 289.0	4	245.75	242.0 - 253.0
Physiological length		5	248.80	241.0 - 262.0	7	226.29	217.0 - 238.0
Minimum circumference of	diaphysis	4	33.5 0	31.0 - 37.0	7	30.36	26.0 - 32.5
Robusticity index		4	13.47	12.86- 14.92?	7	13.45	10.92- 14.68

The averages of maximum length is 274.5 mm for males and 245.75 mm for females. In robustness both the sexes are relatively equal as revealed from their robusticity index.

FEMUR

Four femora (two right and two left) of two male individuals and five (two right and three left) of three females have been considered. Their important mean metrical values are scheduled in Table 78.



Norma Frontalis




Norma Lateralis

Photographs of Type $A_{\rm I}$ Male [SkI, H 487(a)] of Cemetery H St. II (Open burtals)



Norma Frontalis



Norma Verticalis



Photographs of Type A_i Female [Skl. H 88] of Cemetery H St. II (Open burials)

87.27- 117.39

71.43- 85.71

5

5

83 64- 109 26

71.43- 90.00

101.73

78.39

Pilastric index

Platymeric index

Table 78

		Femur				
Character		M A	L E		F E M	A L E
	n	Mean	Range	e n	Mean	Range
Maximum length	_	_		- 3	389.00	378.0 - 410.0
Length in natural position		_		_ 3	388.33	378.0 - 408.5
Sagittal diameter at middle of diaphysis	4	24.12	22.0 - 2	27.0 5	22.10	20.0 - 24.0
Transversal diameter at middle of diaphysis	4	27.25	26.0 - 2	28.5 5	25.50	23.0 - 27.5
Circumference of the shaft at the middle	4	87.75	83.0 - 9	95.0 5	80.40	80.0 - 81.0
Length-thickness index		·	- n	— 3	20.71	19.83- 21.16
Robusticity index	_	_	-	_ 3	13.28	12.61- 13.76

Due to absence of intact male femora, maximum length and length in natural position could not be taken.

99 46

77.05

4

In females however the difference between these two characters is not significant. Robusticity index which expresses the relation of the dimensions at the diaphysis middle and the length in the natural position is available for the females only, the mean being 13.28. Pilastric index is slightly higher in the females. Converse result is shown in platymeric index.

The femora of both the male skeletons H 502(G) and H 698 are robust and strong built as revealed in their relief. Among the females the femur of Skl. H 710 appears to be slender and gracile than others.

It appears that females of the present series were relatively strong built than those of Cemetery R 37.


One conspicuous fact emerges from the table that in most of the indices females show higher value then males thereby indicating that females were comparatively robuster and stronger.

TIBIA


Only six tibiae (three right and three left) of five male individuals and five (four right and one left) have been utilised for consideration. Their important mean values are produced in Table

TABLE 79

		Tibia					
Character		W A	L E		1 1	e le m	A L E
	n	Mean	Rang	je	'n	Mean	Range
Maximum length (spino-malleolar) Physiological length	1 3	410.00 365.83	344.0 - 3		3 5 5	385.67 324.67	323.0 - 352.0? 303.0 - 332.0
Sagittal diameter at the middle Transversal diameter at the middle	6	34.92 22.83	20.0 -	39.5 25.0	5	28.80 18.90	26.5 - 30.0 17.0 - 20.5
Minimum circumference of diaphysis Index of cross-section in the middle Robusticity index	5 6	80.00 65.66	10.0	85:0 71.43	5 5 3	65.58 20.33	66.0 - 70.0 63.33- 68.33
	1	21.12			9	40.00	19:54- 21.07



Figs. 1-5. Anterior or Flexor aspect of the Humerus: Cemetery H St. II (1/2 Natural size)



Figs. 1-3 Radial aspect of Ulna; Figs. 4-7 Volar aspect of Radius : Cemetery H St. II ($\frac{1}{2}$ Natural size)

In mean physiological length both the sexes of the present series are shorter than that of Cemetery R 37. Index of cross-section at the middle, however, of the present series indicates higher values, irrespective of sex than that of Cemetery R 37.

FIBULA

Two fibulae (left) of two male individuals and two (one right and one left) of one female are utilized; some of the mean values have been presented in Table 80 below:

TABLE 80


Character		M A	L E	F	F E M	A L E
	n	Mean	Range	n	Mean	Range
Maximum length	1	369.00		1	323.00	
Minimum circumference of diaphysis	2	40.00	40.0 - 40.0	2	35.00	35.0 - 35.0
Robusticity index	1	10.84		1	10.84	

STATURE AND PROPORTION

It is intelligible from Collective Table ① and P that in most of the cases Dupertuis and Hadden's 'general formulae' yield highest value with regard to calculated stature, and Manouvrier's table shows intermediate results.

The average male stature is found to be 1717.57 mm for six males according to Dupertuis and Hadden, and 1702.37 mm for eight males according to Manouvrier's table. In the case of females it is 1563.92 mm for four individuals according to former and 1575.57 mm for five individuals according to latter.

The mean difference of stature between the males and females is 153.65 mm according to Dupertuis and Hadden's 'general formulae' and 146.66 by Manouvrier's table. These differences are, however, within normal range of sex difference.



Figs. 1-3 Anterior aspect of Tibia & Femur : Cemetery H St. II $(^{1}\!/_{\!2}$ Natural size).

It is not pertinent to predict stature of a population from insufficient data and it is all the more difficult to estimate stature of the sub-groups from even smaller data. However, the trend of the distribution of stature may be appreciated from the following table.

		STATURE ACCORDING TO	="
	MANOUVRIER	PEARSON	DUPERTUIS AND HADDEN
Type A (F)	1523.03 (1)	1504.1 (1)	1538.4 (1)
Type A_1 (F)	1602.74 (2)	1493.7 (1)	1533.6 (1)
Type B_2 (M)	1706.77 (3)	1660.36 (2)	1710.95 (2)

It appears that three male individuals represented in Type B_2 [Skl. H 502 (G), H 695 and H 698] tended to be tall. One female (Skl. H 710) of Type A_1 was perhaps medium or below medium in stature. Similar is the case of a female (Skl. H 488) of Type A.

Extremity bones of two males [Skl. H 184 (K) and H 502 (G)] and of three females [Skl. H 306 (a), H 488 and H 710] have been used for finding out limb proportions of individual skeletons.

In brachial index (humero-radial) a female skeleton (H 488) had remarkably longer radius in proportion to humerus (82.77%).

Three individuals [males: Skl. H 184 (K), H 502 (G) and female: H 306 (a)] were mesatikerkic, giving 79.75, 77.47 and 79.79? respectively as index value. One female (Skl. H 488) was highly delicho-kerkic (82.77).

Crural index (tibio-femoral) which indicates the relation of the leg with respect to the thigh could be worked out in only one female individual (Skl. H 488), gives a value of 85.22.

The humero-femoral index by which the relative lengths of humerus and femur can be compared is 70.63 and 68.05 for Skl. H 488 and H 710 respectively, both being females. The tibio-radial index, which expresses the relation of radial to tibial length, is available for one female skeleton (Skl. H 488) only which is 68.42.

70.68

STATISTIC	CAL	CONSTANT	:	CEMETERY	н	ST	11
(OPEN B	URIA	ALS)					

Length-height index

Table 81						
ADULT MALE TYPE	1	N				
Mean Measurements and Indices of	tne	Neurocranium				
Characters	n	Mean				
Max. cranial length	1	191.00				
Nasion-inion length	1	184.00				
Basion-bregma height	1	135.00				
Min. frontal breadth	1	92.50				
Median sagittal arc	1	383.00				

TABLE 82-Continued

Characters		n	Mean
Nasal index		1	45.45
Maxillo-alveolar	index	1	110.53
Palatal index		1	90.80

TABLE 83

ADULT MALE TYPE A

Mean Indices of the Whole Skull

Characters	n	Mean				
Carry of Eagle of the Eagle of the Carry	177 (A. 17)	CARACTE CONTRACTOR				
Vert. cranio-facial index	1	52.22				
Long. cranio-facial index	1	49.21				

TABLE 82

ADULT MALE TYPE A

Mean Measurements and Indices of the Splanchnocranium

Characters		n	Mean
Prosthion-basion line		1	94.00
Nasion-prosthion line		1	70.50
Nasal height		1	55.00
Nasal breadth		1	25.00
Ant. inter-orbital brea	dth	1	18.50
Orbital breadth (right)	1	39.00
Orbital height (right)		1	31.00
Maxillo-alveolar length	1	1	57.00
Maxillo-alveolar breadt	:h	1	63.00
Palatal length		1	43.50
Palatal breadth		1	39.50
Orbital index (right)		1	79.49

TABLE 84

ADULT MALE TYPE A,

Mean Measurements and Indices of the

Maxillary Molar Teeth

Mean Characters Mesiodistal crown diam. of M_1 1 10.00 1 Labiolingual crown diam. of M1 12.00 Mesiodistal crown diam. of M2 1 8.75 1 Labiolingual crown diam. of M_2 11.50 Mesiodistal crown diam. of M_{S} Labiolingual crown diam. of M_{S} 1 8.25 1 10.75 1 Crown index of M₁ 120.30 Crown index of M_2 1 128.60 1 Crown index of $M_{\rm 3}$ 131.11

Table 85 ${\rm ADULT\ MALE\ TYPE\ B}_2$ Mean Measurements and Indices of the Neurocranium

Characters	n	Mean	±	S.E.	Min.	Max.
						Va. 17
Max. cranial length	4	188.62	\pm	3.48	183.5	193.0
Max. cranial breadth	3	144.67	\pm	4.26	139.0	153.0?
Nasion-inion line	3	174.00	土	2.00	170.0	176.0
Basion-bregma height	3	135.33	\pm	1.20	133.0	137.0
Min. frontal breadth	3	96.83	\pm	2.19	92.5	99.5
Vertical porion height	1	119.00			· .	
Median sagittal arc	2	381.50			_	_
Vertical transversal arc	1	318.00			_	_
Horizontal circumference	3	530.33	土	8.22	514.00	540.00
Cranial module	3	156.17	土	2.37	152.83	161.00?
Calculated cranial capacity	3	1505.19	±	48.77	1440.81	1660.70?
Length-breadth index	3	76.72	\pm	1.18	75.13	79.27?
Length-height index	3	71.82	\pm	1.16	70.37	74.11
Breadth-height index	3	93.68	土	2.40	89.54?	97.84
Length-auricular height index	1	64.85			_	
Breadth-auricular height index	1	85.61			_	-
Tr. fronto-parietal index	3	67.13	土	3.35	60.46?	70.86

TABLE 86 ${\bf ADULT\ MALE\ TYPE\ B}_2$ Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	土	S.E.	Min.	Max.
Prosthion-basion line	2	94.50			94.0	95.0
Nasion-prosthion line	3	69.67	±	2.73	66.0	75.0
Nasion-gnathion line	1	116.5		20		and the same of th
Bizygomatic breadth	1	136.0				-
Nasal height	4	51.88	± -	1.52	47.5	54.5
Nasal breadth	3	24.67	\pm	2.03	21.0	28.0?
Ant. inter-orbital breadth	4	19.50	\pm	0.46	18.5	20.5
Orbital breadth (right)	3	43.00	±	1.16	41.0	45.0
Orbital breadth (left)	3	39.33	±	0.93	37.5	40.5
Orbital height (right)	3	34.00	\pm	0.58	33.0	35.0
Orbital height (left)	3	34.67	\pm	1.01	33.0	36 .5
Maxillo-alveoler length	3	54. 8 3	+	2.53	50.0	5 8 .5
Maxillo-alveolar breadth	3	67.50	+	3.89	62.0	75.0
Palatai length	2	46.75			- 46.0	47.5
Palatal breadth	3	40.50	±	3.76	36.5	48.0

TABLE 86-Continued

Characters	n	Mean <u>+</u>	S.E.	Min.	Max.
				_ , -	
Total facial index	1	85.66		_	-
Superior facial index	1	50.00		_	
Orbital index (right)	3	$79.15 \pm$	1.82	75.56	81.40
Orbital index (left)	3	88.34 ±	4.57	82.50	97.33
Nasal index	2	49.50*		47.62	51.38?
Maxillo-alveolar index	3	$123.30 \pm$	6.36	111.97	133.93
Palatal index	2	77.12		77.89	79.35

 $\begin{array}{c} \text{Table 87} \\ \text{adult male type } \mathbf{B}_2 \end{array}$

Mean Indices of the Whole Skull

Characters	n	Mean	土	S.E.	Min.	Max.
Tr. cranio-facial index	1	97.84			_	
Vert. cranio-facial index	2	52 .37			50.00	54.74
Long. cranio-facial index	3	51.12	\pm	1.07	49.22	52.91
Jugo-frontal index	1	72.43				

Characters	n	Mean	Min.	Max.
				2727.0
Bigonial breadth	2	79.00	75.5	82 .5
Bicondylar breadth	1	112.50?		×
Ht. of mandibular ramus	2	62.00	60.0	64.0
Max. breadth of mandibular ramus	2	39.75	39.5	40.0
Min. breadth of mandibular ramus	2	35.00	34.0	36.0
Ht. at mandibular symphysis	2	30.25	28 .5	32.0
Mandibular length	2	87.00	83.0	91.0
Mandibular angle	2	110.5°	110°	111°
Mandibular index	1	73.78?	All Control	Property
Breadth index of mandible	1	67.11?	and the second	No. was
Jugo-mandibular index	1	55.51	name.	Maria W

^{*} Measurements pertaining to nose of Skl. H 502 (G) are not taken into account due to distortion of nasal bones which are likely to vitiate average value.

TABLE 89 ADULT MALE TYPE B₂ Mean Measurements and Indices of the Permanent Maxillary Molar Teeth

And the latter of the probability of the						
Characters	n	Mean	\pm	S.E.	Min.	Max.
Mesiodistal crown diam. of M_1	3	10.50	\pm	0.38	10.00	11.25
Labiolingual crown diam. of M_1	3	11.17	土	0.30	10.75	11.75
Mesiodistal crown diam. of M2	3	9.33	\pm	0.51	8.50	10.25
Labiolingual crown diam. of M_2	3	10.33		0.21	10.00	10.75
Mesiodistal crown diam, of M_3	3	8.83		0.34	8.50	9.50
Labiolingual crown diam, of M_3	3	10.00	±	0.25	9.50	10.25
Crown index of M ₁	3	106.81		6.35	9 5.55	117.50
Crown index of M2	3	111.79	÷	8.22	98.08	126.47
Crown index of Ma	3	113.92		6.92	100.00	121.18

TABLE 90 ADULT MALE TYPE B. Mean Measurements and Indices of the Permanent Mandibular Molar Teeth

Characters	n	Mean	Min.	Max.
Mesiodistal crown diam. of M.	2	10.75	10.70	11.00
Labiolingual crown diam, of M.		10.75	10.50	
	2	10 .12	9.75	10.50
Mesiodistal crown diam. of M ₂	2	9.50	9.25	9.75
Labiolingual crown diam, of M2	2	9.00	9.00	9.00
Mesiodistal crown diam. of M ₃	2	10.25	10.00	10.50
Labiolingual crown diam, of M ₃	2	8.88	8.75	9.00
Crown index of M ₁	2	94.16	92.86	95.45
Crown index of M2	2	94.87	92.37	97.37
Crown index of Ms	2	86.82	86.14	87.50

TABLE 91 ADULT MALE COMBINED Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean ±	S.E.	S.D.	C. V.	Min.	Max.
Max. cranial length	5	189.10 ±	1.58	3.54	1.87	1 8 3.5	193.0
Max. cranial breadth	3	144.67 生	4.26	7.37	5.09	139.0	153. 0 ?
Nasion-inion length	4	176.50 ±	2.87	5.74	3.25	170.0	184.0
Basion-bregma height	5	13 4.80 ±	0.80	1.79	1.33	133.0	137.0
Min. frental breadth	4	95.75 ±	1.88	3.77	3.94	92.5	99.5

TABLE 91 - Continued

Characters	n	Mean \pm	S.E.	S. D.	c. v.	Min.	Max.
Vertical porion height	1	119.00			_		
Median sagittal arc	3	$382.00 ~\pm$	5.51	9.54	2.50	372.0	391.0
Vertical transversal arc	1	318.00			_	_	
Horizontal circumference	3	$530.33 \pm$	8.22	14.22	2.68	514.0	540.0
Cranial module	3	$156.17\ \pm$	2.48	4.29	2.75	152.83	161.00?
Calculated cranial capacity	3	$1505.19\ \pm$	48.77	84.37	5.61	1440.81	1660.70
Length-breadth index	3	$76.72 \ \pm$	1.29	2.23	2.91	75.13	79.27
Length-height index	4	$71.54 \pm$	0.86	1.73	2.42	70.37	74.11
Breadth-height index	3	$93.68 \pm$	2.40	4.15	4.43	89.54?	97.84
Length-auricular height index	1	⊳ 85.61¬			_	_	
Breadth-auricular height index	1	64.85€			_	_	
Tr. fronto-parietal index	3	$67.13 \pm$	3.35	5.79	8.63	60.46?	70.86

Characters	n '	Mean ±	S.E.	S. D.	c. v.	Min.	Max.
Prosthion-basion line	5	96.60 ±	1.40	3.13	3.24	94.0	100.0
Nasion-prosthion line	5	$70.30 \pm$	1.56	3.49	4.96	66.0	75.0
Nasion-gnathion line	2	116.25		-		116.0	116.5
Bizygomatic breadth	1	136.0			_	-	****
Nasal height	6	$52.67 \pm$	1.10	2.70	5.13	47.5	55.0
Nasal breadth	6	$25.50 \pm$	1.03	2.53	9.92	21.0	28.0?
Ant, inter-orbital breadth	6	$19.75 \pm$	0.56	1.37	6.94	18.5	22.0
Orbital breadth (right)	5	41.90 ±	1.00	2.25	5.37	39.0	45.0
Orbital breadth (left)	4	40.62 ±	1.45	2.90	7.14	37.5	44.5
Orbital height (right)	4	$33.25 \pm$	0.86	1.71	5.41	31.0	35.0
Orbital height (left)	4	$34.50 \pm$	0.74	1.47	4.26	33.0	36.5
Maxillo-alveolar length	6	55.08 ±	1.36	3.32	6.03	50.0	58.5
Maxillo-alveolar breadth	6	65.75 ±	2.02	4.96	7.54	62.0	75.0
Palatal length	5	46.80 ±	0.96	2.14	4.57	43.5	49.0
Palatal breadth	6	39.08 ±	2.14	5.25	13.43	32.5	48.0
Total facial index	1	85.66		-	* second	Statistics	
Superior facial index	1	50.00			a common		THE
Orbital index (right)	4	79.24 ±	1.28	2.57	3.24	75,56	81.40
Orbital index (left)	4	85.36 ±	4.40	8.79	10.30	76.40	97.33
Nasal index	5	47.64 ±	1.30	2.92	6.13	44.21	51.38?
Maxillo-alveolar index		119.53 ±	3.51	8.60	7.19	110.53	133.93
Palatal index	5	79.88 ±	3.78	8.46	10.59	67.71	90,80
	6 5				10.59	67.71	

Table 93 ADULT MALE COMBINED Statistical Constants of the Indices of the Whole Skull

The second secon							
Characters	n	Mean \pm	S.E.	S. D.	c. v.	Min.	Max.
Tr. cranio-facial index	1	97.84		_	_	_	_
Vert. cranio-facial index	3	$52.32 \pm$	1.37	2.37	4.53	50.00	54.74
Long. cranio-facial index	4	$50.64 \pm$	0.90	1.79	3.53	49.21	52.91
Jūgo-frontal index	1	72.43		_	_		-

Table 94 ADULT MALE COMBINED Statistical Constants of the Mandibular Measurements and Indices

Characters	~	n	Mean	Min.	Max.		
Bigonial breadth		2	79.00	75.5	82.5		
Bicondylar breadth		1	112.5?				
Ht. of mandibular ramus		2	62.00	60.0	64.0		
Max. breadth of mandibular ramus		2	39.75	39.5	40.0		
Min. breadth of mandibular ramus		2	35.00	34.0	36.0		
Ht. at mandibular symphysis		2	30.25	28.5	32.0		
Mandibular length		2	87.00	83.0	91.0		
Mandibular angle		2	110.5°	110°	111°		
Mandibular index		1	73.78?	_			
Breadth index of mandible		1	67.11?		-		
Jugo-mandibular index		1		AP			
		1	55.51				

Table 95 ADULT MALE COMBINED Statistical Constants of the Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean 🛨	S.E.	S.D.	c. v.	Min.	Max.
Mesiodistal crown diam. of M ₁	6	$10.12\ \pm$	0.27	0.65	6.42	9.25	11.25
Labiolingual crown diam. of M1	6	$11.21\ \pm$	0.22	0.54	4.82	10.75	12.00
Mesiodistal crown diam. of M2	6	$9.25 \pm$	0.25	0.61	6.59	8.50	10.25
Labiolingual crown diam. of M2	6	$10.54 \pm$	0.18	0.44	4.17	10.00	11.25
Mesiodistal crown diam, of M3	6	$8.75 \pm$	0.20	0.50	5.71	8.25	9.50
Labiolingual crown diam. of Ma	6-	10.00 🛳	0.20	0.50	5.00	9.50	10.75
Crown index of M	6	$111.16~\pm$	3.69	9.04	8.13	95.55	120.30
Crown index of M ₂	6	$114.74~\pm$	4.62	11.31	9.86	98.08	128.60
Crown index of Ma	6	$115.06\ \pm$	4.66	11.41	9.92	100.00	131.11

Characters		n	Mean ±	S.E.	S. D.	c. v.	Min.	Max.
Mesiodistal crown diam. of	M_1	3	$10.75 ~\pm$	0.14	0.24	2.23	10.50	11.00
Labiolingual crown diam. of	M ₁	3	$10.08\ \pm$	0.21	0.37	3.67	9.75	10.50
Mesiodistal crown diam. of	M_2	3	$9.92 ~\pm$	0.44	0.76	7.66	9.25	10.75
Labiolingual crown diam. o	f M ₂	3	$9.25 \pm$	0.24	0.42	4.54	9.00	9.75
Mesiodistal crown diam. of		4	$10.25\ \pm$	0.14	0.28	2.73	10.00	10.50
Labiolingual crown diam. of		4	8.94 ±	0.21	0.42	4.70	8.50	9.50
Crown index of M ₁	1113	3	93.76 ±	0.85	1.47	1.57	92.86	95.45
Crown index of M ₂		3	93.51 ±	1.98	3.43	3.67	90.80	97.37
Crown index of M ₃		4	87.28 ±	1.18	2.37	2.72	85.00	90.48

Table 9	7			TABLE 98 - Conti	nued	
ADULT FEMALE	TYPE A					
Mean Measurements and Indice	s of the	Neurocranium		Characters	n	Mean
Characters	n	Mean		Ant. inter-orbital breadth	1	18.00
				Orbital breadth (right)	1	42.00
Max. cranial length	1	183.50		Orbital breadth (left)	1	40.00
Max. cranial breadth	1	132.00?		Orbital height (right)	1	31.00
Nasion-inion length	1	167.00		Orbital height (left)	1	30.00
Basion-bregma height	1	134.00		Palatal length	1	44.00?
Min. frontal breadth	1	94.00		Palatal breadth	1	34.50
Vertical porion height	1	110.00		Orbital index (right)	1	73.81
Median sagittal arc	1	376.00		Orbital index (left)	1	75.00
Vertical transversal arc	1	297.00		Nasal index	1	66.67
Horizontal circumference	1	505.00?) ⁷	Palatal index	1	78.41?
Cranial module	1	149.83?				
Calculated cranial capacity	1	1298.21?	j:			
Length-breadth index	1	71.93?				
Length-height index	1	73.02	7	m 00		
Breadth-height index	1	101.52?		TABLE 99		
Length-auricular height index	1	59.95		ADULT FEMALE TYP		
Breadth-auricular height index	1	8 3.33?		Mean Measurements and Indices	of the IV	Iandible
Tr. fronto-parietal index	1	71.21?				
partetal mae-				Characters	n	Mean
Table 98				Bigonial breadth	1	84.00
Many 37	YPE A			at af mandibular ramus	1	54.50
Mean Measurements and Indices	of the Sp	lanchnocranium		Max. breadth of mandibular ramus	1	45.00
Characters				Min. breadth of mandibular ramus	1	35.00
acters	n	Mean		Ht. at mandibular symphysis	1	
Nasal L.		25.5		Mandibular length	1	29.50
Nasal height Nasal breadth	1	39.00		Mandibular angle	1	74.00
preadth	1	26.00		Manufactur	1	121°

Nasion-gnathion line

Nasal height

MEMOIR: No. 9

111.11

TAB	LE 100)			TABLE 103-Contin	u e d	
ADULT FE	MALE .	TYPE A					
Mean Measurements and Maxillary	Indic	es of the	Perman	ient	Characters	n	Mean :
Triuming y	1120101				Nasal breadth	1	26.0
Characters		n	M	ean	Ant. inter-orbital breadth	1	21.0
					Orbital breadth (left)	1	37.5
Mesiodistal crown diam. of	M_2	1	10	0.00	Orbital height (left)	1	35 .5
Labiolingual crown diam. o	$f M_2$	1	6	9.00	Maxillo-alveolar length	1	58.0
Mesiodistal crown diam. of	M_3	1	· 8	3.00	Maxillo-alveolar breadth	1	62.5
Labiolingual crown diam.	of M ₃	1	9	9.50	Palatal length	1	48.5
Crown index of M_2		1	90	0.00	Palatal breadth	1	33.0
Crown index of M_3		1	118	3.75	Orbital index (left)	1	94.67
					Nasal index	1	52.53
,					Maxillo-alveolar index	1	107.76
Таг	BLE 10	1			Palatal index	1	68.04
ADULT FE	10	_					
Mean Measurements and			_	4			
Mandibular	. Mola	es of the r Teeth	Perman	ent	Table 104		
					ADULT FEMALE TYP	E A,	
Characters		'n	Me	an	Mean Measurements and Indices		Mandible
Maria Irana					m.can measurements and marces		
Mesiodistal crown diam. of	M_2	1	10	0.00	Characters	n	Mean
Labiolingual crown diam. o	$f M_2$	1	10	0.00	Characters		
Crown index of M ₂		1	100	0.00	Max. breadth of mandibular ramus	1	37.0
					Min. breadth of mandibular ramus	1	30.0
					Ht. at mandibular symphysis	1	34.0
TAR	LE 102)			Mandibular length	1	75.0
ADULT FEM							
Mean Measurements and			Veurocra	nium			
	indice.	, 0.					
Characters	n	Mean	Min.	Max.	Table 105		
May		185.50	177.0	194.0	ADULT FEMALE TYP	E A	
Max. cranial length	2		177.0		Mean Measurements and Indices	of the	Permanent
Nasion-inion length	1	163.0			Maxillary Molar T	eeth	
Min. frontal breadth	1	95.0					
Horizontal circumference	1	492.0			Characters	n	Mean
							10.00
***		0			Mesiodistal crown diam, of M_1	1	10.00
	LE 10				Labiolingual crown diam, of M_1	1	10.25
ADULT FEM Mean Measurements and In	MALE	of the Sr	lanchnos	eranium	Mesiodistal crown diam, of M_2	1	8.50
mean Measurements and In	arces	or me of	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		Labtolingual crown diam, of M_2	1	11.25
Character			Me	an	Mesiodistal crown diam, of Ma	1	9.00
Characters					abiolingual crown diam, of M	1	10.00
Nasion and the		1	GS	9 ()	Crown index of $M_{ m T}$	1	102.50
Nasion prosthion line Nasion-gnathion line		1	12	1.5	Crown index of M2	1	132.64
gnathion inc					Common haden a se	1	11111

19.5

Crown index of Ma

Table 106 ADULT FEMALE TYPE A

Mean Measurements and Indices of the Permanent Mandibular Molar Teeth

Characters	n	Mean
The state of the s		
Mesiodistal crown diam. of M_1	1	10.25
Labiolingual crown diam, of M_1	1	10.25
Mesiodistal crown diam. of M_2	1	9.75
Labiolingual crown diam. of M_2	1	10.00
Mesiodistal crown diam. of M_3	1	10.00
Labiolingual crown diam. of M_3	1	9.75
Crown index of M ₁	1	100.00
Crown index of M2	1	102.76
Crown index of M ₃	1 ′	97.50

Table 108 ADULT FEMALE TYPE B2 Mean Measurements and Indices of the Splanchnocranium

Characters	ř	Mean
Prosthion-basion line	1	100.00
Nasion-prosthion line	1	58.00
Nasion-gnathion line	1	106.00
Bizygomatic breadth	1	115.00
Nasal height	. 1	43.50
Nasal breadth	1	23.00
Ant. inter-orbital breadth	1	15.00
Orbital breadth (right)	1	41.50
Orbital breadth (left)	1	41.50
Orbital height (left)	1	38.00
Total facial index	1	92.17
Superior facial index	1	50.43
Orbital index (left)	1	91.57
Nasal index	1	52.87

Table 109 ADULT FEMALE TYPE B2 Mean Indices of the Whole Skull

68.70

			Mean marces of the whole Skull					
TABLE 1	07		Characters	'n	Mean			
ADULT FEMALE	TYPE \mathbf{B}_2				cuii			
Mean Measurements and Indice	es of the	Neurocranium	Tr. cranio-facial index	1	85.82			
			Vert. cranio-facial index	1	42.96			
Characters	n	Mean	Long. cranio-facial index	1	59.88			
			Jugo-frontal index	1	78.26			
Max. cranial length	1	167.00						
Max. cranial breadth	1	134.00						
Nasion-inion length	1	166.00	Table 110					
Basion-bregma height	1	135.00						
Vin. frontal breadth	1	90.00	ADULT FEMALE TYPE B2					
Vertical porion height	1	109.00	Mean Measurements and Indices of the Mandible					
ranial module	1	145.33		1	rangible.			
Calculated cranial capacity	1	1204.52	Characters	n	ha.			
ength-breadth index	1	83.24			Mean			
ength-height index	1	80.84	Bigonial breadth	1	-			
readth-height index	1	100.75	Min. breadth of mandibular ramus	1	79.0			
ength-auricular height index	1	65.27	Ht. at mandibular symphysis	1	30.0			
- Callh-allrigular haight index	1	81.34	Mandibular length	1	29.0			
r. fronto-parietal index	1	67.16	Jugo-mandibular index	1	70.0			
	1	07.10	Jugo-manaro mach	1	89.76			

TABLE 111 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean	\pm	S.E.	S. D.	c. v.	Min.	Max.
The second secon								100001 12
Max. cranial length	5	181.90	\pm	4.64	10.39	5.71	167.0	194.0
Max. cranial breadth	2	133.00			_		132.0?	134.0
Nasion-inion length	3	165.33	\pm	1.20	2.08	1.26	163.0	167.0
Basion-bregma height	2	134.50			_		134.0	135.0
Min. frontal breadth	3	93.00	\pm	1.53	2.65	2.85	90.0	95.0
Vertical porion height	2	109.50				_	109.0	110.0
Median sagittal arc	1	376.00				_	_	_
Vert. transversal arc	1	297.00			_	_	_	_
Horizontal circumference	2	498.50			_	_	492.0	505.0?
Cranial module	2	147.58				_	145.33	149.83?
Calculated cranial capacity	2	1251.36					1204.52	1298.21?
Length-breadth index	2	76.08				_	71.93	80.24
Length-height index	2	76.93			-	_	73.02	80.84
Breadth-height index	2	101.14				_	100.75	101.52?
Length-auricular height index	2	62.61					59.95	65.27
Breadth-auricular height index	2	82.34					81.34	83.33?
Tr. fronto-parietal index	2	69.18			-	-	67.16	71.21?
							01.10	

Table 112 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Splanchnocranium

Characters	n	Mean	\pm	S.E.	S. D.	c. v.	Min.	Max.		
Prosthion-basion line	2	100.00			-		100.0	100.0		
Nasion-prosthion line	3	66.00	土	4.05	7.00	10.61	58.0	71.0		
Nasion-gnathion line	3	116.17	<u>-+-</u>	5.09	8.81	7.58	106.0	121.5		
Bizygomatic breadth	2	117.50				_	115.0	120.0?		
Nasal height	5	45.80	土	2.14	4.80	10.48	39.0	51.0		
Nasal breadth	4	24.00	土	1.22	2.45	10.21	21.0	26.0		
Ant. inter-orbital breadth	5	18.20	\pm	0.97	2.17	11.92	15.0	21.0		
Orbital breadth (right)	4	40.50	\pm	0.89	1.78	4.40	38.0	42.0		
Orbital breadth (left)	5	40.60	\pm	0.87	1.95	4.80	37.5	42.5		
Orbital height (right)	3	33.17	土	1.49	2.57	7.75	31.0	36.0		
Orbital height (left)	4	33.62	+	1.88	3.77	11.21	30.0	38.0		
Maxillo-alveolar length	2	57:75					57.5	58.0		
Maxillo-alveolar breadth	3	61.33	土	1.17	2.02	3.29	59.0	62.5		
Palatal length	3	46.83	土	1.43	2.47	5.27	44.0?	48.5		

TABLE 112 - Continued

Characters	n	Mean :	± S.E.	S. D.	c.v.	Min.	Max.
Palatal breadth	4	35.25	± 0.97	1.94	5.50	33.0	37.50
Total facial index	2	96.50			_	92.17	100.83
Superior facial index	2	54.80			_	50.43	59.17?
Orbital index (right)	3	82.74 ±	4.58	7.92	9.57	73.81	88.89
Orbital index (left)	4	83.54 ±	5.58	11.16	13.36	72.94	94.67
Nasal index	4	$53.31 \pm$	5.22	10.43	19.56	41.18	66.67
Maxillo-alveolar index	2	108.23			e-man	107.76	108.70
Palatal index	3	$74.86 \pm$	3.42	5.91	7.89	68.04	78.41?

TABLE 113 ADULT FEMALE COMBINED

Indices of the Whole Skull

Characters	n	Mean
Tr. cranio-facial index	1	85.82
Vert. cranio-facial index	1	42.96
Long. cranio-facial index	1	59.88
Jugo-frontal index	1	78.26

TABLE 114 ADULT FEMALE COMBINED Statistical Constants of the Mandibular Measurements and Indices

Characters	n	Mean <u>+</u>	S.E.	S. D.	c. v.	Min.	Max.
Bigonial breadth	4	83.62 ±	1.84	3.68	4.40	79.00	38.00
Bicondylar breadth	2	111.5		-	-	111.00	112.00
Ht. of mandibular ramus	4	57.75 ±	1.74	3.48	6.03	54.5	6.25
Max, breadth of mandibular ramus	5	$41.50 \pm$	1.30	2.92	7.04	37.00	45.00
Min. breadth of mandibular ramus	6	$31.67 \pm$	1.09	2.66	8.40	29.00	35.50
Ht. at mandibular symphysis	7	30.00 ±	1.04	2.75	9.17	25.00	34.00
Mandibular length	7	$74.71 \pm$	1.95	5.18	6.9 3	67.5	81.50
Mandibular angle	4	122.62° ±	0.98	1.97	1.61	121.00°	125.00°
Mandibular index	2	66.79				60.81	72.77
Breadth index of mandible	2	76.92			-	74.55	79.28
Jugo-mandibular index	2	69.14		accepted.	- Allerana	68.70	69.58?

TABLE 115 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean	±	S.E.	S. D.	c. v.	Min.	Max.
Mesiodistal crown diam. of M_{1}	3	9.92	±	0.51	0.88	8.87	9.00	10.75
Labiolingual crown diam. of M_1	3	11.08	\pm	0.42	0.72	6.50	10.25	11.50
Mesiodistal crown diam. of M_2	4	9.62	\pm	0.42	0.85	8.84	8.50	10.50
Labiolingual crown diam. of M ₂	4	10.19	\pm	0.47	0.94	9.22	9.00	11.25
Mesiodistal crow diam. of $M_{\rm 3}$	4	8.75	\pm	0.48	0.96	10.97	8.00	10.00
Labiolingual crown diam. of M ₃	4		土	0.24	0.48	4.99	9.00	10.00
Crown index of M ₁	3	112.44	土	7.79	13.48	11.99	102.50	127.78
Crown index of M ₂	4	107.25	\pm	9.58	19.16	17.86	90.00	132.64
Crown index of M ₃	4	110.59	土	3.90	7.80	7.05	100.00	118.75

TABLE 116 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Permanent Mandibular Molar Teeth

		2			-			
Characters	n	Mean	土	S.E.	S. D.	c. v.	Min.	Max.
Mesiodistal crown diam. of M_1	2	10.25					10.25	10.25
Labiolingual crown diam, of M_1	2	10.25						
					-		10.25	10.25
Mesiodistal crown diam. of M_2	3	9.75	土	0.14	0.24	2,46	9.50	10.00
Labiolingual crown diam. of M ₂	3	9.83	土	0.16	0.28	2.85	9.50	10.00
Mesiodistal crown diam. of Ma	2	9.50					9.00	10.00
	0	9.12					3.00	10.00
Labiolingual crown diam. of M_3	2				_	-	8.50	9.75
Crown index of M _I	2	100.00					100.00	100.00
Crown index of M2	3	101.11	\pm	0.84	1.46	1.44	100.00	102.76
Cl	2	95.97						05.50
Crown index of M ₃	-	00.0			-	-	94.44	97.50

CEMETERY H STRATUM I (JAR BURIALS)

The skeletal remains of 78 individuals were found in Cemetery H Stratum I (Jar burials), but it was possible to examine only 18 skulls for our purpose. The dimensions relating to the length, width and height of the skulls are given in Collective Tables D-E and their morphological observations in those of J-K.

A. CRANIA

SKL. H 206 (B)

It is undoubtedly a male adult; large mastoid processes, strong muscular attachments are masculine features. The skull is incomplete. Nearly the whole of the forehead and part of the right parietal and temporal regions are missing. Both the orbits and nasal bones are somewhat deformed due to the pressure of earth, but most of the measurements are definite. Seen from above, the skull is almost ovoides with protruding tubera parietalia. From the side, the skull appears to be well vaulted and the occipital curve is rounded. The skull is mesocranic, the length-breadth index being 76.67. The palate is broad and evenly parabolic.

All the teeth have erupted and show considerable abrasion, the upper second left permanent molar being lost post mortem.

SKL. H 255 (a) [Fig. 97; Pl. LXVII: $1\cdot 2$]

Adult male, fairly old in age. The skull is in a good state of preservation. Sagittal and coronal sutures are fully closed. Supraorbital ridges are strongly developed, muscular attachments well marked and mastoids moderately big. The occiput shows a pronounced nuchal line. Norma verticalis appears to be an elongated oval. Norma lateralis shows a very prominent glabello-superciliary region, concave nasal ridge and deep nasion. The forehead ascends little inclined up to the metopian and turns abruptly into a flat vertex which slopes obliquely down to a protruding occiput. There is also evidence of some amount of subnasal prognathism, and the meatus lies nearly midway between nasion and inion. In front view, the upper face is of medium height and breadth, the superior facial index being 54.20 (mesen). Orbits are mesoconchal (right OI 80.00 and left OI 77.01) in form with

a tendency towards chameconchal type, whereas nasal aperture is short and wide, hence chamaerrhine (NI. 52.94). The skull is hyperdolichocranic (L-B Index 68.18), orthocranic (L-AH Index 59.60) and acrocranic (B-AH Index 87.41). Palate is short, *paraboloid* in shape and molars are highly eroded. The mandibular fossae are fairly deep.

The lower jaw is well formed with prominent chin. Ramus fairly low and of medium breadth, the minimum breadth of the ascending ramus being 34.5 mm. The body of the mandible is of medium height, the height measured at the site of the second molar is 28 mm. The mandibular angle formed by the posterior border of the ramus and the lower margin of the corpus is 117°. The internal pterygoidal insertional area forms a deep and extensive concavity. All the teeth have erupted and chewing surface of the molars show considerable attrition.

SKL. H 344 [Fig. 98; Pls. XLII, XLIII, XLVIII]

The skull is of an adult male, fairly old in age as judged from the absorption of the upper alveolar margin. Supraorbital ridges are usually prominent muscular attachments well developed and mastoid processes excessively large. In norma verticalis, the skull is ovoid with some bulging at the parietals. Norma lateralis shows massing of bone at glabella, short and concave nasal ridge, depressed nasal root and slight degree of subnasal prognathism. The forehead is little inclined and the sagittal contour not so well arched. The occiput is prominent but more rounded. Norma facialis shows a moderately wide and low upper face, the superior facial index being 44.12 (hypercuryn). Orbits are low and mesoconchal in form, whereas nose is broad with an index of 57.45 (hyperchamaerrhine). The skull is mesocranic (L-B Index 76.63), orthocranic (L-H Index 73.91) and metriocranic (B-H Index 96.45).

Mandible is incomplete and somewhat distorted. The ascending rami are not preserved and broken off little above the gonial angles. Two molars, two premolars and canine are present on the right side; on the left side are the roots of two incisors. Teeth are worn flat about half the way through the tooth structure proper, the wear being greatest on the buccal side.

SKL. H 61

The skull is very much distorted and compressed. Greater portion of the face including nasal bones, glabellar and inter-orbital regions are defective. The frontal bone is pressed in. The occipitomastoid region is distorted on the left side. There is an opening in the left parietal, measures 37 mm from before backwards and 25 mm from above downwards.

The skull is small in size and globular in form. The vertical porion height is 117 mm. Compared to size of the skull slight increase in the vertical porion height is due to occipito-mastoid having been compressed and pushed upwards a few millimeters. The right orbit is squarish and mesoconchal (OI 80.95) in form.

SKL. H 153 (a)

The skull is not well preserved and only a few measurements could be taken accurately. Portion of the frontal bone, upper alveolar margin with all the teeth are missing. Orbits are compressed from above downwards and left parieto-temporal area shows some amount of sidewise compression. Similarly due to pressure, bones of the base have been forced slightly upwards and inwards. The length of the skull is 181 mm. In comparison with length and width the height of the skull appears to be low, the basion-bregma height being about 117 mm, the distorted base making an exact estimate impossible. The forehead is vertical and narrow (minimum frontal breadth 89 mm). The cranial contour in norma verticalis is byrsoides in shape.

SKL. H 154 (a) [Pl. XLIX]

It is an adult female. The skull is incomplete and rather small and gracile in structure. The entire sphenoid bone and the lower border of right orbit are missing. The maxilla lacks the left posterior portion and associated teeth; it retains a right PM_2 , right M_2 and M_3 . In norma verticalis the skull is more or less ellipsoidal in shape, the forehead relatively narrow, parietal eminences being somewhat prominent. From the side, the forehead ascends in a smooth arc into a convex vertex which slopes obliquely down to a prominent tuber occipitale. The facial region is moderately narrow and supraorbital ridges absent. The skull is delichocranic (L-B Index 72.73), hypsicranic (L-H Index 78.98) and acrocranic (B-H Index 108.59). Its cranial capacity is 1300.50 cc.

Mandible is incomplete. Nearly whole of the left half is missing. It is gracile in general appearance. Muscular markings are mildly developed. The mandibular length is 88 mm and the genial angle seems to be evenly rounded. Minimum breadth of the ascending ramus measures 34 mm. The vertical depth of the horizontal ramus in the region of second molar is 26 mm.

SKL. H 206 (d) [Pl. LXVII: 3-1]

The skull belongs to an adult female, between 25 and 30 years of age. It is incomplete having a broken mandible. Supraorbital ridges are only perceptible mesially with feeble muscular impression. Basilar suture is closed while vault sutures are open. Viewing from the top it is ovoides in outline, expanded at the parietals. From the side, the forehead is slightly receding passing back in a uniform expanded at the general contour of the vault; the occipital region is slightly protruding. The skull curve into the general contour of the vault; the occipital region is slightly protruding. The skull is mesocranic (L-B Index 76.88), chamaecranic (L-H Index 72.25) and metriocranic (B-H Index 93.98). The face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left The face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chamaeconchic (left To face is low in relation to its width. Orbits are somewhat rectangular and chama





Fig. 97 Skl. H 255 (a)




MID-SAGITTAL CURVE (1/2 NATURAL SIZE)



Norma Frontalis

DIOPTOGRAPH TRACING (WASH) : MALE SKL. H 344 (Natural Size)



Norma Lateralis

DIOPTOGRAPH TRACING (WASH): SKL. H 344 (Slightly Reduced)

Mandible is medium in size with little muscular markings. Mylohyoid ridge and pterygoid attachments are indistinct. Both the ascending rami are not preserved, the right ascending ramus including portion of the mandibular body is missing. The ascending ramus is not very high (50.5 mm) and wide (30 mm). The mandibular angle is rounded (123°).

SKL. H 245 (e) [Fig. 99]

The skull is incomplete (but reconstructed) and lacks nearly entire basis cranii, both the zygomatic arches, part of the frontal bone including supraorbital margins and nasal bones. The whole cranial surface is very much eroded. The sex is difficult to determine but the extremity bones are rather gracile in structure. It is therefore assumable that the skull belongs to a female. The age




Fig. 99

appears to be fairly high; most of the portion of sagittal suture is ossified. Norma verticalis is wide ovoid, narrowed in front. In norma temporalis the frontal bone ascends obliquely and passes into a somewhat domed vertex. The occiput is fairly rounded with no actual protuberance. The cranial index is 84.00, which places it within the brachycranic or broad-headed class.

SKL, H 246 (c) [Fig. 102]

The skull belongs to an adult female. Supraorbital ridges are median in type, muscle attachments rather small and left mastoid process is small in size. The right zygomatic process, greater part of the maxillae and palate are missing. Norma verticalis is pentagonoid with prominent parietal part of the maxillae and palate are missing. Norma verticalis is pentagonoid with prominent parietal part of the maxillae and palate are missing. Norma verticalis is pentagonoid with prominent when eminences. In norma temporalis the forehead is low and practically vertical to about metopian when it slopes back posteriorly into a low-arched vertex. The occiput is curved with a fairly prominent tuber occipitale. Norma occipitalis is house shaped with sides converging slightly downwards. The nasal ridge is short and concave with a slight depression at the root. The skull is dolichocranic (L-B Index 74.18), orthocranic (L-AH Index 60.99) and metriocranic (B-AH Index 82.22). Intracranial capacity is 1324.35 cc.

SKL. H 247 (a)

Adult skull, between 21 and 25 years. Posterior part of the basis cranii is absent and the skull is flattened at the left side. Low supraorbital ridges and small mastoids suggest it to be a female. All the cranial sutures are open. Viewed from the top, the skull is sphenoides in shape and the right zygomatic arch slightly bowed out while left is broken. The skull appears to be elongated but the cranial index could not be worked out due to absence of landmarks for breadth measurement. It is orthocranic in both the indices (L-H Index 73.86 and L-AH Index 61.65). Forehead is receding and from ophryon it passes backwards. The proportion of length and breadth of the face is middle, superior facial index being mesen (53.36). The right orbit is mesoconchic (OI 79.52). The nose is flat and chamaerrhine (NI 54.17). Subnasal protuberance is marked with muzzle-shaped alveolar region. The palate is deep and narrow and upsiloid in shape.

SKL. H 247 (b) [Pl. LXVI]

The skull is that of an adult female. The cranial sutures are open and comparatively simple. Main parts missing are, the occipital portion of the base, frontal process of the left maxilla, left malar and much of the alveolar margins.

The skull is relatively small in size and shows minimum of bony relief. The cranial capacity is estimated to have been 1253.44 cc. The vault is moderately low (the vertical porion height being 108 mm); while the cranial index is in the border of dolicho- and mesocranic (L-B Index 74.72). The supraorbital ridges are feeble, and the forehead is narrow and almost vertical. Nasal bones are wingshaped with a little or no depression of the nasal root. The upper face height, from nasion to prosthion, measures 60 mm only.

SKL. H 7435 (b) [Figs. 100, 101]

The skull belongs to an adult female, aged approximately 25 years. The right temporo-parietal region is somewhat compressed, the glabellar portion of the forehead including the upper part of the nasal bones is damaged, while the zygomatic arches are entirely missing. All the cranial sutures



0




Fig. 101





Fig. 102 Skl. H 246 (c)



MID-SAGITTAL CURVE (1/2 NATURAL SIZE)

156

are open. Muscular attachments and mastoids are weakly developed. The same is true of supraorbital ridges. When viewed from top, the skull is almost *ellipsoidal* with slightly expanded parietal
eminences. Viewed from the side, the forehead is slightly inclined and passes back into a somewhat
flat vertex. The vault of the skull is strikingly low as shown by the auricular height of the skull
(100.00 mm). There appears to be a small pre-lambdoid flattening, and an occipital bun is moderately
discernible. There is also some amount of sub-nasal prognathism. The maximum cranial length is
178.0 mm, the right parietal region is slightly compressed so that length-breadth index cannot be
derived accurately; but the skull is certainly dolichocranic. The length-auricular height proportion is
within the chamaecranic category, with an index of 56.18. The orbits are rectangular, inclined laterally
downwards, and chamaeconchic.

SKL. Pot No. 11 [Fig. 103]


The calvaria is small and practically devoid of post mortem defects. Supraorbital ridges are indistinct, mastoid processes small and the muscle attachment weak. All the cranial sutures are open but spheno-basilar is united. This, together with the state of dentition suggests that the skull belongs to an adult female about 25 years of age. In norma verticalis the skull is an elongated oval, slightly expanded at the parietals. Seen from the side, the forehead rises almost vertically, but from the ophryon sweeps back into relatively low arched vertex with a small pre-lambdoid flattening. The occiput is convex with a moderately protruding tuber occipitale. Nasion is not deep and the nasal ridge short and concave. In front view, the upper face is of medium breadth, the superior facial index being 52.00 (mesen) and nose is mesorrhine with an index of 48.48. The skull is mesocranic (L-B Index 75.72) and orthocranic (L-AH Index 59.54).

SKL. Pot No. 12 (a) [Pis. XLIV, XLV]


The skull appears to be that of an young adult of less than 25 years of age. It is small in size, the calculated cranial capacity being 1203.30 cc. Vault sutures as well as spheno-basilar suture are not ossified. The sex is undoubtedly female; small mastoids, poor glabella and arcus superciliaries are feminine features. The skull is notably smooth in surface relief. In norma verticalis, the cranial contour is byrsoides in shape. In norma temporalis the forehead rises vertically and from obelion slopes backwards in the parieto-occipital region. The occipital squama is prominent and projected behind the inion. The vault of the skull is relatively low, basion-bregma height being 119 mm only.

SKL. Pot No. 12 (b)

Adult female. The skull is small rather gracile in structure. It is incomplete. The entire facial region including maxillae and palate, part of the frontal bone and zygomatic arches are absent. The cranial surface is eroded and all the cranial sutures cannot be seen perfectly. As far as can be seen all the sutures appear to be un-united. The head shape, seen from above, is elliptical. In profile the vertex is well arched and the occiput receds obliquely with pronounced occiput. Norma verticalis is house-shaped with slightly downward converging side walls.



Norma Frontalis



Date latts

DIOPTOGRAPH TRACING (WASH) : FEMALE SKL. Pot No. 12(a) (Natural Size)

ANALYSIS

Of the 18 crania measured 3 were adult males, 12 adult females, 2 juveniles and 1 infant. On the basis of metric and visual observations 4 adult crania (2 males and 2 females) have been classified as belonging to the long-headed Type A, 9 adult crania (all female) as small-headed Type A_2 and 2 adult crania (1 male and 1 female) as round-headed Type B_2 .

Distribution of the adult crania according to type and sex is as follows:

SEX	TYPE A	TYPE A,	TYPE \mathbf{B}_2
MALE	H 255 (a) H 344		Н 206 (В)
FEMALE	H 246 (c) H 7435 B	H 154 (a) H 247 (a) H 153 (a) H 247 (b) H 61 Pot No. 11 H 206 (d) Pot No. 12 (a) Pot No. 12 (b)	H 245 (e)

TYPE A

In Tables 120-125, 136-138 the mean measurements and indices of the male and female Type A skulls from Cemetery H st II Jar burials are given in detail. The salient cranio-facial features of the Type A skulls from jar burials are as follows:

	M	IALE	FEMA	LE:
Length-breadth index Length-height index Breadth-height index Length-auricular height index Breadth-auricular height index Transverse fronto-parietal index Superior facial index Orbital index	Dolichocranic Orthocranic Metriocranic Orthocranic Metriocranic Eurymetopic Euryn Mesoconchic	(72.40) (70.80) (97.86) (62.14) (85.90) (70.17) (49.16) (77.56 ± 1.05)	Dolichocranic Chamaecranic Metriocranic Orthocranic Tapeinocranic Eurymetopic — Hypsiconchic Chamaerrhine	(73.32) (69.96) (95.35) (58.58) (79.87) (70.82) — (84.62) (53.41)
Breadth-height index Length-auricular height index Breadth-auricular height index Transverse fronto-parietal index Superior facial index	Orthocranic Metriocranic Eurymetopic Euryn	(62.14) (85.90) (70.17) (49.16)	Orthocranic Tapeinocranic Eurymetopic — Hypsiconchic	(58.5 (79.5 (70.5 (84.6

In Plate XLVI and in Figures 104-106 are presented photographs and dioptographic drawings of Type A male cranium. In the male strong supraorbital ridges, relatively flat vertex, prominent occiput, highly depressed nose, rather short and moderately wide nasal aperture and prominent cheek bones are well represented. Viewed from the top, the skull looks like an elongated oval. In the front view, the upper face is of medium height and width.

TYPE A₂

In Tables 139 to 144 are presented the mean cranial measurements and indices of the female skulls belonging to Type A_2 . Any male skull representing this type is conspicuously absent.

The essential cranio-facial features of female belonging to Type A_2 is as follows:

Length-breadth index	Mesocranic	(75.15)	\pm	0.70)
Length-height index	Orthocranic	(71.36)	\pm	1.68)
Breadth-height index	Acrocranic	(98.42	\pm	2.61)
Length-auricular height index	Orthocranic	(61.00	\pm	0.58)
Breadth-auricular height index	Metriocranic	(82.80	\pm	2.29)
Superior facial index	Mesen	(51.36	\pm	1.38)
Orbital index	Mesoconchic	(81.09	土	1.44)
Nasal index	Chamaerrhine	(55.44	\pm	2.23)
Cranial Capacity	Euencephal	(1235.7	6	± 24.24)
Palatal index	Brachystaphylin	(85.37)		
Transverse fronto-parietal index	Eurymetopic	(70.07	\pm	2.00)

Female skulls of Type A_2 from jar burials are dolichocranic to sub-mesocranic, with an average cranial index range of 73-75. They represent a people who are relatively small-headed with a small cranial capacity, averaging 1235.76 cc for the females. In Plate XLVII and Figures 107-109 are given photographs and dioptographic drawings of the female belonging to Type A_2 in jar burials. The sagittal contour shows that the vault of the skull is low with moderately protruding occiput. The forehead is narrow and vertical.

TYPE Bo

Only two crania (1 male and 1 female) have been classified as Type B₂. Both the skulls are damaged. Morphologically these skulls closely conform to round-headed skulls found at Cemetery H Open Burials.

DISCUSSION

From the foregoing account of the crania, it appears that a new physical type is represented in Cemetery H st I (jar burials). It is a small-brained, weakly mesocephalic people, designated by us as Type A_2 . In addition, four long-headed skulls [Nos. H 255 (a), H 344, H 246 (c) and H 7435 (b) I seem to resemble, both metrically and morphologically, Type A skulls of Cemetery R 37 and two round-headed skulls [Nos. H 206 (B) and H 245 (e) correspond more closely with the round-headed Type B_2 (open burials) than the round-headed Type B_1 (Area G).

On the otherhand, crania of Type A_2 skulls are remarkable on account of small size, sub-rounded head (the skulls gravitate around the indices of 73-75), low vault with small face, vertical and pinched forehead, poorly developed supraorbital ridges and feeble muscular attachments.

Differences in mean measurements between Type A_2 finds of Cemetery H st I (jar burials) and Type A_1 females of Cemetery R 37 is clearly demonstrated in Table 117.

Table 117 Showing deviations of Type A_2 from Type A_1 in salient cranio-facial dimensions

Characters	Female Type A ₁ Cemetery R 37	Female Type A ₂ Cemetery H (Jar burials)	Difference
Max. cranial length	180.08	175.71	- 4.37
Max. cranial breadth	129.92	129.43	- 0.49
Vertical porion height	111.36	108.80	- 2.56
Median sagittal arc	374.80	354.33	-20.47
Vertical transversal arc	308.40	289.42	-18.98
Cranial module	147.33	143.97	- 3.36
Cranial capacity	1281.12 cc	1235.76 cc	-45.36
Nasion-prosthion length	62.33	59.80	- 2.53
Bizygomatic breadth	124.00	121.50	- 2.50
Nasal height	48.90	46.50	- 2.40
Nasal breadth	25.10	25.00	- 0.10
Palatal length	42.00	41.50	- 0.50
Palatal breadth	39.33	35.00?	- 4.33

It appears from Table 117 that the Type A_2 is smaller than Type A_1 in all major dimensions. Though metrically there is no indication of close similarity between Type A_1 and Type A_2 , nevertheless in general appearance the difference between Type A_2 and Type A_1 is not remarkably large. One of the important problem now emerges whether Type A_2 may be considered as variant of Type A_1 of Cemetery R 37 or they represent a separate division. Type A_2 skulls may be identified as a smaller form of Classic Mediterranean (to use Coon's expression) and in all probability they were not identical with earlier population of Harappan culture.

B. EXTREMITY BONES

Since most of the burial jars of Cemetery H st I contained fractional burials, the quantity of post-cranial bones deposited in them was less in comparison to regular burials of R 37 and Cemetery H st II. The extremity bones in perfectly measureable condition were poorly represented in the jar




Fig. 104 Norma Frontalis




Fig. 105 Norma Verticalis




Fig. 106 Norma Lateralis

Figs. 104, 105 and 106. Dioptographic contours of Type A Male [Skl. H 255 (a)] of Cemetery H St. I (Jar burials)




Norma Frontalis

Norma Verticalis



Norma Lateralis

Photographs of Type A Male [Skl. H 255(a)] of Cemetery H St. I (Jar burials)




Norma Frontalis



Norma Verticalis

Mar



Norma Lateralis

Photographs of Type A2 Female [Skl. Pot No. 11] of Cemetery H St. I (Jar burials)




Fig. 107 Norma Frontalis




Fig. 108 Norma Verticalis




Fig. 109 Norma Lateralis

Figs. 107, 108 and 109. Dioptographic contours of Type A_2 Female $[\,Skl.\,$ Pot $\,No.\,$ 11 $]\,$ of Cemetery H St. I $\,(Jar\,$ burials)

burials. Primary data on them have been produced in the Collective Table N, while their average values have been shown in Tables 118 and 119.

HUMERUS

Three humeri (two right and one left) of two females are measured.

TABLE 118

Character	Female				
	n	Range	Mean		
Maximum length	3	279.0-319.0	305.33		
Breadth of proximal epiphysis	3	43.0- 45.0	43.83		
Breadth of distal epiphysis	3	53.5-55.5	54.33		
Circumference of the shaft at the middle	2	52.0- 59.0	55.50		
Minimum circumference of diaphysis	3	51.0- 55.0	52.33		
Robusticity index	3	15.99-18.28	17.19		





Fig. 110 Skl. H 206 (d)
Female: (Rt)
Volar projection

Fig. 111 Skr. H 206 (d) Female: (Rt) Lateral projection

In relation to open burial female humeri of st II, those of jar burials st I were longer in size. Robusticity index, however, reveals that jar burial humeri (17.19) were weaker than open burial (19.14). The range of the former varies from 15.99 to 18.28 and that of the latter between 18.46 to 20.22.

Septal aperture is found in two cases, viz., Skl. H 206 (d) and in a juvenile H 206 (a) (both being right humeri).



Frontal view



Vertical view




Left lateral view

SKL. H 344 : CEMETERY H St. 1 (JAR BURIALS)




Frontal view


111



Vertical view



Left leteral view



Occipital view

SKL, H 154(a) : CEMETERY H St. I (JAR BURIALS)

RADIUS AND ULNA

Measurements of a single radius (left) of a female individual [Skl. H 247 (a)] and one ulna of a male [Skl. H 601 (a), left] are shown in Collective Table N. Another ulna [H 206 (d), Figs. 110 & 111], without its styloid process, shows a physiological length of 223 mm and a minimum circumference of diaphysis of 32 mm.

FEMUR

Only two femora of two females [Skl. H 154 (a), H 206 (c) i are measured of which mean values are tabulated in Table 119.

Table 119

F	0	~	٠	12

Character	Female				
	n	Range	Mean		
Sagittal diameter at middle of diaphysis	2	25.5 - 26.00	25.75		
Transversal diameter at middle of diaphysis	2	25.0 - 28.00	26.50		
Circumference of the shaft at the middle	2	80.0 - 83.00	81.50		
Pilastric index	2	91.07 - 104.00	97.54		
Platymerie index	2	73.33 - 85.71	. 79.52		

None of the femora yielded the measurement of maximum length. In both the bones anteroposterior diameter of the proximal shaft is slender in relation to the transverse diameter at the same point; in platymeric index, one is platy- and the other is eurymeric. Pilastric index indicates primitiveness when compared with other modern populations (Wilder 1920: 126).

Only one female tibia (Skl. H 61, left) has been measured whose maximum legnth is $365~\mathrm{mm}$ and robusticity index is 20.61.

STATURE

Stature of a single male |Skl. H 601 (a)| and three females |Skl. H 61, H 206 (c)| and |Skl. H 601 (a)| and three females |Skl. H 61, H 206 (c)| and |Skl. H 601 (a)| and |Skl. H 601 (a)| and three females |Skl. H 61, H 206 (c)| and |Skl. H 601 (a)| and |Skl

STATISTICAL CONSTANT CEMETERY H (JAR BURIALS)

TABLE 120

ADULT MALE TYPE A

Mean Measurements and Indices of the Neurocranium

Characters	n	Mean	Min.	Max.
Max. cranial length	2	191.00	184.0?	198.0
Max. cranial breadth	2	138.00	135.0	141.0
Nasion-inion line	2	172.50	165.0	180.0
Basion-bregma height	2	135.00	134.0	136.0
Min. frontal breadth	2	98.75	95.0	98.5
Vertical porion height	2	118.50	118.0	119.0
Median sagittal arc	2	381.50	379.0	384.0
Vertical transversal arc	2	311.50	307.0	316.0
Horizontal circumference	2	533.50	522.0	545.0
Cranial module	2	154.67	153.67	155.67
Calculated cranial capacity	2	1470.89	1459.13	1482.65
Length-breadth index	2	72.40?	68.18	76.63?
Length-height index	2	70.80?	67.68	73.91?
Breadth-height index	2	97.86	96.45	99.26
Length-auricular height index	2	62.14?	59.60	64.67?
Breadth-auricular height index	2	85.90	84.40	87.41
Tr. fronto-parietal index	2	70.17	67.38	72.96

Table 121 ADULT MALE TYPE A Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	Min.	Max.
Prosthion-basion line	2	102.50	98.0	107.0
Nasion-prosthion line	2	65.50	60.0	71.0
Nasion-gnathion line	1	123.00		
Bizygomatic breadth	2	133.50	131.0	136.0
Nasal height	2	49.00	47.0	51.0
Nasal breadth	2	27.00	27.0	27.0
Ant. inter-orbital breadth	2	22.00	22.0	22.0
Orbital breadth (right)	2	40.75	39.0	42.5
Orbital breadth (left)	2	41.75	40.0	43.5
Orbital height (right)	2	32.25	30.5	34.0
Orbital height (left)	2	31.75	30.0	33.5
Maxillo-alveoler length	2	56.00	56.0	56.0
Maxillo-alveolar breadth	1	63.00		
Palatal length	1	47.00		
Palatal breadth	1	41.00	_	

TABLE 121 - Continued

Characters	n	Mean	Min.	Max.
Total facial index	1	93.89	_	_
Superior facial index	2	49.16	$44.12^{'}$	54.20
Orbital index (right)	2	79.10	78.21	80.00
Orbital index (left)	2	76.00	75.00	77.01
Nasal index	2	55.20	52.94	57.45
Maxillo-alveolar index	1	112.50		

Table 122

ADULT MALE TYPE A

Mean Indices of the Whole Skull

Characters	n	Mean	Min.	Max.
~			00.45	97.04
Tr. cranio-facial index	2	96.74	96.45 41.91	52.99
Vert. cranio-facial index	2	47.45	53.26	54.04
Long. cranio-facial index	2	53. 6 5 72.52	69.85	75.19
Jugo-frontal index	2	72.52	00.00	

Mean Measurements and Indices of the Mandible

Characters	n	Mean	Min.	Max.
Bigonial breadth a Bicondylar breadth	2	87.25 111.00	74.50 —	100.00
Ht. of mandibular ramus Max. breadth of mandibular ramus Min. breadth of mandibular ramus	1 1 2	62.00 46.00 34.75	34.50	35.00
Ht. at mandibular symphysis Mandibular length	2 2 2	27.75 84 .25 121°	24.00 .80.50 117°	31.50 88.00 125°
Mandibular angle Mandibular index Breadth index of mandible Jugo-mandibular index	1 1 2	79.28 90.09 65.56	54.78	76.34

Table 124 ADULT MALE TYPE A

Mean	Measurements	and	Indices	of	the	Permanent
	Maxill	arv	Molar ?	Feetl	1	

Characters	n	Mean
Mesiodistal crown diam. of M_1	1	11.00
Labiolingual crown diam, of M_1	1	12.00
Mesiodistal crown diam, of M_2	1	10.50
Labiolingual crown diam. of M_2	1	11.75
Mesiodistal crown diam. of M_3	1	9.00
Labiolingual crown diam. of M ₃	1	11.00
Crown index of M ₁	1	113.64
Crown index of M2	1	112.28
Crown index of M ₃	1	122.22

TABLE 125

ADULT MALE TYPE A

Mean Measurements and Indices of the Permanent Mandibular Molar Teeth

Mean
10.50
11.50
11.50
10.50
109.52
91.30

TABLE 126 ADULT MALE TYPE B2

Mean Measurements and	Indices of	the	Neurocranium
Characters		n	Mean
Max. cranial length		1	180.07
Max. cranial breadth		1	138.0
Vertical transversal arc		1	312.0
Herizental circumference		1	140.0
Length-breadth index		1	76.6 7?

Table 127 ADULT MALE TYPE \mathbf{B}_2

Mean Measurements and Indices of the Splanchnocranium

Characters		n	Mean
Maxillo-alveolar	length	1	50.0
Maxillo-alveolar	breadth	1	62.0
Palatal length		1	42.0
Palatal breadth		1	39.0
$Maxillo\hbox{-} alveolar$	index	1	124.00
Palatal index		1	92.86

Table 128

ADULT MALE TYPE \mathbf{B}_2

Mean Measurements and Indices of the Permanent Maxillary Molar Teeth

Characters	n	Mean
Mesiodistal crown diam, of M ₁	1	10.75
Labiolingual crown diam, of $ m M_1$	1	12.25
Mesiodistal crown diam. of M_2	1	9.50
Labiolingual crown diam, of $ m M_2$	1	12.50
Mesiodistal crown diam, of M ₃	1	9.50
Labiolingual crown diam, of $ m M_{\odot}$	1	11.50
Crown index of M ₁	1	113.96
Crown index of M ₂	1	131.58
Crown index of Ma	1	122.40

Table 129

ADULT MALE TYPE B2

Mean Measurements and Indices of the Permanent Mandibular Molar Teeth

Characters	n	Mean
Mesiodistal crown diam, of M_{I}	1	12.00
Labiolingual crown diam, of M ₁	1	11.25
Mesiodistal crown diam, of M2	1	10.75
Labiolingual crown diam, of $ m M_2$	1	11.25
Mesiodistal crown diam, of $ m M_3$	1	10.00
Labiolingual crown diam, of $ m M_3$	1	10.75
Crown index of M ₁	1	93.82
Crown index of M2	1	104.66
Grown index of M ₃	1	107.64

TABLE 130

ADULT MALE COMBINED

Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean	<u>+</u> S.E.	S. D.	c. v.	Min.	Max.
Max. cranial length	3	187.33	\pm 5.46	9.45	5.04	180.00	198.00
Max. cranial breadth	3	138.00	\pm 1.73	3.00	2.17	135.00	141.00
Nasion-inion length	2	172.50		_	_	165.00	180.00
Basion-bregma height	2	135.00			_	134.00	136.00
Min. frontal breadth	2	96.75		_	_	95.00	98.50
Vertical porion height	2	118.50		-		118.00	119.00
Median sagittal arc	2	381.50		_		379.00	384.00
Vert. transversal arc	3	311.67	± 2.61	4.51	1.45	307.00	316.00
Horizontal circumference	3	523.67	± 11.88	20.55	3.92	504.00	545.00
Cranial module	2	154.67		_		153.67	155.67
Calculated cranial capacity	2	1470.89				1459.13	1482.65
Length-breadth index	3	73.83	± 2.83	4.89	6.62	68.18	76.67?
Length-height index	2	70.80		_		67.68	73.91?
Breadth-height index	2	97.86				96.45	99.26
Length-auricular height index	2	62.14				59.60	64.67?
Breadth-auricular height index	2	85.90				84.40	87.41
Tr. fronto-parietal index	2	70.17			_	67.38	72.96

 ${\bf TABLE~131}$ ${\bf ADULT~MALE~COMBINED}$ Statistical Constants of the Measurements and Indices of the Splanchnocranium

Characters	n	Mean <u>+</u>	S.E.	S. D.	c. V.	Min.	Max.
						98.00	107.00
Prosthion-basion line	2	102.50		-		60.00	71.00
Nasion-prosthion line	. 2	65.50					
Nasion-gnathion line	1	123.00			-	131.00	136.00
Bizygomatic breadth	2	133.50					
Nasal height	2	49.00			_	47.00	51.00
Nasal breadth	2	27.00			-	27.00	27.00
	2	22.00			****	22.00	22.00
Ant. inter-orbital breadth	2					39.00	42.50
Orbital breadth (right)		40.75				40.00	43.50
Orbital breadth (left)	2	41.75			-	30.50	34.00
Orbital height (right)	2	32.25			-	30.00	
Orbital height (left)	2	31.75				50.00	33.5 0
Maxillo-alveolar length	3	54.00	2.00	3.46	6.41		56.00
	2	62.50			2.000	62.00	63.00
Maxillo-alveolar breadth	2	44.50		-		42.00	47.00
Palatal length						39.00	41.00
Palatal breadth	2	40.00					

TABLE 131 - Continued

Characters	n	Mean	S. D.	c. v.	Min.	Max.
Total facial index	1	93.89		_	_	
Superior facial index	2	49.16	-	_	44.12	54.20
Orbital index (right)	2	79.10	_	_	78.21	80.00
Orbital index (left)	2	76.00		_	75.00	77.01
Nasal index	2	55.20		_	52.94	57.45
Maxillo-alveolar index	2	118.25	_	_	112.50	124.00
Palatal index	1	92.86	_			_

TABLE 132
ADULT MALE COMBINED

Mean Indices of the Whole Skull

Characters n Mean Min.	Max.
1.1	
Tr. cranio-facial index 2 96.74 96.45	97.04
Vert. cranio-facial index 2 47.45 41.91	52.99 •
Long. cranio-facial index 2 53.65 53.26	54.04
Jugo-frontal index 2 72.52 69.85	75.19

TABLE 133

ADULT MALE COMBINED

Statistical Constants of the Mandibular Measurements and Indices

						**		
n	Mean	<u>+</u>	S.E.		S. D.	c. v.	Min.	Max.
2	87.25				-		74.50	100.00
1	111.00				No.	- Constitution of the Cons		
1	62.00	±	± 2.68		_	Militar		-
3 1	46.00							_
2	34.75						34.50	35.00
3	2 9 .33				4.64	15.82	24.00	32.50
2	84.25					-	80.50	88.00
2	121°				(1) (1) (1) (1)		117°	125°
1	79.28						manufine.	
\underline{I}_{il}	90,09	£			-		-	
2	65.56				-		54.78	76.34
	2 1 1 5 1 5 2 3 2 2 1	2 87.25 1 111.00 1 62.00 6 1 46.00 6 2 34.75 3 29.33 2 84.25 2 121° 1 79.28 1 90,09	2 87.25 1 111.00 1 62.00 6 1 46.00 6 2 34.75 3 29.33 ± 2 84.25 2 121° 1 79.28 1 90,09	2 87.25 1 111.00 1 62.00 5 1 46.00 5 2 34.75 3 29.33 ± 2.68 2 84.25 2 121° 1 79.28 1 90,09	2 87.25 1 111.00 1 62.00 5 1 46.00 5 2 34.75 3 29.33 ± 2.68 2 84.25 2 121° 1 79.28 1 90,09	2 87.25	2 87.25 1 111.00 1 62.00 5 1 46.00 5 2 34.75 3 29.33 ± 2.68 4.64 15.82 2 84.25 2 121° 1 79.28 1 90,09	2 87.25 — 74.50 1 111.00 — — — — 1 62.00 — — — — — 5 1 46.00 — — — 34.50 3 29.33 ± 2.68 4.64 15.82 24.00 2 84.25 — 80.50 2 121° — 117° 1 79.28 — — — —

Table 134

Adult male combined

Statistical Constants of the Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean	Min.	Max.
Mesiodistal crown diam, of M_1	2	10.88	10.75	11.00
Labiolingual crown diam. of M ₁	2	12.38	12.25	12.50
$Mesiodistal$ crown diam. of M_2	2	10.00	9.50	10.50
Labiolingual crown diam. of $ m M_2$	2	12.12	11.75	12.50
Mesiodistal crown diam. of M ₃	2	9.25	9.00	9.50
Labiolingual crown diam. of M ₃	2	11.25	11.00	11.50
Crown index of M ₁	2	113.80	113.64	113.96
Crown index of M2	2	121.93	112.28	131.58
Crown index of M_3	2	122.31	122.22	122.40
Crown index of M ₃	2	122.31	122.22	122.40

Table 135

Adult Male Combined

Statistical Constants of the Measurements and Indices of Permanent Mandibular Molar Teeth

n	Mean	Min.	Max.
2	11.25	10.50	12.00
2	11.38	11.25	11.50
2	11.12	10.75	11.50
2	10.88	10.50	11.25
1	10.00	-	-
1	10.75	_	
2	101.67	93.82	109.52
2	97.98	91.30	104.66
1	107.64	-	
	2 2 2 2 2 1 1 1 2	2 11.25 2 11.38 2 11.12 2 10.88 1 10.00 1 10.75 2 101.67 2 97.98	2 11.25 10.50 2 11.38 11.25 2 11.12 10.75 2 10.88 10.50 1 10.60 — 1 10.75 — 2 101.67 98.82 2 97.98 91.30

TABLE 136

ADULT FEMALE TYPE A

Mean Measurements and Indices of the Neurocranium

Characters	'n	Mean	Min.	Max.
			170.0	100.0
Max. cranial length	2	180.00	178.0	182.0
Max. cranial breadth	2	132.00	129:0?	135.0
	4		165.0	167.0
Nasion-inion length	2	166.00		
Basion-bregma height	2	126.00	117.0	135.0
Min. frontal breadth	-	93.50	91.0	96.0
	2		100.0	111.0
Vertical porion height	2	105.50	100.0	111.0

TABLE 136 - Continued

Characters	n	Mean	Min.	Max.
Median sagittal arc	2	360.50	350.0	371.0
Vertical transversal arc	2	290.50	283.0	298.0
Horizontal circumference	2	536.00	499.0	573.0
Cranial module	2	146.00	141.33?	150.67
Calculated cranial capacity	1	1324.35	_	_
Length-breadth index	2	73.32?	72.47?	74.18
Length-height index	2	69.96	65.73	74.18
Breadth-height index	2	95.35?	90.70?	100.00
Length-auricular height index	2	58.58	56.18	60.99
Breadth-auricular height index	2	79.87?	77.52?	82.22
Tr. fronto-parietal index	2	70.82?	70.54?	71.11

Table 137 ADULT FEMALE TYPE A Mean Measurements and Indices of the Splanchnocranium

Characters	n	Mean	Min.	Max.
Prosthion-basion line	1	100.50?	_	_
Nasion-prosthion line	1	57.00	_	_
Nasal height	1,	44.00?		
Nasal breadth	1 '	23.50	_	
Ant. inter-orbital breadth	2	19.25	18.5	20.0
Orbital breadth (right)	2	39.00	39.0	39.0
Orbital breadth (left)	2	40.25	39.0	41.5
Orbital height (right)	1	33.00		-
Orbital height (left)	1	30.00		
Maxillo-alveolar length	1	57.00		
Maxillo-alveolar breadth	1	64.00		-
Orbital index (right)	1	84.62		
Orbital index (left)	1	72.29		
Nasal index	1	53.41?		
Maxillo-alveolar index	1	112.28		

Table 138 ADULT FEMALE TYPE A

Mean Indices of the Whole Skull

Characters			n	Mean
Vert. cranio-facial Long. cranio-facial	index	*	1	48.72? 56.46?

Table 139 ${\tt ADULT\ FEMALE\ TYPE\ A_2}$ Mean Measurements and Indices of the Neurocranium

Characters	n	Mean	±	S. E.	Min.	Max.
Max. cranial length	7	175.71	土	1.15	173.0	181.0
Max. cranial breadth	7	129.43	\pm	1.85	119.0	133.0
Nasion-inion length	5	166.50	\pm	1.80	162.0	173.0
Basion-bregma height	9	126.39	\pm	2.46	117.0?	139.0
Min. frontal breadth	7	91.50	\pm	2.31	85.0	104.0
Vertical porion height	5	108.80	\pm	2.27	103.0	117.0
Median sagittal arc	6	354.33	\pm	2.62	344.0	362.0
Vertical transversal arc	6	289.42	\pm	3.26	282.0	304.0
Horizontal circumference	8	492.62	\pm	2.40	483.0	504.0
Cranial module	5	143.97	\pm	1.20	141.00	147.67
Calculated cranial capacity	5	1235.76	\pm	24.24	1160.87	1300.50
Length-breadth index	5	75.15	土	0.70	72.73	76.88
Length-height index	7	71.36	\pm	1.68	64.64?	78.98
Breadth-height index	7	98.42	土	2.61	90.84	108.59
Length-auricular height index	4	61.00	\pm	0.58	59.54	62.14
Breadth-auricular height index	4	82.80	±	2.29	78.63	89.31
Tr. fronto-parietal index	6	70.07	±	2.00	65.79	79.39

Table 140 ${\bf ADULT\ \ FEMALE\ \ TYPE\ \ A_2}$ Mean Measurements and Indices of the Splanchnocranium

Characters		Mean ±	S.E.	Min.	Max.
Prosthion-basion line	5	92.40 ±	2.64	85.0	100.0
Nasion-prosthion line	5	59.80 ±	2.06	54.0	65.0
Nasion-gnathion line	1	96.00?		-	
Bizygomatic breadth	4	$121.50 \pm$	2.40	116.0	126.0
Nasal height	7	46.50 ±	1.72	40.0	54.0
Nasal breadth	6	$25.00 \pm$	0.86	22.0	28.0
Ant. inter-orbital breadth	7	$18.43 \pm$	0.92	15.0	22.0
Orbital breadth (right)	6	$39.42 \pm$	1.02	35.0	42.0
Orbital breadth (left)	5	38.40 ±	1.32	34.0	41.5
Orbital height (right)	6	$32.67 \pm$	0.73	30.0	35.0
Orbital height (left)	4	31.88 ±	0.83	30.0	34.0
Orbital height length	3	55.00 ±	3.22	49.0	
Maxillo-alveolar length	3	64.00 ±	0.29		60.03
Maxillo-alveolar breadth	3			6 3.5?	64.5

TABLE 140 - Continued

Characters	n	Mean \pm S	i.E.	Min.	Max.
			-1-20-1		
Palatal length	4	41.50 ± 1	.04	39.0	44.0
Palatal breadth	1	35.00?		48.71	
Superior facial index	3	51.36 ± 1	.38	75.00	53.36
Orbital index (right)	5	79.91 ± 1	.37	75.90	83.33
Orbital index (left)	4	82.56 ± 2	2.82	48.48	88.24
Nasal index	6	55.44 ± 2	2.23 1	05.83?	65.12
Maxillo-alveolar index	3	$117.21 ~\pm~ 7$	7.23	_	130.61
Palatal index	1	85.37?			_

TABLE 141 ADULT FEMALE TYPE \mathbf{A}_2

Mean Indices of the Whole Skull

Characters	n	Mean \pm S.E.	Min.	Max.
Tr. cranio-facial index Vert. cranio-facial index Long. cranio-facial index Jugo-frontal index	2 5 5 3	91.98 48.12 ± 1.61 52.90 ± 1.28 73.94 ± 2.02	88.55 43.20 49.13 70.63	95.42 53.28 56.82 77.59

TABLE 142 ADULT FEMALE TYPE \mathbf{A}_2 Mean Measurements and Indices of the Mandible

Characters	n	Mean	Min.	Max.
Ht. of mandibular ramus Max. breadth of mandibular ramus Min. breadth of mandibular ramus Ht. at mandibular symphysis Mandibular length Mandibular angle	2 2 2 1 2 2 2	53.75 40.25 32.00 21.00 84.50 123°	50.5 39.0 30.0 — 81.0 123°	57.0 41.5 34.0 — 88.0 123°

 ${\bf TABLE~143}$ ${\bf ADULT~FEMALE~TYPE~A}_2$ ${\bf Mean~Measurements~and~Indices~of~the~Permanent~Maxillary~Molar~Teeth}$

Characters	n	Mean	Min.	Max.
Mesiodistal crown diam. of M_1	2	10.88	10.75	11.00
Labiolingual crown diam. of M ₁	2	11.62	11.25	12.00
Mesiodistal crown diam. of M_2	2	9.75	9.50	10.00
Labiolingual crown diam. of M_2	2	11.00	11.00	11.00
Mesiodistal crow diam. of M_3	1	9.00	_	_
Labiolingual crown diam. of M_3	1	10.00	_	_
Crown index of M ₁	2	106.92	104.76	109.09
Crown index of M ₂	2	112.90	110.00	115.79
Crown index of M_3	1	111.11	_	-

* * =					
Characters	n	Mean	S. E.	Min.	Max.
Mesiodistal crown diam. of M_1	2	10.38		10.00	10.75
Labiolingual crown diam. of M_1	2	10.38	x.	9.75	11.00
Mesiodistal crown diam. of M ₂	3	9.83 ±	0.60	9.00	11.00
Labiolingual crown diam. of M_2	3	10.00 ±		8.50	12.00
Mesiodistal crown diam. of M ₃	3	9.33 ±	4	9.00	9.50
Labiolingual crown diam. of M ₃	3	9.83 ±		9.00	11.00
	2		0.00	90.80	110.00
Crown index of M ₁		100.40	F 07	89.72	109.09
Crown index of M ₂	3	$101.46 \pm$	- 10	94.74	122.22
Crown index of M ₃	3	$105.65 \pm$	8.43	04.14	144.22

TABLE 145

ADULT FEMALE COMBINED

Statistical Constants of the Measurements and Indices of the Neurocranium

Characters	n	Mean	±	S. E.	S. D.	c. V.	Min.	Max.
Max. cranial length Max. cranial breadth Nasion-inion length Basion-bregma height Min. frontal breadth	10 10 7 11 9	176.50 131.70 166.36 126.32 91.94	± ± ± ± ±	1.03 2.19 1.26 2.33 1.84	3.24 6.93 3.35 7.73 5.53	1.84 5.26 2!01 6.12 6.01	173.00 119.00 162.00 117.00 85.00	182.00 147.00 173.00 139.00 104.00

TABLE 145 - Continued

Characters	n	Mean	\pm	S. E.	S. D.	c. v.	Min.	Max.
West's 1								
Vertical porion height	7,	107.86	\pm	2.06	5.47	5.07	100.00	117.00
Median sagittal arc	8	355.88	\pm	2.94	8.31	2.34	344.00	
Vert. transversal arc	8	289.69	\pm	2.78	7.86			371.00
Horizontal circumference	10	501.30	土	8.22	25.97	2.71	282.00	304.00
Cranial module	7	145.05	<u>+</u>	1.31		5.18	483.00	573.00
Calculated cranial capacity	9	1235.45	+	25.72	3.48	2.40	141.00	150.67
Length-breadth index	8	75.80	+		68.15	5.52	1145.03	1324.35
Length-height index	9	77.05	_	1.29	3.64	4.80	72.47?	84.00
Breadth-height index	9		±	1.48	4.43	5.75	64.64?	78.98
Length-auricular height index	6	97.74	土	2.19	6.57	6.72	90.70?	108.59
		60.20	\pm	88.0	2.16	3.59	56.18	62.14
Breadth-auricular height index	6	81.82	土	1.69	4.13	5.05	77.52?	
Tr. fronto-parietal index	8	70.26	\pm	1.47	4.17	5.94		89.31
		- 1				5.94	65.79	79.39

Table 146 ADULT FEMALE COMBINED Statistical Constants of the Measurements and Indices of the Splanchnocranium

								
Characters	n n	Mean	±	S. E.	S. D.	c. v.	Min.	Max.
Prosthion-basion line Nasion-prosthion line Nasion-gnathion line Bizygomatic breadth Nasal height Nasal breadth Ant. inter-orbital breadth	6 6 1 4 8 8	93.75 59.33 96.00? 121.50 46.19 24.88 18.61	+ + + + + +	2.54 1.75 2.40 1.52 2.08	6.23 4.29 — 4.80 4.31 5.90	6.65 7.23 — 3.95 9.33 23.71	Min. 85.00 54.00 — 116.00 40.00? 22.00	100.50 65.00 — 126.00 54.00 28.00
Orbital breadth (right) Orbital breadth (left)	8 7 7	39.31 38.93 32.71	_ ± ± ±	0.73 0.75 1.01	2.18 2.12 2.67	11.71 5.39 6.86	15.00 35.00 34.00	22.00 42.00 41.50
Orbital height (right) Orbital height (left) Maxillo-alveoler length	5 4 4	31.50 55.50 64.00	± ± ±	0.62 0.74 2.33 0.20	1.63 1.66 4.66	4.98 5.27 8.40	30.00 3 0 .00 49.00	35.00 34.00 60.00?
Maxillo-alveolar breadth Palatal length Palatal breadth Superior facial index	4 1 3	41.50 35.00 51.36	土	1.04	0.41 2.08 — 2.39	0.64 5.01 —	63.50? 39.00 —	64.50 44.00
Orbital index (right) Orbital index (left) Nasal index	6 5 7	80.70 80.50 55.15	± ± ±	1.37 3.00 1.91	3.35 6.71 5.05	4.65 4.15 8.34 9.16	48.71 75.00 72.29 53.33	53.36 84.62 88.24 65.12
Maxillo-alveolar index Palatal index	4	115.98 85.37	土	5.26	10.51	9.06	105.83?	130.61

TABLE 147
ADULT FEMALE COMBINED

Statistical Constants of the Mean Indices of the Whole Skull

Characters	n	Mean	\pm	S. E.	S. D.	c. v.	Min.	Max.	
Tr. cranio-facial index	2	91.98			_	_	88.55	95.42	
Vert. cranio-facial index	6	48.22	\pm	1.32	3.23	6.70	43.20	53.28	
Long. cranio-facial index	. 6	53.49	土	1.33	3.27	6.11	49.13	56.82	
Jugo-frontal index	3	73.94	\pm	2.02	3.49	4.72	70.68	77.59	

TABLE 148

ADULT FEMALE COMBINED

Statistical Constants of the Measurements and Indices of the Mandible

Characters	n	Mean	\pm	S. E.	S. D.	C. V.	Min.	Max.
Bigonial breadth	1	86.00					_	
Ht. of mandibular ramus	3	51.50	\pm	2.93	5.07	9.84	47.00	57.00
Max. breadth of mandibular ramus	3	39.50	\pm	1.04	1.80	4.56	38.00	41.05
Min. breadth of mandibular ramus	4	32.00	\pm	1.16	2.31	7.22	30.00	34.00
Ht. at mandibular symphysis	3	23.33	土	1.20	2.08	8.92	21.00	25.00
Mandibular length	3	81.83	\pm	3.35	5.79	7.08	76.05	88.00
Mandibular angle	3	126.33	±	3.34	5.77	4.57	123°	133°

TABLE 149

ADULT FEMALE COMBINED

Mean Measurements and Indices of Permanent Maxillary Molar Teeth

Characters	n	Mean	Min.	Max.
Mesiodistal crown diam. of $\mathrm{M_1}$	2	10.88	10.75	11.00
Labiolingual crown diam, of M_1	2	11.62	11.25	12.00
Mesiodistal crown diam. of M_2	2	9.75	9.50	10.00
Labiolingual crown diam. of M_2	2	11.00	11.00	11.00
Mesiodistal crown diam. of M_3	1	9.00		
Labiolingual crown diam. of M3	1	10.00		Titlera
Crown index of M ₁	2	106.92	104.76	109.09
Crown index of M ₂	2	112.90	110.00	115.79
Crown index of M ₃	1	111.11	wa 1980 M	

TABLE 150 ADULT FEMALE COMBINED

Statistical Constants of the Measurements and Indices of the Permanent Mandibular Molar Teeth

Characters	n*	Mean	土	S. E.	S. D.	c. v.	Min.	Max.
Mesiodistal crown diam. of M_1	2	10.38			_	_	10.00	10.75
Labiolingual crown diam. of M_1	2	10.38				_	9.75	11.00
Mesiodistal crown diam, of M_2	3	9.83	\pm	0.60	1.04	10.58	9.00	11.00
Labiolingual crown diam. of M_2	3	10.00	\pm_{1}	1.04	1.80	18.00	8.50	12.00
Mesiodistal crown diam. of M_3	3	9.33	\pm	0.16	0.28	3.00	9.00	9.50
Labiolingual crown diam. of M_3	3	9.83	\pm	0.60	1.04	10.58	9.00	11.00
Crown index of $M_{\rm I}$	2	100.40			_	_	90.80	110.00
Crown index of M ₂	3	101.46	±	5.97	10.32	10.17	89.72	109.09
Crown index of M_3	3	105.65	土	8.43	14.59	13.81	94.74	122.22
					- 1.00	10.01	94.74	122.22

^{*} Althrough n denotes number of skulls (in case of tooth)

In sum, the population of mature Harappan culture (Cemetery R 37) was long-headed of which one type was tall, rugged and sturdy-built, having pronounced eye-brow ridges, receding forehead, broad nose with depressed root (Type A); and the other was gracile, comparatively shorter, finer and weaker (Type A_1).

In a narrow trench at Area G some round-headed crania (Type B_1) were found huddled together with the long-headed gracile type (Type A_1).

At Cemetery H Stratum II, skeletons similar to mature Harappan culture (Cemetery R 37) were discovered (Type A and Type A_1) in addition to a rather tall, large- and round- headed type (Type B_2). In Stratum I, besides the long-headed and round-headed peoples (Type A, Type A_1 and Type B_2), another rather medium statured small- and medium-headed, low-faced people (Type A_2 ; females only) was found.

GENERAL OBSERVATIONS

We have so long described the skulls according to the traditional way of breaking up samples into morphological types. We must, however, remember that the majority of the samples come from a population living within a restricted area of a large and populous city (ante, page 10) where some intra-breeding was presumably involved. Under these circumstances the types as based upon a combination of characters should not be taken to be identical with "races" in the modern sense of that term. Obviously, in any sample of skulls, dimensions, indices and observations will vary to some extent. Variation will be greater, the larger the sample. We have, therefore, also treated the material in the present section in a way that takes into account this variability, by calculating mean, standard deviation and standard error of the characters of samples from different strata of Harappa. It is likely that some interesting results hitherto unnoticed may become noticeable in this way.

The mean values of stature, head length, head breadth, head height, length-breadth index, length-auricular height index, superior facial height, facial breadth, superior facial index, nasal height, nasal breadth, nasal index, cranial module and cranial capacity for the male skeletons have been given in Table 151.

It is observed that the estimated statures as calculated from long bones coming from different burial sites indicate that the people of R 37 were somewhat taller than those of H.

Skulls from R 37 are distinguishable from those of H by the comparatively smaller cranial capacity of the former. The circumferences of both are however not very different. That would indicate that there is some difference in height.

TABLE 151

AVERAGE VALUES OF MEASUREMENTS AND INDICES OF MALE CRANIA FROM CEMETERY R 37 AND CEMETERY H

	HEAD LENGTH	HEAD BREADTH	HEAD HEIGHT	HOR.	CRANIAL CAPACITY IN CC.	L-B INDEX	L-AH INDEX	B-AH INDEX	UPPER FACE HEIGHT	BIZYGO- MATIC BREADTH	SUPERIOR FACE INDEX		NASAL BREADTH	NASAL INDEX	STATURE
Cemetery H St. I	187.33 ±5.46	138.00 ±1.73	118.50	523.67 ±11.88	1470.89	73.83 ±2.83	62.14	85.90	65.50	133.50	49.16	49.00	27.00	55.20	_
St. II	189.10 ±1.58	144.67 ±4.26	119.00	530.33 ±8.22	1505.19 ±48.77	76.72 ±1.29	64.85	85,61	70.30 ± 1.56	136.00	50.00	52.67 ± 1.10	25.50 ± 1.03	47.64 ± 1.30	1717.57 ± 13.10
Cemetery R 3	7 187.54 ±1.09	133.32 ± 0.83	115.00 ±0.84	520.00 ± 2.99	1387.57 ± 11.53	71.05 ±0.65	61.37 ±0.68	86.26 ±0.74	70.62 ±1.20	131.25 ± 2.64		51.96 <u>+</u> 0.90	26.68 ±0.74	51.08 ±1.42	1757.10 ± 23.60

Moreover, certain other distinctions are also noticeable. While the absolute lengths of the skulls of R 37 are not very different from those of H, the head is distinctly narrower, height is less, and the upper face is slightly narrower. The cranial index of R 37 is lower (71.05) than in the case of H (76.72; 73.83). This implies that people of R 37 were more long-headed and had narrower faces than those represented in Cemetery H.

We shall now consider the skulls found in only Cemetery H. If we discard the female skulls and limit ourselves only to the male skulls a slight difference is noticeable between Stratum I and II. For example, skulls from Stratum I are smaller in size and have higher nasal index than those of Stratum II. But what is significant is that a few round-headed skulls are present in both the strata, while this element is so far completely absent from Cemetery R 37. From this point of view the population of R 37 appears to be more homogenous than those represented in Cemetery H.

One of the major questions which arises from the above observations is this: Were the people of R 37 and of H genetically related to one another or not? Did the people of H replace the earlier population of R 37, or do they represent a somewhat differentiated group which lived socially separated from the rest and thus probably maintained their difference till later times? As far as the present evidence goes, it is not possible to answer either of the two questions satisfactorily. There is no positive evidence to suggest that the population of R 37 was subjected to heavy intermixture which might have brought about drastic changes in their cranial characteristics. This is in spite of the fact that between R 37 and H a very sharp change is noticeable in regard to culture.

Harappan culture was fairly widespread in north and north-western India. If samples were available from all the discovered sites it would have given us some clue about the regional differences of the population in that quarter of India in the past; but the available skulls are mainly from Rupar, Harappa, Mohenjo-daro and Lothal. It is interesting that the Harappan population of Mohenjo-daro has similarities with present-day long-headed population of Sind, so far as the cephalic index is concerned. The same relationship holds between the ancient population of Harappa and the present-day population of the Punjab. The skulls from Lothal (Sarkar, unpublished) are on the average round-headed (79.68; value calculated by us on seven better preserved skulls). This is remarkably close to the cephalic index of the present-day inhabitants of Gujarat.*

When we consider these facts it appears striking that the differentiation in regard to cephalic index observable in the Harappan age is so remarkably similar to the differentiation in the present-day population in the corresponding areas. In other words, the implication is that the population in this widespread region of India has remained more or less stable since Harappan times to the present day.


^{*} The occurrence of round-head people at Lothal in the Harappan age cuts seriously across Riskey's hypothesis that the round-headedness of Gujarat or Maharashtra was due to the Scythian invasion of India during the 1st and 2nd Century A.D.

This may or may not be borne out by fresh discoveries, or if other physical characters are taken into account; but the above significant similarity in cephalic index cannot be overlooked in any case.

Such stability is perhaps comparable only to what is observed in connection with the population of Egypt which seems to have persisted over four or five millenia or more in the same manner as in the case of Harappa. In other parts of the world, marked changes of culture have sometimes been the result of invasion by people bearing other physical characteristics, so that cultural change has been accompanied by changes in physical characters also. But in the case of Harappa, although cultural change is marked, yet, it does not seem to have been the result of large-scale invasion; the physical types remaining, more or less, constant. We can, therefore, suggest that even if there were invasion, the number of invaders must have been too small to bring about marked changes in physical characters, or that most of the changes were due to cultural forces of internal origin.

Acknowledgments: We wish to acknowledge our indebtedness to Dr D. K. Sen, Head of the Department of Anthropology, University of Lucknow for many profitable discussions and to Dr S. S. Sarkar, F.N.I., of the Department of Anthropology, University of Calcutta for advice and for allowing us to utilize his unpublished data on Lothal skeletons.

Thanks are due to Mrs Anima Roy, Mr A. Pal and Miss Bela Sinha for their co-operation in the laboratory; to Dr G. S. Mody for helping us in determination of age of the skeletons and in preparing the list of skeletal materials, and to Mr H. Mukherjee for his assistance. Finally, we also express our thanks to Mr A. K. Chakrabarti, In-charge of Publication, Anthropological Survey of India, for his keen interest in bringing out the present volume expeditiously.




Frontal view



Vertical view



Left lateral view



Occipital view

ŠKL. H 695 : ČEMĒTĒRY H ST. 11 (ÕPEN BURIALS)



Skl. H 698



Fig. 1 Occlusal aspect

Fig. 2 Left lateral aspect



Fig. 3 Occlusal aspect




Fig. 4 Left lateral aspect

Occlusal and Left lateral aspect of Mandible : Cemetery H St. II (Open Burials)



Fig. 1 Flexor aspect of Humerus $\left[\text{ H } 502(\text{G}) \right]$ a, right; b, left



Fig. 2 Extensor aspect of Humerus [H 502(G)] a, right; b, left



Fig. 3 Flexor aspect of Humerus (H710) a, right; b, left




Fig. 4 Extensor aspect of Humerus (Hi710) a. right: b. left

Flexor and Extensor aspect of Humerus : Cemetery H St. II (Open Burlats)



Frontal view



Vertical view



Right lateral view



 $Occipital\ view$

SKL. H 247 (b) : CEMETERY H ST. I (JAR BURIALS)




Fig. 1 Occlusal aspect




Fig. 2 Left lateral aspect



Fig. 3 Occlusal aspect




Fig. 4 Left lateral aspect

Occlusal and Left lateral aspect of Mandible : Cemetery H St. I (Jar Burials)

Skl. H 206 (d)

Skl. H 255(a)



Fig. 1 Infant in embryonic position : Jar burials




Fig. 2 Occipital view showing cranial injury (Skl. H 698) : Open burlals



Fig. 3 Endocraniar cast of Skl. H 5440 (D): Mound AB




Fig. 1 Flexor aspect of Humerus | Skl. H 206 (d) | (Right) : Jan burials




Fig. 3 Volar aspect of articulated Radius & Ulua (Skl. H 710) : Open burials a, left; b, right



Fig. 2 Radial aspect of Ulna [Skl. H 206 (d)] : Jar burials a, left; b, right




Fig. 4 Volar aspect of articulated Radius & Ulna (Skl. H 698): Open burials a, left; b, right




Fig. 1 Flexor aspect of Humerus: R 37 a, Skl. 10 (rt.); b. H 796 (B) (rt.); c, H 818 (lt.)



Fig. 2 Anterior aspect of Fumer: Open burials a. H 698 (left); b. H 710 (left)




Fig. 3 Anterior View of Tibia: Open burials a H 710; b, H 698



Fig. 4 Anterior view of Tibia & Fibula in articulation (H 710): Open burials



Fig. 1 Hand (H 184 k)



Fig. 2 Foot [306 (a)]

REFERENCES

Boyd, J. D. and J. C. Trevor	1953 Race, Sex, Age and Stature from Skeletal Material. In: Modern Trends in Forensic Medicine, edited by Keith Simpson, pp. 133-152. London, Butterworth.
Buxton, L. H. Dudley and D. Talbot Rice	1931 Report on the Human Remains found at Kish. J. Roy. Anth. Inst., Vol. LXI, pp. 57-119.
Coon, Carleton S.	1939 The Races of Europe. New York, Macmillan.
Dixon, Roland B.	1923 Racial History of Man. New York. Charles Scribner's.
Dupertuis, C. Wesley and John A. Hadden, Jr.	1951 On the Reconstruction of Stature from Long Bones. Amer. J. Phys. Anthrop., N. S. Vol. 9, pp. 15-54.
Dwight, Thomas	1894 Methods of estimating the height from parts of the skeleton. Med. Rec., N. Y., Vol. 65, p. 16.
Frederichs, Heinz F. and Heinrich W. Muller	1933 Die Rassenelemente im Indus. Anthropos, Band XXVIII, pp. 383-406.
Guha, B. S. and P. C. Basu	1938 Report on the Human Remains Excavated at Mohenjo-daro in 1928-29. In: Further Excavations at Mohenjo-daro, by E. H. Mackay, Vol. I, pp. 613-638. Delhi, Govt. of India Press.
Hadden, A. C.	1919 The Wanderings of Peoples. First ed., 1911, Reprinted (with corrections) 1912, 1919. Cambridge, Univ. Press.
Hargreaves, H.	1935 Excavations in Baluchistan, Mem. Arch. Surv. of India, No. 35. Delhi, Govt. of India Press.
Hooton, Earnest Albert	1946 Up from the Ape. New York, Macmillan.
Kappers, C. U. Ariëns	1934 An Introduction to the Anthropology of the Near East in Ancient and Recent Times, with a chapter on Near Eastern Blood groups by Leiand W. Parr. Amsterdam, N. V. Noord-Hollandsche Uitgeversmaatschappij.
Keith, Sir Arthur	Review of Mohenjo-daro and the Indus Civilization', Hlust. London News, Dec. 19, p. 1002.

Krogman, Wilton Marion	1937	••	Cranial types from Alisar Hüyük and their relations to other racial types, ancient and modern, of Europe and Western Asia. In: The Alishar Hüyük, seasons of 1930-32, by Hans Henning von der Osten, Pt. III, (OIP XXX), pp. 213-293. Chicago, Chicago Univ. Press.
	1940	••	Racial types from Tepe Hissar, Iran, from the Late Fifth to the Early Second Millennium, B.C. Kon. Ned. Akad. Wet., Verh. (Tweede Sectie), Dl. XXXIX, No. 2, pp. 1-87, Amsterdam.
	1946	••	Proc. Inst. Med. Chicago, Vol. 16, p. 154
Lee, Alice and Karl Pearson	1901		A first study of the Correlation of the Human Skull. Phil. Trans. Roy. Soc. London, Series A, Vol. cxcvi, pp. 225-264.
Manouvrier, L.	1893	••	La dètermination de la taille après les grandes os des membres. Mem. de la Soc. d' Anthropol. <i>de Paris</i> , 2 sér, Tome 4, pp. 347-402.
Martin, Rudolf	1928		Lehrbuch der Anthropologie. Jena, Gustav Fischer Verlag.
Martin, Rudolf and Karl Saller	1956		Lehrbuch der Anthropologie. 3 Leif. Stuttgart, Gustav Fischer Verlag.
Møllisøn, Theodor	1938		Spezielle Methoden anthropologischer Messung. In: Handbuch der biologischen Arbeitsmethoden , edited by Emil Abderhalden, Abt. VII, Lfg. 468, pp. 523-682. Berlin & Wien, Urban & Schwarzenberg.
Mukherjee, Ramkrishna,			
C. Radhakrishna Rao and J. C. Trevor	1955	٠.	The Ancient Inhabitants of Jebel Moya (Sudan). Cambridge, Univ. Press.
Pearson, Karl	1898	* *	Mathematical Contributions to the Theory of Evolution, V: On the Reconstruction of the Stature of Prehistoric Races. Phil. Trans. Roy. Sec. London, Series A, Vol. excii, pp. 169-244.
Sergi, G.	1899		Specie e varieta umane. Riv. di Sci. Biol. I, Part I, p. 20.
Sewell, Col. R. B. Seymour and B. S. Guha	1931	* *	Human Remains. In: Mohenjo-daro and Indus Civilization, edited by Sir John Marshall, pp. 599-648. Lendon, Arthur Probsthan.
Vats, Madho Sarup	1940	* *	Excavations at Harappa. Vol. I. Delhi, Govt. of India Press, Calcutta.
Wheeler, R. E. M.	1947		
Wilder, Darris H.	1920		A Laboratory Manual of Anthropometry. Philadelphia, Blakiston.

APPENDIX

CATALOGUE OF SKELETAL REMAINS

CEMETERY R 37

Skl. H 779. Adult; male. Rib fragments; left humerus in pieces; right ischium, united apophysis; left femur fragments; proximal and distal pieces of left tibia, shaft of left fibula.

Skl. H 779*. Adult; female. Body fragments of cervical vertebrae; carpal left 3rd, metacarpal left 3rd; proximal

and distal parts of left tibia.

3 Skl. H 779 (a). Adult 25-30 years; female. Skull incom plete; fragments of thoracic and lumbar vertebrae; clavicle represented by medial two-thirds; right and left shaft fragments of humerus; distal portion of right ulna; distal end of right radius; fragments of pelvis; distal ends of right tibia and right fibula; talus right and left; calcaneum right, left medial half fused apophysis; left navicular; left cuboid; cuneiform right 1st, left 1st and 2nd; metatarsals right 1st, 2nd, 4th, 5th, left 1st and

2nd, Extra femora pieces. Skl. H 779 (b). Juvenile; female. Decayed skull presented by broken part of face and left temporal part, left half of mandible; fragments of cervical vertebrae.

Skl. H 779 (c). Adult, about 30-35 years; female. Skull incomplete but restored; head and axillary border of scapula. right

Adult; female. Skull incomplete and Skl. H 779 (d). disintegrated.

Skl. H 779 (e). Adult, about 25-30 years; male. Nearly

complete skull. Skl. H 779 (f). Adult; female. Skull fragments represented by facial part and broken part of temporal bone with

zygomatic arch. Skl. H 780. Adult, 30-40 years; female. Skull complete; rib fragments; proximal part of left ulna; right hip-bone broken; right femur slightly twisted distaly and medialy; distal parts of left fibula; right calcaneum, right 1st metatarsal, proximal phalanx 1.

metatarsai, product product skil H 781. Adult, above 25 years; male. Skull incomplete. Skil. H 781. Adult, above 25 years; male. Skull incomplete, broken piece of mandible; thoracic 10th, 11th, 12th and lumbar vertebrae 1st to 5th; left humerus (distal end missing); sacrum in one mass; fragments of pelvis and

Skl. H 782. Adult; female. Disintegrated skull fragents; typical cervical vertebrae 5, thoracic vertebrae 3; proximal piece of sacrum; rib fragments; head of right proximal piece of Sacrum; rib fragments; nead of right scapula; clavicle middle third and lateral third (left); shaft fragments of right humerus; right shaft fragments of ulna; right radius fragments 4, distal end of left radius; right metacarpals 1st, 2nd, and left 3rd; proximal radius; (hand) 1; fragments of picht illum; shaft and radius; right include paid 13t, 2nd, and left ord; proximat phalanx (hand) 1; fragments of right ilium; shaft and phalanx (hand) 1, tragments of right illum; shaft and condylar fragments of right and left femur; broken parts of right and left tibia; distal end of left fibula; left talus, of right and left tipla; distal end of left fibula; left talus, left calcaneum, navicular 2, cuneiform 4, cubeid 2, metatarsal 11, proximal phalange (foot) 4.

Skl. H 784A. Adult, about 20 years; male. Skull represented by parietal pieces, broken mandible, fourteen receibbles teeth have erupted but some of them are

represented by Parison pieces, proken mandible, fourteen mandibular teeth have erupted but some of them are missing from the socket and M3 did not cut.

missing from the specket and M 3 did not cut.

Skl. H 788. Adult, 25-30 years; female, Skull compressed side to side, right half of the mandible with all the 8 side to side; cervical vertebrae 6, thoracic vertebrae 10, teeth intact; cervical vertebrae 4. teeth intact, terminate of the interest of the intact, terminate of the intact, terminate of the intact, terminate of the intact, the interest of the intact, the intact, the interest of lumbar vertesiae (spanish and acromion processes of scapula; medial twocoracoid and actions processes of Scapula; mediat two-thirds of right clavicle; humerus right, humerus left thirds of right staticle, numerus right, humerus lett preximal two-thirds and distal pieces; left ulna 3 pieces; right radius, left radius shaft pieces; carpals right, lunate right radius, left 1st and 3rd, phalanges proximal 9, in-2. metacarpai lett bone right pieces, es pubis right, frag-termediate 2; hip-bone; femur right head and neck, shaft ments of left hip-bone; femur left head and neck, shaft ments of left hip-bene, remur right head and neck, shart and distal end, femur left head and medical condyle; patella left; tibla right pieces, tibla left proximal piece; fibula right and left proximal; navicular right and left

(one extra left); talus right and left, cuboid right piece; cuneiform right and left 1st & 3rd; calcaneum pieces, left sustentaculum tali; metatarsal right 5th, left 1st, 2nd, 4th and 5th fragments; phalange proximal 5,

Skl. H 790. Adult; female. Proximal part of left femur broken right talus.

15 Skl. H 790 (a). Adult; probably male. Fragment of right ulna; fragments of right hip-bone; shaft fragments of right and left femur; fragments of right and left tibia; distal end of right fibula, nearly complete left fibula (without head); right and left talus; matatarsal 1st. Skl. H 791. Adult, about 25 years; female. Left half of

mandible with lateral incisor, canine, PM 1, M 1, M 2 in the sockets; 1st to 7th carvical vertebrae, 1st, 2nd, and 3rd thoracic vertebrae in one mass, and 3 typical and 11th; 1st to 5th lumbar vertebrae with sacrum piece; broken parts of right and left clavicle; distal end and head of left humerus; left ulna fragments; fragments of right and left radius; broken parts of pelvis; shaft pieces of left femur; shaft fragments of right and left tibia; shaft fragment of left fibula.

Skl. H 791/A. Adult, 25 to 30 years; female. Skull nearly complete; right ulna fragments.

Skl. H 791 (b). Adult; sex(?). Skull fragments with broken teeth sticking.

Skl. H 792. Adult; female. Complete right ulna; distal half of right radius, proximal two-thirds of left radius. Skl. H 793. Adult, 25-30 years; male. Skull not well preserved and had to be restored before measurement; preserved and had to be restored before headstrement, cervical vertebrae in one mass; thoracic vertebrae 1st, 2nd, 3rd in one mass and 2 typical rib fragments; 2 lumbar vertebrae typical in one mass; fragment of right sacrum having osteophytic growth at superior surface; right scapula with head-spine and axillary border piece, fragments of left scapula; fragment of right elavicle; shaft of right humerus, shaft fragments of left humerus; right and left ulna fragments; right and left radius fragments; fragment of pelvis; shaft fragments of right femur, preximal two-thirds of left femur; right and left patella; right and left tibia; eroded right fibula, shaft fragments of left fibula; right and left talus; left calcanium; right navicular.

Skl. H 793 (A). Adult, 30-35 years; male. Skull nearly complete and had to be reconstructed; typical cervical vertebrae 6; 1st, 2nd, 5th, 6th and 7th thoracic vetebrae; lumbar vetebra 1, and 4th and 5th lumbar vertebrae in one mass with sacrum fragment; rib fragments; right scapula; medial third of right clavicle, middle one-third of left clavicle; proximal helf of right humerus; shaft fragment of right and left ulha; distal ends of right and left radius; carpal 3, metacarpal 7, and phalange 10; right lilum maca left right recent humans. left radius: carpal 3, mefacarpal 7, and phalange 10: right lilum piece, left publs piece; broken part of right and left femur; shaft of right tibia, proximal part of left tibia;

shaft of left fibula; phalange (foot) 4.

Skl. H 793 A*. Adult; female. Represented by isolated molar tooth; scapula fragments; medial third of right clavicle; head and distal end of left humerus; distal threefourth of right radius, shaft fragments of left radius; phalange proximal 2 and intermediate 1; fragments of left pelvis; right and left fibula; phalange (feet) preximal 1

and intermediate 1.

Skl. H 793 (B). Adult, 25-30 years; male. Skull complete. Skl. H 794, Adult, 25-30 years; male, Skull complete but not well preserved, mandible present; cervical 6, theracic 6th to 12th, and lumbar vertebbae 3rd to 5th; fragments 6th to 12th, and lumbar vertemae and to out; fragments of sacrum; rib fragments; broken parts of right and left scapula; broken parts of right and left calvicle; preximal hand distal end of right humerus; shaft of right ulna distal third of left ulna; fragments of right radius; carpai 3, phalanx 1; fragments of pelvis; right femur with patella, 3, phalaint t, seeman of right tibia, preximal two-third of left tibia; fibula fragments; number of foot bones.

^{*}Asterik mark indicates presence of second skeleton bear ing the same field number.

sphenoid fragments and portion of right maxilla with all the teeth of right side excepting M $_3$; few cervical vertebrae, fragment of a thoracic vertebrae; spine piece of left scapula; lateral two-thirds of left clavicle; fragments of right and left humerus; shaft of left ulna; distal three-fourth of right radius, shaft of left radius: fragments of pelvis; shaft fragments of left and femora; shaft fragments of left and right tibae; shaft frag-ments of left and right fibulae. Skl. H 795. Adult; sex (?). Medial two-third of left

Skl. H 795. Adult; sex (?). Medial two-third of left clavicle; shaft of right radius; shaft fragments of right

tibia and fibula fragments.

Skl. H 795/A. Adult, about 21-25 years; female. Skull not well preserved, mandible broken and had to be restored; mib fragments; right scapula; shaft fragments of left and right humerus; proximal and distal three-fourth of right ulna; distal end of right radius; phalange 4; broken shaft parts of right and left femur; right and left tibia fragments; distal end of right fibula.

Skl. H 795 A 2. Adult (?); female. Shaft and crumbled condyle of femur.

Skl. H 796. Adult; female. Typical vertebrae 2, disintegrated left hip-bone, shaft fragments of femora, shaft of Skl. H 796 left tibia.

Skl. H 796 A. Adult, above 25 years; male. Lumbar vertebrae 4, right scapula; left clavicle; proximal three-Skl. H 796 A. 25 years; male. Lumbar fourth of right ulna; shaft fragments of left ulna; shaft fragments of right and left radius; fragments of pelvis; proximal piece of femur; shaft of right fibula.

Skl. H 796 (B). Adult; male. Skull and mandible in locked-jaw condition; cervical vertebrae 3, thoracic vertebra 1st; rib fragments; fragments of scapulae; right clavicle; right humerus; right 3rd metacarpal.

Skl. H 797. Above 18 years; female. Left femur frag-

33 Skl. H 798 (i). Adult; male. Represented by few skull pieces; right half of the mandible; right ulna shaft; right metacarpal 3; proximal phalange (hand) 4; broken parts of right and left innominate; fragments of left femur; right femur head in acetabulum; left tibia; fibula fragments.

- Skl. H 798 (ii). Skl. H 798 (ii). Adult; female. Represented by skull fragments; broken mandible; 7th cervical vertebra, all thoracic and one lumbar; right clavicle; rib fragments; right humerus; right ulna fragments; right and left radius fragments; metacarpal left third; phalange (hand) intermediate 4 and distal 1; broken left femur; shaft of right tibia; proximal half of left tibia; shaft fragments of left
- 35 Skl. H 798 (iii). Adult; sex (?). Thoracic vertebra 1: lumbar vertebra 1; broken sacrum; shaft fragment of left shaft pieces of humerus; right ulna broken; proximal half of left tibia; right 3rd metatarsal; fragments of right femur shaft.
- Skl. H 798 (a). Adult, about 30 to 40 years; probably female. Skull complete; fragments of left pelvis; right and
- Skl. H 798 (A). Adult; male. Skull complete; rib fragments; cervical vertebrae 1st to 7th; theracic vertebrae 1st to 10th and 12th; lumbar vertebrae 1st-3rd and 5th; ribs right 1st and 2nd; body of the sacrum; distal three-fourth of the left humerus; right hip-bone having fused crest epiphysis and ischial apophysis 7 complete left hip-bone; and ischial apophysis apophysis and ischial apophysis apophysis and ischial apophysis apophysis apophysis and ischial apophysis apoph bone; right femur shaft fragments and left chips; shaft
- of left tibia. Skl. H 798/A1. Adult, about 30 years; female. Skull complete.
- 9kl. H 798/A2. Juvenile; female. Skull complete. 11. H 788/A3. About 21 years; probably male. Mundible 40 token in two halves; fourteen intact mandibular toth,
- did not out H 798 (B). Acult/ between 25 and 30 years; male dail incomplete; greater portion of the facial region and ranii missing; els fragments 2; phalanx 1.

 798 (C). Adult, but oen 25 and 30 years; male. 41
- 12
- 43 Skl. 10 d. Adult, above 65 years; male. Skull Skl. 10 defective.
 repr. 6. Adult. below 30 years; female. Skull frontal bone only. " Couplete.

- 45 Skl. H 798 f. Adult, 25-30 years; probably male. Skull represented by few pieces.
- Skl. H 798 G. Adult; male. Skull represented by frontal bone mixed up with lump of earth, superciliary arches well developed.
- Skl. H 799. Adult, above 25 years; sex (?). Few fragmentary pieces of thoracic and lumbar vertebrae; right scapula in pieces; right humerus fragments; broken part of ulna; olecranon piece of ulna attached with trochlear surface in hyperextention; fragments of radius; shaft of left tibla; distal three-fourth of right fibula; shaft of left fibula; right talus; phalanges.
- Skl. H 800. About 18 years; male. Medial-third of costoclavicular ligament of clavicle raised; fragments of left ulna; proximal phalanx (hand) 1; shaft and condylar fragments of right and left femur; proximal end of right and left tibia; fragments of right fibula; broken talus; cuneiform 3rd; metatarsal 3; proximal phalange 2.

Skl. H 801. Adult, about 25 years; probably female. Vertebrae cervical 5th to 7th, thoracic 1st, 12th and typical 3, lumbar 1st to 5th; left hip-bone; femur right shaft, and lateral condyle intact; broken shafts of right and left fibula.

- Skl. H 801A. Adult, 30 to 35 years; female. complete; mandible broken in 3 pieces; thoracic vertebrae 2: rib fragments; fragmentary pieces of right and left scapula; fragments of right and left clavicle; trochlear fragments of right humerus; distal third of left humerus; fragments of right ulna; fragments of right radius; carpal 2; metacarpal 6; phalange (hand) 6; left ilium fragments; proximal end of right femur; fragmentary left femur; right and left patella; proximal half of right tibia and proximal three-fourth of left tibia; shaft fragments of right and left fibula.
- Skl. H 801AI*. Adult; sex (?). Cervical vertebrae 1st to 5th in one lump; rib fragments; shaft fragments of left humerus; shaft fragments of right and left ulna; fragmentary shaft of left radius; shaft fragment of left femur; left fibula fragments.

- Skl. H 801A2. Adult; sex (?). Defective right humerus. Skl. H 801A3. Adult; female. Incomplete right hip-bone. Skl. H 801B. Adult; male. Skull bone represented by parietal fragments, petrous portion of left temporal and mastoid processes; distal half of right humerus; shaft of right femur; fragments of right tibia; distal piece of left fibula.
- Skl. H 801B*. Adult; female. Represented by skull frag-
- ments and shaft of right femur. Skl. H 801 b(l). Adult; sex (?). (Skeleton damaged by flood). Skull represented by parietal fragments and broken
- Portions of maxillae with few teeth in the socket.

 Skl. H 801 b(ii): Adult; female. (Severely damaged by flood). Skull represented by few parietal fragments, left temporal bone, and isolated molar tooth.

 Skl. H 801 b(iii): Adult; female. Benecontain.
- Skl. H 801 b(iii). Adult; female. Represented by small pieces of skull-bone.
- Skl, H 501 b(iv). Juvenile; male. Skull incomplete. lower facial part and base broken; mandible incomplete. left ascending ramus being broken; 14 mandibula teeth, M_3 did not erupt.
- Ski, H 802 (i). Adult, about 25 to 80 years; female. Skull badly warped; mandible in fragments; few vertebrae; rib fragments; fragments of right clavicle; shaft fragment of right and left humerus; hip-bone fragments broken; parts of left femur and broken shaft or right tibia.
- Sid. H 802 (ii). Adult; male. Skull represented by parietal and occipital fragments and isolated tooth, broken mandible; head and coracoid process of left scapula; shaft framents of right and left femur; shaft fragments of high tibia; distal end of right fibula and left fibula.

 Sid. N 802 (til). Adult; female. Right femur shaft; left fibula.
- figure with defective head.
- with defective head.

 No. 11 802. Adult; male. Shaft fragments of right and left tipla mixed up with animal bones.

 Skl. 18 802. Adult; female. Skull fragments represented by panicula and petrous portions of right temporal bone; left female; shafts of right and left tibla; distal portion of right think.

- 65 Sk!, H 802 A1. Adult, sex (?). Represented by axillary border of a scapula.
- 66 Skl. H 803. About, 25 years; male. Broken pieces of vertebrae; rib fragments; medial 2/3rd of right clavicle; right humerus (head broken); shaft pieces of left humerus; right and left ulna fragments; right and left radius shaft fragments; right carpal 8, metacarpal medial 4 and intermediate 2; left carpal 4, left metacarpal 4, proximal phalange 4; disintegrated right femur; shaft fragment of left femur; right and left patella; proximal end of right tibia; distal 2/3rd of left tibia with talus adherent (arthritis ankle joint); fragments of fibula and footbones.

Skl. H 803A. Adult; male. Skull represented by fragments of occipital bone and one lumbar vertebra.
 Skl. H 804. Adult; female. Skull incomplete. Other bones

- Skl. H 804. Adult; female. Skull incomplete. Ther bones include rib-fragments; right and left humeri fragments; left ulna; shaft fragments of right end left radius; few Wrist bones; right and left os coxae; shaft fragments of right femur; left femur; right and left patella; right tibia; fragments of right tibia with talus and calcanium; shaft fragments of right and left fibula; right cunciform 3rd, cuboid 1, and metatarsal 3.
- 69 Skl. H 804. Adult; male. Proximal 2/3rd of right femurand shaft of left radius.
- 70 Skl. H 805. Adult, fairly old in age; female. Skull not well preserved and had to be restored; 2nd cervical vertebra; part of right scapula; part of right clavicle; proximal 2/3rd of right humerous; distal ends of a radius; right tibia; shaft pieces of fibula; left talus; some hand and foot bones.
- Ski. If 805A. Adult, about 21-25 years; female. Skull represented by right frontal bone only. Other skeletal parts include cervical vertebrae 15th in one mass; theracic vertebrae 4th-8th in one mass and 9th and 10th in one mass; fragments of lumbar vertebrae; rib fragments; pieces of right scapula; lateral 2/3rd of right clavicle; shaft fragments of humerus; shaft fragments of right and left ulna; complete left radius; metacarpal 5; proximal phalange 3; shaft fragments of left (bula: navigular 1 and left cuboid.
- left fibula; navicular 1 and left cuboid.

 72 Skl. H 806. Adult; male. Skull with mandible complete; cervical vertebrae 5, rib fragments, acromion processes of right and left scapula; middle part of left clavicle; shaft fragments of humerus; proximal parts of right and left ulna; broken parts of left radius; right carpal bone 1, metacarpal 1, proximal phalange (hand) 3; fragments of peivis; shaft fragments of right femur; left femur with patella adherent to distal end; shaft fragments of right and left tibia; talus 1; calcaneum 1; metatarso-phalangeal
- joint intact; terminal phalanx (foot) 1.

 73 Skl. H 806A. Adult; female. Complete skull with mandible; several vertebrae; fragments of ribs; acromion and coracoid processes of scapulae; broken parts of clavicle; distal 3/4th of right humerus; complete left humerus; proximal 1/2 of right and left ulna; fragments of right radius; carpal 2, metacarpal 4, phalange 2; broken parts of right os coxae; shaft fragments of right and left femur; right and left patella; shaft fragments of tibiae; shaft fragments of left fibulae; left foot with intact cuneiform, cuboid, metators may left to be alanges.
- tarsal medial 4, phalanges.

 74 Skl. H 807. Adult; sex (?). Shaft of right humerus; distal algorithms of right ulna; proximal 1/4th of left radius; shaft fragments of femur; right tibia fragments and shaft fragments.
- ments of left tibia.

 75 Skl. H 808. Above 18 years; probably female. Rib fragments; coracoid process of scapula; shaft fragments of right humerus; shaft of left radius; carpal 3, metacarpal 2, phalange (hand) 8; broken piece of right hip-bone; fragments of femora; fragments of right tibia; right calcanuem 1, cuboid 1, right cuneiform 2; left navicular 1, left cuneiform 2, left cuboid 1; metatarsal 3, phalange 5.
- cuneiform 2, left cuses 1, sheaters 3, phalange 5.

 76 Skl. H 810. 17-21 years; probably female. Skull incomplete but preserves part of parietal, left zygoma, fragments of occiput, broken part of right manilla with 7 teeth (excepting M 3) intact; broken mandible showing all the teeth excepting M 3 which did not cut. Other skeletal parts consist of several vertebrae; rib pleces; head of scapula; nearly complete right humerus; shaft fragment of left humerus; ulna fragments; radius fragments; carpal 4.

- parts of right os conae; fragments of right femur, distal end of left femur, patella 2, shaft fragments of tibia, shaft fragments of fibula, metatarsals 7, calcaneum (right) 1, right talus 1, right cunciforms 3 and left talus.
- 77 Skl. H 811. Adult; male. Skull complete. Skeletal materials consist of broken left clavicle, right femur, fragments of fibula and first cervical vetebra.
- 78 Skl. H 811. Female adult, fragments of right humerus, shaft of right radius, fragments of pelvis, left scapula, right ulna fragments, left patella, shaft of left tibia and a few phalanges.
- 79 Skl. H 812. Adult; female. Skull complete and had to be restored. Other skeletal parts represented by odontoje process of second cervical vertebra, lumber vertebrae 3, broken part of sacrum, head and shaft pieces of right humerus, proximal 3/4th of left humerus and shaft fragments of right and left tibia.
- 80 Skl. H 813. Adult, about 18-21 years; female. Skull represented by temporal bone and left zygoma. Other skeletons consist of broken mandible with some of its teeth in position, number of vertebrae, body of sacrum, rib fragments, broken parts of both right and left scapulae, fragments of clavicle, nearly complete right humerus, distal part of left humerus, ulna fragments, shaft of right radius, proximal phalanx 1, distal phalanx 1, right ischium piece, shaft fragments of right and left femur, left patella, shaft of left tibia and shaft fragments of left fibula.
 81 Skl. H 814. Adult; female. Cervical vertebrae 5th, 6th and
- 81 Skl. H 814. Adult; female. Cervical vertebrae 5th, 6th and 7th in one mass; theracic vertebrae 3 in one mass and 2 in another; and lumber vertebrae 2 in one mass; many rib fragments, head and coraceid process of left scapula, complete right and left clavicle, broken right humerus, distalend of right radius, metacarpal (left) 1, proximal phalanx 1, es pubis, right patella, distalend of right tibia, fragments of left tibia, shaft fragments of fibula, cuneiform 1, metatarsal 1.
- metatarsal 1.

 82 Skl. H 814 a. Adult; male. Skull incomplete, represented by pertion of frontal bone, right yzgomatic arch and greater mart of the left parietal.
- part of the left parietal.

 83. Skl. H 814 b. Skull represented by few fragmentary pieces; no mandible is preserved; other skeletal parts include rib fragments, left scapula, middle piece of left clavicle, head of right humerus, shaft fragments of left humerus, left ulna fragments, fragments of left radius, fragmentary pieces of pelvis, fragments of left tibia, metatarsal 2 and left talus.
- metatarsal 2 she left takes.

 84 Skl. H 814 c. Adult: female. Nearly complete right os coxac.
- 85 Skl. H 815. Adult, about 20 years; male. Complete mandible with 14 teeth intact, M 3 on both sides did not crupt. Teeth show considerable wearing. High degree of wearing of incisors definitely suggest an edge to edge bite. Other skeletal parts are represented by fragments of left femurally.
- seniy.

 Ski. H 816. Skull nearly complete, skeletal materials consist of few vertebrae, head of right scapula, right and left clavicle, fragments of humerus, shaft fragments of both right and left ulna, shaft fragments of right radius, preximal phalanges (hand) 2, part of left hip-bone, shaft fragments of both right and left femur, shaft fragments of tibla and distal half of left fibula.
- ef tibla and distal half (18-21 years); female. Skull not Skl. H 817. Adult, about 18-21 years; female. Skull not well preserved—greater portion of the occiput and occipital well preserved—greater portion of the occiput and occipital part of the basis cranii had to be reconstructed. Mandible is present. Skeletal materials consist of rib fragments, few pieces of theracie and lumbar vertebrae, medial 2/3rd of right clavicle, shaft fragments of left humerus, left ulna fragments, shaft fragments of both radii, metatarsals 2 fragments of polvis, fragments of femora, fragments of tibliae, broken shaft of fibula and proximal phalanges (foot) 4.
- 88 Skl. H 818. Adult; male. Skull complete. Other skeletal materials include few vertebrae, sacrum piece with completely sacralized 5th lumbar vertebra, few ribs, broke, pieces of right and left scapula, fragments of clavicle, shaft of right humerus, complete left humerus, shaft of a radius, phalanges 2, fragments of peivis, fragments of right femur, right and left patella, complete left tible, shaft fragments of fibula, left calcaneum 1, left talus, left

- 104 Skl. 5. Female, about 18-20 years. Skull with mandible
 - incomplete and badly warped.
 Skl. 7. Adult, sex (?). Two typical cervical vertebrae and few rib fragments.

MEMOIR: No. 9

- Skl. 8. Adult; sex (?). Head of a right femur.
- 107 Skl. 9. Adult; sex (?). Metacarpal fragments and phalanges (hand).
- 108 Skl. 10. Adult; male. Skull complete with mandible. Other skeletons include several vertebrae, left side of a sacrum, broken ribs, acromion process of right scapula, acromion and coracoid processes of left scapula, complete right humerus, complete left ulna, distal end of left radius, carpals 6, metacarpals 5, phalanges 9, broken parts of right and left os coxae, right and left femora, both tibia, left fibula, left calcaneum, left tarsal, phalanges (foot) 5 and metatarsal 1.

cuboid 1, right talus, navicular 1, right metatarsal 1 and

- proximal phalanx (foot) 1. Skl. H 819. Juvenile; female. Skull represented by right sagital half without mastoid process. Other skeletal parts include right-half of mandible, few vertebra, 32 rib fragments, clavicle fragments, fragments of radius, matatarsal 1, metacarpal 1, proximal phalanges (hand) 3, shaft fragments of femur, shaft fragments of right tibia, broken shaft of left fibula, right and left talus, right navicular, right cuneiform 1, metatarsals 2, proximal phalanx (foot) 1 and distal phalanx (foot) 1.
- Skl. II 820. Adult; about 25 years. Skull represented by few fragments and isolated incisor tooth. Other skeletons are as follows; 2nd cervical vertebra, fragments of few thoracic and lumbar vertebrae, sacrum, many rib frag-ments, broken part of right scapula, complete left scapula parts of left clavicle, broken parts of humerus, shaft fragments of radius, shaft of an ulna, carpals 7, proximal phalanges 2, intermediate phalanges 2, broken parts of right and left os coxae (left os coxae having fused ischial tubresity apophysis), proximal half of left femur, shaft fragments of right femur, complete right and left patella. shaft fragments of both right and left tibia, shaft fragments of fibula, right calcaneum 1, right talus, right navicular, proximal phalanges 3 and cuneiforms 5.
- Skl. H 820 L. Adult; female. Complete right tibia. Skl. H 820 C. Adult; female. Nearly complete left ulna distal end missing.
- H 820 (i). Adult; female. Skull complete. No mandible.
- Skl. H 820 (ii). Adult; female. Skull incomplete, mandi-
- Skl. H 820 (iii). Adult; male. Skull nearly complete and well-preserved. Only portion of left frontal and parietal bone is missing.
- Skl. H 821. Adult; male. Skull fragments, several vertebrae, sacrum piece, sternum, rib fragments, left scapula, fragments of right clavicle, distal portion of right humerus, left humerus fragments 3, fragments of right ulna (belonging to two individuals), shaft fragments of right radius, carpals 2, phalanges 2, fragments of os coxae, shaft fragments of right and left femur, right and left patella, fragments of right tibia, nearly complete left tibia, fibula fragments 5, left calcaneum 1, sustentaculum tiali 1, right and left talus, right navicular, left cuboid, left cunelform 2, metatarsals 4, phalanges 4.
- Unnumbered skeleton. Juvenile. Represented by shaft fragments of right femur.
- Unnumbered skeleton. Juvenile. Shaft fragments of left femur 3.
- Unnumbered skeleton. Probably female adult. Skeleton severely damaged; skull represented by portion of parietal and right masteid process.
- Skl. 1. Adult; male. Skull and mandible complete. Other skeletons include number of vertebrae, broken parts of sacrum, few ribs, parts of scapula, fragments of right clavicle, complete right humerus, fragments of right ulna, shaft fragments of right radius, carpals 2, metacarpals 4. fragments of right pelvis, nearly complete femora, right patella, shaft fragments of tibia, shaft fragments of right and left fibula 7, calcaneum 2, right talus, right and left navicular, cuneiforms 3, metatarsals 5, phalanges and sesamoid bones 2.
- 101 Skl. 2. Adult: female. Skull complete but not well preserved. Other skeletal parts represented by several vertebrae, sacrum fragments 3, rib fragments, scapula fragments, part of right clavicle, shaft fragments of humerus right and left ulnae, fragments of radius, right carpal 1. fragments of metacarpais, left carpais 8, fifth metacarpai, Proximal phalanges (hand) 3, fragments of both right and preximal phalanges (hand) 3, fragments of both right and left os coxae, shafts of right and left femur, left patella, shaft fragments of tibiae, fragments of fibula, right and left talus, right cuboid, left navicular, cuneiforms 4, metatarsais 4, proximal phalanges (foot) 5.

 Ski. 3. Adult; sex (?). Skeleton broken parts of caiva, few vertebrae, rib fragments and chips of bones.

 Ski. 4. Adult, 60-85 years; male. Skull badly preserved representated by complete vault, left parietal, left temporal, portion of occiput and small portion of facial bones.
- ral, portion of occiput and small portion of facial bones.

AREA G

DIVISION I

- 1 Skl. I S 1. Child, about 5-6 years. Skull complete, without mandible; size small, frontal bone marked with a depressed groove in the mid-sagittal plane (pl. XXXI fig. 1). Parietooccipital region also shows depressed fracture; first and second milk molars of both the sides are intact, and permanent first molar lies deep in the socket.

 Skl. I S 11. Adult, 30-40 years; male. Skull complete with
- mandible.
 Skl. I S 13. Young adult, 21-25 years; female.
- without mandible, size small, not well preserved.
 Skl. I S 15. Child, 7-9 years. Skull very small without mandible; not well preserved.

DIVISION II

- Skl. II S 2. Adult; sex (?). Skull fragments with several teeth.
- Skl. II S 5. Adult, 25-30 years; female. Skull complete without mandible.
- Skl. II S 18. Adult, 25-30 years; male. Skull complete. Skl. II S 36. Adult; female. Skull incomplete and in pieces; all the sixteen maxillary teeth intact.
- Skl. II S 42. Adult, 25-30 years; male. mandible.

DIVISION III

- Skl. III S 1. Infant, 2-5 years. Skull very small, without mandible; reconstructed; maxillary teeth represented by right deciduous canine, first and second milk molars, and in addition to this crowns of both upper permanent first molars sprouting.
- Skl. III S 2. Adult, 25-30 years; male. Skull complete. Skl. III S 3. Adult. Skull in fragments without mandible;
- Skl. III S 3. Adult. Skull in fragments without mandible; palate and maxillary dental arch wide.

 Skl. III S 4. Child, about 3-6 years. Skull small in size, entire facial part being missing. A deep oblique fracture is present in the frontal bone; bregma fontanelle closed.

 Skl. III S 7. Adult; probably male. Skull represented by fragments of parietal bones only.

 Skl. III S 8. Child, 6-7 years. Skull small in size. No
- Skl. III S 8. Child, 6-7 years. mandible.

- Skl. III S 13. Child. Skull. Fragments.
 Skl. III S 21. Adult, below 25 years; female.
 Skl. III S 22. Adult, 25-30 years; male. Skull full and without mandible.
- Skl. III S 23. Adult; male. Skull complete with mandible. Skl. III S 24. Adult. Sex (?). Fragments of parietals. Skl. III S 25. Adult. Represented by fragments of parietal bones.
- Ski. III S 46. Adult. Skull represented by left sagittal half with some of the maxillary teeth.
- Skl. III S 47. Adult; male. Skull complete without mandible. In addition to the above cranial materials many frag-mentary long bone pieces along with a complete radius (Skl. II S 51) were found mixed up with heaps of animal

bones.

MOUND AREA

1 Skl. Mound A. Aduit; male. Skull represented by frontal and parietal fragments; proximal 2/3rd of right femur and

- 2 Skl. B 10. Adult; sex (?). Fragments of left temporal bone, occipital bone; left condyloid process of mandible. right shaft of humerus.
- Skl. B 722. Adult; female. Parts of occipital bone; right half of the mandible; rib fragments, scapula pieces 2, broken part of right clavicle, 5th metacarpal (rt.), acetabulum fragment, shaft fragments of right and left femur, fragment of right tibia, distal end of left fibula. Skl. B 1556. Child. Skull—lower half of the frontal bone.
- Skl. Mound AB (Recent). Adult, 21-25 years; probably
- female. Skull with mandible complete. Skl. HP XXX No. 421. Child; about 8-9 years. Skull incomplete, greater portion of the face and right side being missing.
- Skl. 783. Adult; sex (?). Left patella found mixed up with animal bones.
- Skl. 789. Adult; probably female. Left Os coxae. Skl. 4899. Adult; male. Broken parts of pelvis.
- Skl. 5440. Child. First and second cervical vertebrae, coracoid process of right scapula; shaft fragments of both humeri, fragments of radius, shaft fragments of right and left femur, broken parts of fibulae.
- Skl. 5440 A. Adult; probably male. Incomplete mandible having inverted gonial angles and prominent chin.
- Skl. 5440 C. Adult, 40-50 years; male. Skull represented by left sagittal half; no mandible. Skl. 5440 D. Adult, 25-30 years; probably female. Skull
- incomplete. Endocranial brain cast. Skl. 5461. Adult; sex (?). Ninth thoracic vertebra. Skl. 7740. Adult; sex (?). Broken part of left fibula.

- Skl. 11635. Adult, 52-30 years; male. Skull complete with mandible; second cervical vertebra and body of a typical thoracic vertebra, broken part of right scapula, medial 2/3rds of right clavicle, shaft fragments of right and left humerus.
- Skl. P III 76. Young adult; probably female. Skull represented by parietal, temporal and maxilla of the right
- represented by parietal, temporal and maxila of the right side; 5th metacarpal (right), proximal phalange (hand) 1. Skl. P III 76. Child. Proximal 2/3rd of right humerus. Skl. P IV 95 b. Infant. Skull parts include parietal and orbital bone of the right side with fragments of mandible; vertebral bodies 5, tiny ribs 12, left scapula right and left clavicle, complete right and left humerus, proximal part of right ulna, complete right radius, phalanx 1, complete right femur, broken parts of left femur, shaft fragments of tibiae, distal 2-3rd of right fibula and complete left fibula.
- Mound F IV. Adult, Sex (?). First cervical vertebra. rib fragments 2, fragments of left scapula, broken parts
- of rights clavicle, distal part of right tibia.

 Skl. Mound F X. Adult; male. Skull fragments; rib fragments 2, broken part of right ulna, part of a radius, shafts of right and left femur, fragments of tibiae and first; metatarsal (right).
- Skl. Mound F 3686 A. Adult; female. Nearly complete 22 left tibia.
- Skl. Mound J (I). Infant. Right orbit and right zygoma; cervical vertebra 1, head and spine of right scapula, proximal part of right ulna, shaft fragment of left femur, tibia and fibula fragments.
- Ski. Mound J (II). Child, 3-6 years. Right orbit, part of Skl. Mound J (II). Child, 3-6 years. Right orbit, part of parietal bone; fragments of vertebra, rib fragments, complete left humerus, nearly complete right femur, proximal part of right tibia distal part of left fibula, Skl. Mound J. Child, 6-9 years. Skull represented by right orbit, right parietal and right temporal, right-half of
- right orbit, right parietal and right temporal, right-half of mandible; left scapula, proximal part of right humerus, shaft of left ulna, shaft of right radius, complete left radius, metacarpals 3, proximal phalanges (hand) 5, fragments of pelvis, shaft fragments of right tibia, broken parts of right fibula, proximal phalanges (foot) 3.

CEMETERY H STRATUM II (Open Burials)

1 Skl. H 88. Adult, about 25:30 years; female. Skull incomplete; mandible in two pieces; cervical vertebrae 6th and 7th, theracic 1st, lumbar 1st to 3rd in one mass; osteo-arthritic change noticed at 4th body piece; frag-mentary left & right scapulae; lateral third of right clavicle; complete left clavicle; fragmentary humen;

- right ulna in two pieces; left ulna broken; right and left radius incomplete; metacarpal 1st, 2nd and 5th; proximal phalange 5; intermediate 4 and distal 4; incomplete pelvis having wide greater sciatic notch; right and left femora; right patella; both the tibiae in pieces; incomplete fibulae:
- left talus; left calcaneum and phalange 1. Skl. H 184 (K). Adult, about 25-30 years; male. Incomplete disintegrated skull; cervical vertebrae 1st to 3rd in fragmentary condition, thoracic 3rd to 7th in one mass and 12th, one lumbar having osteo-arthritic change; sternum; 12th, one lumbar having osteo-artific change; sternum; fragmentary scapulae; medial third of right clavicle; distal half of right humerus; complete left humerus; right ulna having a fracture at distal end; left ulna in pieces; broken radii; incomplete pelvis having greater sciatic notch; both the femora in pieces; incomplete tibia; right and left fibulae in pieces; head of right talus fractured and twisted; 2nd cuneiform of feft; right carpal 1st, 3rd and 4th; metacarpal 1st, and 5th; proximal phalange 3; left carpal 1st, 2nd, 5th to 8th; metacarpal 1st, and, 4th and
- 5th; proximal phalange 5, intermediate 4, distal 4. Skl. H 306 (a). Adult, about 18-25 years: female. Disinte-Skl. H 306 (a). grated skull; broken mandible; cervical vertebrae 1st, 2nd, and 3rd to 5th in one mass; thoracic typical, 2nd to 12th in five separate masses; lumbar 3; sacrum in pieces; both scapulae in fragmentary condition; right clavicle in two pieces, and medial two third of left; both humeri having fused epiphyses; incomplete right and left ulna; right radius distal half and left incomplete right metacarpal (hand) 5; left carpal 2nd, 7th and 8th; metacarpal 5, phalange proximal 5, intermediate 4; broken femora in pieces; pieces of right and left patella; decayed right tibia; left tibia slightly bowed medially having fused epiphysis; right and left fibulae in pieces; tarsals, calcaneum, talus, navicular, cunieform 3, cuboid, metatarsal 5
- foot: tarsals, navicular, cumeform 3, cuboid, metatarsal medial 4 and phalange (toe) one of left foot.

 Ski. H 307 (a). Adult, about 40-50 years; male. Decayed skull; no mandible; cervical vertebrae typical 2, thoracic 2, lumbar 1 to 5 body only; right hip-bone with fused epiphysis; broken femora.
- cpiphysis; broken remora.

 Skl. H 484 (a). Adult, about 25-30 years; male. Incomplete skull; cervical vertebrae 5th, 6th, 7th in one mass; thoracic 1st to 12th in five masses; flattened sacrum piece; theracic 1st to 12th in five masses; flattened sacrum piece; sternum not fused; fragmentary scapulae; incomplete humeri; right ulna full; left ulna distal third; broken and incomplete radii; right carpal 1st, 3rd to 5th, 7th, 8th; left carpal 1st, 3rd and 4th; phalange proximal 3, intermediate 3, terminal 1; pelvis incomplete having fused crest epiphyses; fragmentary femora; tibiae in pieces; fibulae incomplete; right tarsals, navicular, cunieform 3, fibulae incomplete. The considering all right calcaneum cuboid; left navicular, cunieform 3; right calcaneum broken having big sustentaculum tali; left calcaneum broken: left talus: metatarsals right 5, phalange proximal 3, intermediate 4, terminal 3.
- 3, intermediate 4, terminal 3.
 Skl. H 485. Juvenile, 12-14 years; female. Fragmentary skull and mandible bones; 14 intact mandiblar teeth without 3rd molars; theracic vertebrae 3 in one mass; without 3rd modars; theracic vertebrae 3 in one mass; without 3rd modars; theracic vertebrae 3 in one mass; other fragmentary pieces of vertebrae; sternum; incomplete scapulae; clavicle right complete; two third of right humerus; shaft piece of left humerus; shaft of right ulna, right capitate; phalange proximal 3, intermediate 1; left hip-bone without epiphysis union; shaft pieces of femora; shaft pieces of tibiae; calcaneum right, apophysis not fused; right and left talus fragments; left and right navicular; right cunciform 3, left 3rd; left cuboid; right metatarsal 5, left 1st and 2nd; phalange proximal 7. Skil. H 488 [H 487 (a)]. Adult, 25-30 years; male. Skull damaged by the action of saltpetre. No reference to number H 488 could be found and in place No. H 487 (a) has been considered as indicated by Vat (1940; 224). Skil. H 488*. Adult, 25-30 years; female. Incomplete skull; cervical vertebrae 4th to 7th and body piece of 1st and 2nd; thoracic vertebrae 1st to 1th; lumbar 1st to 5th in
- cervical vertebrae and to the and pace of 1st and 2nd; theracic vertebrae 1st to 11th; lumbar 1st to 5th in certain control of the state of

2nd to 5th, finger phalange proximal 3, intermediate 1; pelvis incomplete having fused crest epiphyses; complete femora; patella right piece; left tibia; proximal and distal ends of right fibula; left tarsal, navicular; right calcaneum; right metatarsal 1st to 5th, toe phalange proximal 5, intermediate 1.

5, intermediate 1.

Skl. H 501(a). Adult, 25-30 years; female. Incomplete and and bilaterally compressed skull; fragmentary mandible pieces; vertebrae in several masses; sacrum; fragmentary scapulae; complete claviculae; both humeri with united epiphyses; shaft pieces of radii; fragmentary ulna pieces; left metacarpal 3rd; phalange proximal 1, intermediate 2. terminal 1, pelvis incomplete; left femur; right femur m pieces; left patella; right tibia; crumbling left tibia; fragmentary fibulae pieces; right calcaneum broken anteriorly; disintegrated right and left talus.

10 Skl. H 502 (G). Adult, above 25 years; male. Skull somewhat damaged; incomplete mandible; 16 intact mandibular teeth; cervical vertebrae 1st to 7th in separate masses; thoracic 1st to 12th in one mass; lumbar typical 2; disunited manubrium and body of sternum; complete 1st rib of both sides; decayed scapulae; right clavicle, left in two pieces; humerus left and right; radius right; left and right ulna; disintegrated pelvis having narrow greater sciatic notch; both femora repaired; broken right patella left complete; left tibia; fragmentary fibulae; left calcaneum medially eroded; left cuboid.
Skl. H 503 I. Adult; male. Fragmentary skull bones; head

and spine portion of right scapula; middle portion of left

clavicle; broken pieces of left humerus; head and trochanter piece of one femur.

Skl. H 503 II. Child, about 6-12 years. Coccyx piece very small; humerus right shaft; proximal shaft of radius; hip-bone, ilium piece separate.

Skl. 636 Adult probably male Distorted and incomplete.

Adult, probably male. Distorted and incomplete skull; right clavicle; fragmentary shaft piece of humeri; pieces of left and right ulnae; pieces of right and left femur; left patella; right tibia; talus right and left; calcaneum left; right cuneiform 3; left 2nd and 3rd navicular; cuboid left piece; right metatarsal 3rd; left 5th and abolement processing the left piece; right metatarsal 3rd; left 5th and abolement processing the left piece; right metatarsal 3rd; left 5th and abolement processing the left piece; right metatarsal 3rd; left states and abolement processing the left piece process and pieces are respectively.

5th and phalange proximal 1.
Skl. H 694. Adult, above 25 years; male. Head, coracoid process and spine of left scapula; complete left humerus having united epiphysis.

15 Skl. H 695. Adult, 25-30 years; male. Skull slightly disintegrated by action of saltpetre; scapula right and left in pieces; complete right clavicle; right humerus; proximal pieces; complete right clavicle; right numerus, proximat pieces of right of left ulnae; proximal pieces of both radii; right metacarpal 2nd and 3rd; phalange proximal 4, intermediate 1, distal 1; left hip-bone disintegrated; femur left shaft; right tibia in pieces; distal pieces of one extra tibia; proximal part of left tibia; fragmentary

fibulae; phalange proximal 2, distal 1. Skl. H 696. Adult, 25 30 years; male. Fragmentary skull portion; theracic vertebrae 11th and 12th, lumbar 1st; right humerus; distal half of left humerus; pieces of right and left ulnae; distal end of left radius; metacarpal right 2nd to 5th, left 1st to 5th; phalange proximal 9, intermediate 5 and distal 1, fragmentary hip-bone; fragmentary pieces of femora; broken right tibia; pieces of left tibia; left fibula; shaft piece of right fibula; calcaneum, talus and navicular of both sides; right cunieform 2nd and 3rd, left 3rd; cuboid right and left; right metatarsal 1st to 5th, left 1st to 4th, toe phalanges Ist marked by lipping of edges; phalange proximal 8. intermediate a terminal 1. Extra left navicular 1st; left capitate 1st; distal shaft piece of right femur.

capitate lat; distal shaft piece of right femur.

Skl. H 697. Adult, above 25 years; male. Skuil badly crushed, incomplete and flattened bilaterally; mandible surving partly; cervical veriabree lower and theracic upper 2 in one mass; lumbar fragment 5 5th, fragment of sacrum; right scapula; left clavical; shaft of right humarus; left humers in pieces; right and left ulnae; fragmentary left predicts. fragmentary left radius; right navicular, lunate, triquetrum pisiform, capitate all in one mass; hammate; metacarpal left 1st, 6th to 8th; phalange proximal 8, intermediate 5.

distal 3; right hip-bone with fused crest epiphysis; left hip-bone with separate piece of os pubis; piece of right femur; shaft pieces of left tibia; fragmentary fibulae; right calcaneum; navicular, cuneiform 2nd and left 3rd. left cuboid 1st; talus fragments; right metatarsal 1st.

Skl. H 698. Adult, 25-30 years; male. Complete skull and mandible; cervical vertebrae 1st to 4th; thoracic 3rd to 12th in three masses; lumbar 1st and 2nd; complete 12th in three masses; lumbar 1st and 2nd; complete sacrum; manubrium and body of sternum fused; head of right scapula with acromion and coracoid processes; head right scapula with acromion and coracold processes; nead piece of left scapula, left clavicle; lateral third of right clavicle; shaft piece of right humerus; left humerus; right radius; left ulna; fragmentary right ulna, right carpal 1st to 3rd, 5th, 6th, 7th, and trace of 6th; incomplete pelvis with greater sciatic notch; right and left femora; shaft portion of tibiae; right and left patella; two pieces of left fibula; left calcaneum; left hammate; right talus.

Skl. H 699. Adult, 21-25 years; female. Skull with mandible in a locked-jaw condition; cervical vertebrae 3rd, 7th and typical 4; theracic 1st to 7th, having disunited annular contact. nular epiphyses; sternum; pieces of ribs, axillary border of right scapula; fragmentary left scapula; pieces of claviculae; head piece and distal end of right humerus; shaft of left humerus; fragmentary pieces of left ulna; proximal piece of right radius; fragment of 1st left metacarpal; phalange proximal 4; a lot of fragmentary

mentary long bones.

Skl. H 700. Adult, below 25 years; female. Fragmentary bieces; 14 mandibular teeth. cervical vertebrae 1st, 2nd, 6th and typical 2 of which annular epiphyses not united; thoracic 1st; right clavicle lateral third; pieces of right humerus; shaft piece of left humerus; shaft piece or right and left ulnae having fueed having fue fuel having fueed having fue fueed having fueed hav having fused epiphyses; fragmentary pieces of radii; right metacarpal 1st; fragments of left fibula; toe phalange proximal 1.

Skl. H 704. Child, about 12 years. Left parietal bone with

prominent parietal eminence.

Skl. H 707. Adult, sex? Skull survived by parietal pieces: right half of mandible; medial part of right clavicle, shaft fragments of left humerus; fragmentary left and right.

Skl. H 708*. Adult, sex? Lumbar vertebrae 1st, 2nd, 3rd in one mass; lateral two third of right clavicale, left femur shaft of the shaf

shaft of a child; shaft piece of fibula.

Skl. H 710. Adult, above 40 years; skull complete with mandible; cervical vertebra 1st to 7th; caudal piece of sacrum. sacrum; fragmentary left scapula; right clavicle in two pieces; left clavicle full curved and twisted slightly; humerus right full having sharp margins of bicipital groove. groove; left radius complete; left carpal 1st to 3rd and 5th to 8th, metacarpal 5, phalange 4; right carpal 1st. 5th to 8th, metacarpal 5, phalange proximal 5 intermediate 3, distal and thumb one each; broken pelvis having wide greater sciatic notches; right femur without head; left forms and the state of the scient patella; left and right patella; left and right patella; left and right patella; left forms and the scient patella; left and right patella; left and right patella; left forms and the scient patella; left and right patella; left forms and the scient patella; left and right patella; left and right patella; left and right patella; left forms and right patella; left and right patella; head; left femur complete; left and right patella; left and right patella; left and right patella; left fibula without head; and right tibia; right fibula; left fibula without head; left falue left talus, navicular, cuneiform 3 and cuboid; left metatargal 5, phalange proximal 5, distal great toe 1; right targal, navicular, cuneiform 3, cuboid; phalange 3, distal 2

Skl. H 734 (a). Adult, sex? Shaft of left humerus in two

Skl. H 739* 1. Adult, sex? Rib fragments: piece of left scapula; left elavicle in two pieces; distal end of right humerus; fragmentary left humerus; shaft of left ulna in three palaceters. in three peices; a lot of unidentified fragmentary bones.

CEMETERY H STRATUM I (Jar Burials)

IETERY H STRATUM I (Jar Burials)
Lists of skeletal materials have been furnished by M. S.
Vats in 'Excavations at Harappa', Vol. I, pp. 212-219,
242-245; and by R. E. M. Wheeler in 'Harappa 1946:The
Defences and Cemetery R 37', pp. 89-96.

COLLECTIVE TABLES

				ГЛ 1	M 8	M 2 a	M 17	M 9	M 21	M 25	M 24 b
No.	Skl No.	Sex	Age	Maximum Cranial length	Maximum Cranial breadth	Nasion-inion length	Basion-bregma height	Minimum frontal breadth	Vertical porion height	Median sagittal arc	Vertical transversal arc
CEMI	ETERY R 37										
1	H 779(e)	М	A	191.0	130.0	182.5	133.5?	94.0	112.5		295.0
2	H 793	M	A	193.0	133.0	176.0	129.0	_	111.0?	3 9 5. 0	298.0
3	H 793(A)	M	AH	190.5	152. 0	173.0	137.0	102.0	121.0	397.0	322.0
4	H 793(B)	M	A	186.0	131.0	174.0	129.0	99.0	116.0	385.0	301.0
5	Н 794	M	A	192.0	138.5	175. 0	138.0	90.0	118.5	3 8 5. 0	309.0
6	H 796(B)	М	A	178.0	135.0	164.0	132.5	95.0	114.0	353. 0	298.0
7	H 798(A)	M	A	189.0	131.0	177.0	131.5	97.0	113.0	37 6.0	299.0
8	H 798(B)	M	A		135.0		_	96.0	_	_	
9	H 798(C)	M	A	188.0?	131.0		137.0	90.0			
10	Н 806	M	A	189.0	131.0	174.5	128.0	93.0	110.5	376.0	294.0
11	H 811	М	A	184.0	132.0?	176.0	136.0	97.0	119.0	356.0	_
.12	H 818	M	A	188.0	130.0	170.0	_		117.0	378. 0	
13	H 820(iii)	Μ	A	184.0	134.0	157.0	135. 0	97.0	118.5	377.0	3 0 3. 0
14	Skl. 1	Μ	A.	188.0	140.0	180.0	131.0?	97.0	115.0	380.0	3 0 5. 0
15	Skl. 10	М	A	188.0	135.0	179.0	145.0	97.0	115.0	367.0	304.0
16	H 801(B)	M	Juv.		-	-	********				
17	H 779,(a)	F	A	182.0	-	171.0	126.0		110.5	362.0	291.0?
18	H 779(c)	F	A	184.0	122.0?	167.0	130.0	91.5	108.0	3 78.0	-
19	H 779(d)	F	A) comment		83.0			*******
20	H 780	F	A	181.0	137.0	173.0	127.0	96.0	112.0	364.0	314.0
21	H 788	F	A					Prince.		-	Promise
22	H 791/A	F	Α	174.0	127.0	16 5. 0		92.0	104.0	parents*	297.0

	M 23	M 40	M 48	M 47	M 45	M 55	M 54	M 50	M 51	M 51	M 52	2 M 5
No.	Horizontal circumference	Prosthion-basion line	Nasion-prosthion line	Nasion-gnathion line	Bizygomatic breadth	Nasal height	Nasal breadth	Ant. interorbital breadth	Orbital breadth (right)	Orbital breadth (left)	Orbital height (right)	Orbital height (left)
CEME	TERY R	37										
1	522.0	96.0?	71.0	_	_	53.0	22.0	18.0	42.0	41.5?	37.0	34.0?
2	530.0	106.0	71.0	-		48.0		_		_	_	_
3	548.0	106.0?	67.5?	_	141.0	56.5	27.0	23.0	49.0	46.0	39.0	41.0
4	517.0	98.0	68.0	_	_	51.0	25.0	21.0	40.5	41.5	36.0	32.0
5	526.0	98.0?	75.5	135.0	132.0	53.0	27.0	19.0	44.5	40.0	35.0	35.5
6	500.0	101.0	77.0	132.0		53.5	28.0	23.0	39.0	40.0	36.0	38.0
7	523.0	104.0	69.5	_	125.0	50.0	26.5	19.0	41.5	41.5	30.5	30.5
8		-	_	_	_					43.0	_	35.5
9	_		_	_	-		-			-	-	
10	518.0	104.0	69.5	118.5	133.5?	53.0	30.0	20.0	44.0	44.5	35.5	35.€
11	508.0	113.0	74.0	_	_	54.0	26.0?	19.0	42.5	_	34.0	33.5
12		_	68.5	_	122.0	48.0	25.0	18.0	43.5	42.0	34.0	31.5
13	514.0	98.5	61.0	_	138.0	48.0	26.0	20.0	42.0	41.5	28.0	31.5
14	532.0	102.0	69.5	123.0	137.0	53.5	31.0	18.0	42.0	45.5	33.5	32.5
15	530.0	102.0	73.0	120.0		5 8 .5	27.0	18.5	44.0		36.5	37.5
16		89.0	69.0	116.5	-	53.5	25.0	19.0		43.0	_	
17	493.0?	93.0	70.0	_		49.0	23.0	15.0?	45.0?	43.0	38.0	37.0
18		94.0	63.0	_		45.0	24.0?	21.0	40.0	0 -	34.0	34.0
9			-	-	_		-		41.0			32.0
0	503.0		65.0		*****	48.0	23.0	16.5	42.0	44.5	33.0	_
1			65.5	-		45.0	25.0 ?		-		32.0	33.0
2	491.0	-	Principal Control of C	-	125.0			20.0	36.0?	_	-	-
					-20.0			30,0	00.0	-	33.07	-

	M 60	M 61	M 62	M 63	M 66	M 65	M 70	M 71 (1)	M 71	M 69	M 68	M 79
No.	Maxillo-alveolar length	Maxillo-alveolar breadth	Palatal length	Palatal breadth	Bigonial breadth	Bicondylar breadth	Ht. of mandibular ramus	Max. breadth of mandibular ramus	Min. breadth of mandibular ramus	Ht. at mandibular symphysis	Mandibular length	Mandibular angle
CEMETER	X R 3	7										
1	58.0	61.5	46.0	35.0	_	_	_	_	_	_	_	_
2	61.0	68.0	_	45.0	83.0	99.0	60.0	44.0	33.5	38.5	90.0	127°
3	59.0?	70.0	46 p ?	46.0	90.0	_		_	39.0	_	91.0	_
4	63.0		47.0	36.0	_	_	65. 0	_	37.0	_	78.0	118°
5 ·	_	_	49.0?	_	96.0	121.0	67.0	44.5	30.0	40.0	75.5	126°
6	60.0	69.0	53.0?	40.0	93.0	130.0	_	42.0	32.0	37.0	78.5	_
7	58.0	62.0	50.0	39.0	85.0?		63.0	41.0	31.0	31.0	74.0	116.5°
8					_	Tomore	_	_	_			_
9						_	_		_	-		
10	60.0	68.0	55.0	41.0	81.0	117.5	69.0	49.0	36.0	35.0	85.0	117.5°
11	_	-	47.0	37.0		_	_	_		_	_	_
12	_	63.0	43.0	41.0	97.0	114.0?	_	39.0?	31.0	26.0	74.0	120°
13	56.5	67.0	44.0	43.0		_	-				_	_
14	6 3. 0		_		101.0	135.0	74.0	_	33.0	34.0?	82.0	118°
15	58.0	64.0	50.0	37.0	96 .5	128.0	65.0	52.0	36.0	29.0	82.0	128°
16	50.0	64.0?	numerod .				-	37.5	30.5	29.5		-
17	53. 0	62.5	48.0	41.0		-	News	_		36.0	71.0?	-
18	50.0	64.0	42.0	40.0	_		Proper	_	-			
19	_	Statement .				-	_	emesa	Primary		-	
20	58.0	62.0	48.0	41.0			Menny		20		-	
21	54.0	5 9 .5	46.0	39.0		_		_	_	named to		
22	******	-				-			-			

	-	M (1+8+1)	7)			,		J			
		3	M 8/M 1	M 17/M 1	M 17/M 8	M 21/M 1	M 21/M 8	M 9/M 8	M 47/M 45	M 48/M 45	M 66/M 45
No.	Calculated cranial capacity in c.c.	Cranial module	Length-breadth index	Length-height index	Breadth-height index	Length-auricular height index	Breadth-auricular height index	Transverse fronto-parietal index	Total facial index	Superior facial index	Jugo-mandibular index
CEMI	ETERY R	37									
1	1355.66	151.50?	68.06	69.90?	102.69?	58.90	86.54	72.31	_	_	_
2	1375.33?	151.67	68.91	66.84	96.99	57.51?	83.46?	_	_	_	-
3	1605.68	159.83	79.79	71.92	90.13	63.52	79.61	67.11		47.87?	6 3. 8 3
4	1367.30	148.67	70.43	69.35	98.47	62.37	88.55	75.57		MATRIANA.	-
5	1481.60	156.17	72.14	71.88	99.64	61.72	85.56	64.98	10 2.27	57.20	72.73
6	1336.67	148.50	75.84	74.44	98.15	64.04	84.44	70.37	_		
7	1357.20	150.50	69 .31	69.58	100.38	59.79	86.26	7 4.0 5		55. 60	68.00?
8	_		_	_	_	_		71.11		_	_
9	1422.09?	152.00?	69.68?	72.87?	104.58			68.70		_	
10	1335.42	149.33	69.31	67.72	97.71	58.47	84.35	70.99	88.76	52.06	60.67?
11	1389.77?	150.67?	71.74?	73.91	103.03?	64.67	90.15?	73.48?	-		
12	1378.93		69.15	_		62.23	90.00	_		56.15	79.51
13	1400.84	151.00	72.83	73.37	100.75	64.40	88.43	72.3 9	_	44.20	-
14	1437.82		74.47	69.68?	93.57?	61.17	82.14	69.29	89.78	5 0 .73	73.7 2
15	1399.77	156.00	71.81	77.13	107.41	61.17	85.19	71.85	-		-
16	_		_						Muddleto		-
17			-	69.23		60.71	_		quidhis	-	-
18	1198.35?	145.33?	66.30?	70.65	106.56?	58.70	88.52?	75. 00 ?	-	-	-
19	_		-	-	-		and the second		Manage		
20	1345.15	148.33	75.69	70.17	92.70	61.88	81.75	70.07	_	-	-
21	_		***************************************		**************************************	_	a		Ness		-
22	1145.86	nastrine	72.99	Name		5 9 .77	81.89	72.44		Promote	-
							*				

	M_52/M 51	M 52/M 51	M 54/M 55	M 61/M 60	M 63/M 62	M 68/M 65	M 66/M 65	M 45/M 8	M 48/M 17	M 40/M 1	M 9/M 45
No.	Orbital index (right)	Orbital index (left)	Nasal index	Maxillo-alveolar index	Palatal index	Mandibular index	Breadth index of mandible	Trans, craniofacial index	Vert. craniofacial index	Longitudinal craniofacial index	Jugo-frontal index
CEMETI	ERY R 3	7				A CONTRACTOR OF THE PARTY OF TH					
1	88.10	81.93?	41.51	106.03	76.09	_	_		53.18?	50.26?	_
2				111.48	_	90.91	83.84		55.04	54.92	
3	79.59	89 .13	47.79	118.64?	100.00?			92.76	49.27?	55.64?	72.34
4	88.89	77.11	49.02	_	76.60		_		52.71	52.69	· —
5	78.65	88 .75?	50.94			62.40	79.34	95.31	54.71	51.04?	68.18
6	92.31	95.00	52.34	115.00	75.47	60.38	71.54		58.11	56.74	_
7	73.49	73.49	53.00	106.90	78.00	_	_	9 5.42	52.85	55.03	77.60
8		82.56	_			-	_		_	_	_
9	*******					_	_	-			
10	80.68	78.65	56.60	113.33	74.55	72.34	68.94	101.91?	54.30	55.03	69.66?
11	80.00	_	48.15?		78.72		_	-	54.41	61.41	
12	78.16	75.00	52.08		9 5.3 5	64.91	85.09	93.85			
13	66.67	75.90	54.17	118.58	9 7.73	_	_	102.99	45.19	53.53	70.29
14	79.76	71.43	57. 9 4	_	_	60.74	74.81	97.86	53.05?	54.26	70.80
15	8 2. 9 5	83.33	46.15	110.34	74.00	64.06	75.39		50.34	54.26	_
16	Pilling	86.0 5	46.73	128.00?	erezza.	-	_			_	
17	84.44?	79.07	46.94	117.92	85.42		_		55.56	51.10	
18	85.00	82.05	53.33?	128.00	95.24		_ ,		48.46	51.09	_
19	80.49	W-1000	parame		-	-	_				
20	76.19	79.52	47.92	1000	85.42		_		51.18	-	
21		Manager 1	55. 56 ?	110.19	84.78	Printing	_				
22	91.67?	**************************************		approved		-	-	98.43		-	73. 60

Maxillary teeth

	M 81	M 81 (1)	M 81	M 81 (1)	M 8\$	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)
No	Mesiodistal crown diam. Right M1	Labiolingual crown diam. Right M 1	Mesiodistal crown diam. Left M 1	Labiolingual crown diam. Left M1	Mesiodistal crown diam. Right M 2	Labiolingual crown diam. Right M 2	Mesiodistal crown diam. Left M2	Labiolingual crown diam. Left M2	Mesiodistal crown diam. Right M3	Labiolingual crown diam. Right 'M 3	Mesiodistal crown diam. Left M3	Labiolingual crown diam. Left M3
CEME	TERY F	R 37										
1	9.0	11.5	10.0	11.5	9.5	12.0	7.5	11.0	9.5	11.0	8.5	11.0
2	10.0	10.5	11.0	12.0	10.0	12.0	10.0	11.5	11.0	12.0	10.0	12.0
3	10.5	11.5	11.0	12.0	10.5	10.0	10.0	10.5	9.5	10.5	10.0	11.0
4	9.5	11.5	10.0	11.5	10.5	11.5	10.5	11.5			_	-
5	_		_		10.5	11.5	_	_	10.5	11.0		
6	*****	_		_	_	_	-		_	-		. —
7	10.0	11.0	10.0	10.5	10.5	11.0	10.0	10.5	8.5	9.5	_	-
8	-	_	_	_	_	_		-			_	_
9	_	_	_	-				_		and the	-	
10	11.0	12.0	_	_	10.0	11.5	10.0	10.5				
11	10.0	10.5	_		10.0	10.5	_	-	7.5	10.0		
12	11.0	11.0	9.5	9.0	10.5	11.0	9.0	9.0		_	al control	
13	11.0	11.0	10.5	11.5	10 .5	_	10.0	12.5	9.0	10.5	LAURING	
14	VIIII POR			_						_		
15	11.0	11.5	10.0	11.5	8.5	10.0	9.0	11.0		named .	_	
16	10.0	10.5	11.0	12.0	errore.		9.5	10.0			8 .5	8.0
17	10.0	10.0	photos	-	9.5	9.0		-			7.0	6.5
18				-	-	generated	9.0	8.0				
19	District	annual .				name of the same o	-		-	9.0	8.5	10 =
20			10.0		10.0	9.0	8.5	10.0	9.0	7.0	7.0	10.5
21	10.0	9.0	10.0	9.0	9.0	8.0	7.0	8.0		_		7.5
22		NAME OF TAXABLE PARTY.	wydraeth	Attivida	Product		-	-			_	Medican

																						CH		1
10) 10)	21	26	19	18	17	16	10	14	13	12	11	10	9	00	7	. 6	ΟΊ	4	ω	N		CEMETERY	No.	
1	1	1	1		1	11.0	11.0	-	I	10.0	Ì	10.0?	Þ	l	l	I	10.5	11.0	l	10.5	b	R	Mesiodistal crown diam.	
1	1	1	1	ł		10.0	11.0	I	1	10.5	1	10.5	1	1	I	1	10.5	9.0	I	11.0	1	37	Labiolingual crown diam.	3
was	1	1	Ť	1	l	11,5	10.5	9.0?	1	10.0	l	L	1	1	4 =	1	ı	10.0	ı	10.5	1		Mesiodistal crown diam.	a
	1	No.	l			10.0	11.0	11.5	ı	10.0	I	ı	ı	ļ	I	ı	i	9.0	1	11.0	1		Labiolingual crown diam.	<u> </u>
	ŀ	Ĭ	ł	1	9.0	10.0	9.0	1	I	10.0	1	10.5	1	1	1	1	11.0	9.0	ı	10.0	1		Mesiodistal crown diam.	g E
1	1	1	4	1	7.5	9.0	10.5	1	1	9.0	1	9.5	I	1	1	1	10.5	7.5	1	10.5	I		Labiolingual crown diam.	۵ -
ľ	l	Ĺ	İ	1	9.0	10.5	9.0	L	1	10.0	1	I	I	1	1	1	11.0	10.0	1	11.0	1		Mesiodistal crown diam.	7
1	ı	1	1	1	7.5	9.0	10.5	I	L	9.0	1	I	1	I	1	1	10.5	7.5	I	11.0	I		Labiolingual crown diam.	o o
1	I	;	I		& 51	1		I	4	10.5		8.0	I	l	1	1	11.5	9.0	ı	10.0	I		Mesiodistal crown diam.	0
1	1	.	l	1	7.0	1	9.0	ı	١	9.0	1	9.5	ŀ	1	I	I	10.5	7.5	ı	10.0	1		Labiolingual crown diam.	ד
ı	1	I	I	1	9.0	1	9.0	10.5	I	10.5	1	1	1	1	ı	1	12.0	8.51	I	10.0	1		Mesiodistal crown diam.	
1	1	1	1	1	7.0	1	9.5	9.0	1	9.0	1	1	1	1	ľ	1	10.5	7.5	1	11.0	1		Labiolingual crown diam.	

	Crov	vn inc	lex of M81	maxill I (1)/M 81	ary t	eeth	Crown	inde	x of m M 81	andibu (1)/M 81	lar to	eth
							i H					
No.	Right M 1	Left M 1	Right M2	Left M2	Right M3	Left M3	Right M1	Left M1	Right M2	Left M2	Right M3	Left M3
CEMETH	ERY R 3	7										
1	127.78	115.00	126.32	146.67	115.79	129.41	_	_	_		_	_
2	105.00	109.09	120.00	115.00	109.09	120.00	104.76	104.76	105.00	100.00	100.00	110.00
3	109.52	109.09	95.24	105.00	110.53	110.00	_				_	-
4	121.05	115.00	109.52	109.52	_	_	81.82	90.00	8 3.33	75.00	83.33	88.2
5	_	_	109.52	_	104.76	_	100.00	_	95.45	95.45	91.30	87.50
6	_		_	_	_	_				-	_	
7	110.00	105.00	104.76	105.00	111.76	_			~	_		
8		_		_						,	_	
9	_			_	_		_		_			_
10	109.09		115.00	105.00	-	_ ,	105.00 ?		90.48	and the same of th	1,18.75	
11	105.00		105.00	-	133.33					A.		-
1 2	100.00	94.74	104.76	100.00	-		105.00	100.00	90.00	.90.00	85,71	85.7
13	100.00	109.52	-	125.00	116.67	_			_	-	المعارات	
14	-		_	-			-	127.78?	7	-		85.71
1 5	104.55	115.00	117.65	122.22		******	100.00	104.76	116.67	11667	105.88	105.56
16	105.00	109.09		105.26		******	90.91	86.96	90.00	85.71	Nacion .	
1 7	100.00		94.74			94.12			- 83,33	88:38	82:35	77.7
18			-	88.89	_	92.86			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ 	14	_	Hanna
19	-		-	-		er-mail				-		
20		_	90.00	117.65	100.00	123.53			_		1	-
21	90.00	90.00	88.89	114.29	87.50	107.14				-	-	
22	· ·	-	THE	******	-			~ _		Million	-	

В

Linear measurements on craniograms

11					on cramo				
	W 42	W 43	W 44	W 48	W 49	W 50	W 51	W 52	W 53
			75	ht	calvarial height	position line	dicular	ndicular	endicular
	Frontal chord	Parietal chord	Occipital chord	Calvarial height	da calvar	na positio	Frontal perpendicular	Parietal perpendicular	Occipital perpendicular
No.	Front	Parie	Occip	Calva	Lambda	Bregma	Front	Parie	Occip
EMETER	Y R 37			,					
1	117.0	123.0		101.0	75.0	95.5	24.5	24.5	_
2	_	_	_	_	_	_			_
3	117.5	142.5	85.5	125.0	92.5	109.0	25.0	38.0	23
4	109.5	125.0	95.5	109.0	74.5	98.5	27.5	25.5	30
5	112.5	120.5	100.0	109.5?	73.5	100.0	26.0	23.0?	28
6	111.0	115.0	92.0	106.0	72.0	97.5	19.5	23.5	22
7	112.5	121.5	89.5	107.0	74.5	97.5	24.5	26.0	30
8	-		_	_	_	_			_
9	:		_	_	_		~	_	_
10	110.0	124.0	99.5	105.5	71.0	9 5.5	25.5	24.0	25
11	107.0	103.0	102.0	92.5	59.5	85.5	22.0	19.5	33
12	114.0	124.5		110.5	78.0	102.0	27.5	24.5	
13	107.0	122.0	99.0	115.5	75. 0	99.5	22.0?	26.5	29
14	108.0?	131.5?	94.0	104.5?	77.5?	94.0?	25.0	29.5	25
15	112.5	114.5	96.5	102.0	70.5	95.0	22.5	25.5	24
16		ana-rooms			***************************************	*******			_
17	105.5	104.0	105.5	102.0	62.0	92.5	21.0	19.0	38
18	115.0	100.5	104.5	107.0	67.0	102.0	29.5	14.5	3
19		province (0,5	11.	- Americans			-
20	113.0	106.5	108.0	110.0	66.0	94.5	24.0	22.5	2
21	Seems		_		PT-100				
22	Services.					turing.	and the same of th	-	-

			Angı	ular meas	urements o	on cranio	ırams			
	W 3	W 4	W 5	W 9	W 11	W 12	W 13	W 14	W 15	W 7
No.	Frontal inclination angle	Occipital inclination angle	Facial profile angle	Calvarial base angle	Frontal curvature angle	Parietal curvature angle	Occipital curvature angle	Occipital flexional angle	Superior facial length angle	Alveolar profile angle
CEME	TERY R	37								
1	60.5°	88°	84.5°	10.5°	134°	136.5°	_	_	42°	78°
2		_	_							-
3	68°	95.5°	_	19°	133°	123°	120°	120°		-
4	64.5°	84.5°	86°	13.5°	125°	134.5°	113.5°	113°	40.5°	76°
5	63°	84°	89°	11.5°	128°	137.5°?	118°	118°	43.5°	75°
6	62.5°	88°	81°	18°	141°	134.5°	125°	125°	44°	75.5°
7	60.5°	85°	81°	12°	132.5°	133°	111.5°	112°	3 9 .5°	8 3°
. 8			_	_		_		_	_	
9		and the second			enaments.	- '	_			_
10	61°	90.5°	78°	15°	129.5°	136.5°	124°	124°	40°	5 6°
11	53°	77.5°	8 3°	3.5°	135°	138.5°	124°	118°	3 8°	75°
12	63.5°	89.5°	80°	16°	128°	13 6 .5°				72°
13	68°	92°	77°	18°	133.5°?	132°	127.5°	125°	36.5°	64.5°
14	60.5°	88°	85°	10.5°	130°	132.5°	122°	122°	39°	7 9 .5°
15	57.5°	79.5°	7 9 .5°	11°	136°	131°	126°	1 26 .5°	40°	73°
16		and the same of th								-
17	62°	75°	81.5°	14°	13 6 .5°	13 9 °	108°	110°	43°	72°
18	61.5°	80°	86.5°	12°	124.5°	145°	1 0 4.5°	106.5°	38°	90°
19								, -		
20	57°	82°	8 3.5°	9 .5°	132.5°	134.5°	121°	121°	3 9 °	79"
21			remains			and the same of th			-	379864
22				retion					-	

				M 1	M 8	M 2a	M 17	M 9	M 21	M 25	M 24b
No.	Skl No.	Sex	Age	Maximum Cranial length	Maximum Cranial breadth	Nasion-inion length	Basion-bregma height	Minimum frontal breadth	Vertical porion height	Median sagittal arc	Vertical transversal arc
CEME	TERY R 37										
23	H 795/A	F	A		_			93.0	_		_
24	H 798/AI	F	A	171.0	138.0	162.0	_	89.0	111.0	358.0	313.0
25	H 798(a)	F	A	178.0	129.0	169.0	131.0	96.0	111.5	360.0?	297.0
26	H 801/A	F	Å	184.0	131.5	169.0	135.0	90.0	119.0	387.0	308.0
27	H 804	F	A			_	_	89.0	_	_	_
28	H 805	F	A	186.5	132.0	174.0	128.0		117.0	385.0	315.0
29	H 806/A	F	A	195.0	139.0	181.0		93.0	120.0	_	325.0
30	H 812	F	A	182.0	125.0?	170.0	125.0	100.0	109.0	372.0	294.0
31/	H 816	F	A	167.0	130.0	162.0	_	91.0	109.0	346.0	295.0
32	H 817	F	A				125.0?	93.0	107.5?	_	_
33	H 820(I)	F	A	181.0	129.0	166.0	125.0	94.0	113.0	366.0	309.0
34	H 820(II)	F	A	172.0?				97.0	_		_
35	Skl. 2	F	A	178.0	128.5	171.5	124.0	92.0	106.0?	353.0	289.0
36	H 798/A2	F	Juv.	166.0	128.0	155.0	116.0	87.0	102.0	330.0	275.0
a !											
AREA											
37	I S II	М	A	188.5	135.0	174.0?	126.5	96.0	115.0	387.0	307.0
38	II S 18	M	A	1790	142.0	170.0	133.0	106.5	115.0	365.0	315.0
39	II \$ 42	IVI	Α	178.0	129.0	166.0	139.5	90.0?		363.0	303.0
40	III S 2	M	A	171.0	139.0	162.0	137.5	105.0	115.0		310.0

	M 23	M 40	M 48	M 47	M 45	M 55	M 54	M 50	M 51	M 51	M 52	M 52
No.	Horizontal circumference	Prosthion-basion line	Nasion-prosthion line	Nasion-gnathion line	Bizygomatic breadth	Nasal height	Nasal breadth	Ant. interorbital breadth	Orbital breadth (right)	Orbital breadth (left)	Orbital height (right)	Orbital height (left)
CEMET	ERY R	37										
23	_	101.0	64.0	_		47.5	23.0	20.5	38.0	37.0	34.0	34.5
24	498.0	_	64.0	_	123.0	47.0	26.0	17.0	42.0	42.0	37.0	37.0
25	501.0	_	66.0		127.0	51.0	26.0	18.5	41.0	41.0	31.5	31.5
26	508.0	_	_	_	127.0	53.0	24.0	17.0	38.0	39.0	33.0	34.0
27	_	_	60.0	_		46.0	22.5	18.0	40.0	38.0	35.0	34.0
28	514.0		_	_	_	51.0	27.0		38.5	44.0		38.0
29	528.0	_	70.0	120.0	_	47.0	25.0		43.0	42.0	33.0	34.0
30	506. 0	85.0	72.0	_	_	53. 0	23.0	20.0	46.0	44.0	39.0	36.5
31	483.0	_	65.0	-	128.0	48.5	25.5	18.0	39.0	41.0	32.0	31.5
32	_	98.0	60.0	_	121.0?	48.5	24.5	14.0	41.0	40.0	33.0	33.5
33	504.0	_	******		_	42.0	28.5?	21.0	_	39.0	31.0	31.0
34			64.0	107.0		48.0	27.0	18.0	46.0	_	35.0	35.0
35	500.0	95.5	63.0	108.0	121.0	46.0	25.0	22.5	38.5	_	32.0	
3 6	466.0	*88.0	56.0	_	112.5	44.0	23.0	17.0	37.0	36.5	31.5	31.5
												*
AREA	G											
37	520.0	93.5?	69.0?	119.0		51.0?	26.0	18.0	43.0	40.5	32.0	32.0
38	523.0	99.0	66.0		128.0	51.0	25.0	23.0	42.0	44.5	29 .5	29 .5
3 9	495.0	94.0	66.5	water		51.5	25.0	17.0	40.0	38.0	32.5	32.5
40	505.0	*******	man	Vestigat	nonequi.	47.0	24.0	23.0	40.0	41.5	32.0	32.0

***	M 60	M 61	M 62	M 63	M 66	M 65	M 70	M 71 (1)	M 71	M 69	M 68	M 79
No.	Maxillo-alveolar length	Maxillo-alveolar breadth	Palatal length	Palatal breadth	Bigonial breadth	Bicondylar breadth	Ht. of mandibular ramus	Max. breadth of mandibula r ramus	Min. breadth of mandibular ramus	Ht. at mandibular symphysis	Mandibular length	Mandibular angle
EMETI	ERY R 3	37										
23	59.0	63.0	46.0	40.0	_	_		_	_	_	_	_
24 .	48.0	58.0	43.0	38.0		_	_	_	_	-	_	
25	56.0	63.5	41.0	39.0	_	_		_			_	_
26	_		1 -	_	89.0	117.0	62.0	44.0	36.0	31.0	80.0	120.5°
27			_	_	78.0?		57.5	38.0	30.0	24.0	66.0	119.5
28	_				_		-	_	_		_	_
29	56.0	59.0	47.0	40.0	79.0	104.0?	58.0	41.5	33.5	35.0	81.0	117.5
30	_			-	_		_	_	***	_		
31			-		_			_			_	
32	55.0	65.5	41.0	40.0	_	_		_		EW .		_
33			_				· · _	_		_	_	
34	57.0	61.0	48.0	40.0	87.0		60.0	38.0	30.0	28.0	73.0	120°
35	55.5	62.0	40.0	40.0	70.0	104.0	57.5	38.0	30.5	26.0	78.0	116°
36	56.5	58.0									-	_

A	Hib.	7077	A		~
/A	-846	101	Z3.	- 1	1.2

37	54.0	56.0	52.0	39 0	85.0	120.0	67.0	47.0	37. 0	35.0	83.5	113.5°
38	56.0	62.5	48.0	41.0	-			-				
39	49.0	67.5	46.0	44.0		-	-	_			-	
40	_	66.0		42.0			manufacture.			-	-	-

	N TOO SEEMEN ON THE PART OF TH	M (1+8+17)	M 8/M 1	M 17/M 1	M 17/M 8	M 21/M 1	M 21/M 8	M9/M 8	M 47/M 45	M 48/M 45	M 66/M 45
No.	Calculated cranial capacity in c.c.	Cranial module	Length-breadth index	Length-height index	Breadth-height index	Length-auricular height index	Breadth-auricular height index	Transverse fronto-parietal index	Total facial index	Superior facial index	Jugo-mandibular index
CEME	TERY R 3	7									
23	: te	· _	-	<u></u>	_	_			_	<u></u>	
24	1279.47	_	80.70	_	_	64.91	80.43	64.49	_	52.03	
25	1254.88	146.00	72.47	73.60	101.55	62.64	86.43	74.42	_	51.97	_
26	1387.61	150.17	71.47	73.37	102.66	64.67	90.49	68.44			70.08
27		_		_		_		_		_	
28	1388.02	148.83	70.78	68.63	96.97	62.73	88.64	<u> </u>		_	
29	1542.89	_	71.28	_		61.54	86 .33	66.91		-	and a
30	1221.39?	144.00?	68.68?	68.68	100.00?	59.89	87.20?	80.00?		_	
31	1174.23		77.84		_	65.27	83.85	70.00	_	50.78	
32				-			-	_		49,59?	
33	1287.40	145.00	71.27	69.06	96.90	62.43	87.60	72.87	-		
34	_							-	شت		-
35	1198.42?	143.50	72.19	69.66	96.50	5 9 .55?	82.49?	71.60	89.26	52.07	57. 8 5
36	-	136.67	77.11	69.88	90.62	61.45	79.69	67.97	_	49.78	5
AREA (Ž.						05.12	77 14			
37	1402.50	150.00	71.62	67.11	9 3.7 0	61.01	85.19	71.14		51 50	
38	1401.31	151.33	7 9 .33	74.30	93.66	64.25	80.99	75.00	- Control of the Cont	51.56	Weather.
39	1376.65	148.83	72.47	78.37	108.14	and the same of th		69.77?	- Transit	America	
40	1334.56	149.17	81.29	80.41	98.92	67.25	82.78	75.54		Manual	-

	M 52/M 5	1 M 52/M	51 M 54/M 5	5 M 61/M 60	M 63/M 62	2 M 68/M 6	55 M 66/M 6	5 M 45/M 8	M 48/M 17	M 40/M 1	M 9/M 45
No.	Orbital index (right)	Orbital index (left)	Nasal index	Maxillo-alveolar index	Palatal index	Mandibular index	Breadth index of mandible	Trans. craniofacial index	Vert. craniofacial index	Longitudinal craniofacial index	Jugo-frontal index
CEMET	ERY R	37									
23	89.47	93.24	48.42	106.78	86.96	_	_	. —	_	_	· —
24	88.10	88.10	55.32	120.83	88.37	_		89.1 3		-	72.36
25	76.83	76.83	50.98	113.39	95.12			98.45	50.38	_	75.59
26	86.84	87.18	45.28		_	68 .38	76.07	96.58			70.87
27	87.50	89.47	48.91	_		_	_				_ ×
28	_	86.36	52.94				-		_	agening.	-
29	76.74	80.95	53. 19	105.36	85.11	77.88?	75.96?			_	-
30	84.78	82.95	43.40		_				57.60	46.70	_
31	82.05	76.8 3	52.5 8					98.46	_	_	71.09
32	80.49	83.75	50.52	119.09	97.56				48.00?	-	76.86?
33	_	79.49	67.86						-		_
34	76.09		56.25	107.02	83.33	Principal Control	glasses	_		_	_
35	8 3.12	-	54.35	111.71	100.00	75.00	67.31	94.16	50.81	5 3. 6 5	76.03
36	8 5.14	86,30	52.27	102.65	Ministry		_	87.89	48.28	53.01	77.33
AREA	G										
37	74.42	79:01	50.98?	103.70	75.00	ß9.58	70.83		54.55?	49.60?	
38	70.24	66.29	49.02	111.61	85.42		******	90.14	49.62	55.31	83.20
39 40	81.25	85 .53	48.54	137.7 6	9 5. 6 5			-	47.67	52.81	
70	80.00	77.11	51.06				-	-	francis.	2)	

M a	х	i	1	1	a	r	V	t	е	е	t	h
-----	---	---	---	---	---	---	---	---	---	---	---	---

	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81 (1)	M 81
No.	Mesiodistal crown diam. Right M 1	Labiolingual crown diam. Right M 1	Mesiodistal crown diam. Left M 1	Labiolingual crown diam. Left M1	Mesiodistal crown diam. Right M 2	Labiolingual crown diam. Right M 2	Mesiodistal crown diam. Left M2	Labiolingual crown diam. Left M2	Mesiodistal crown diam. Right M 3	Labiolingual crown diam. Right M3	Mesiodistal crown diam. Left M3	Labiolingual crown diam. Left M3
CEM	ETERY	R 37										
23	10.0	9.0	10.0	10.0	9.0	10.0	10.0	10.0	10.0	10.0	10.0	10.0
24	_	-		-		_	_	_		ë <u> </u>	-	
25	10.0	12.0	10.0	12.0	10.0	11.0	11.0	11.5	9.0	10.0	9.0	9.0
26	_	-	_	_	_	_		_			-	_
27	_	_	_	. <u> </u>	_		_	_				
28	_	_	_	_				_			_	_
29	10.0	10.5	10.0	11.0	10.0	10.0	10.0	10.5	8:0	8.5	8:0	8,0
30	_	_		_	_			_		_	-	
31	_	-		_			_			_	-	-
32	10.5	11.0	10.5	11.0	10.0	11.5	11.5	11.0		_		_
33	_		_				_	-	_	-		
34		_		_) 			_	· ·		_
35	10.0	11.5	10.0	11.0	9.5	10.5	9.5	11.0	admit.	一种	,\	
36	9.5	10.5	10.5	10.0	9.5	10.0	9 .5	10.0		-	-	
				a								
								~				
	EA G								16.5	1:1.0	-	
37	10.0	11.0	11.0	11.5	10.0	12.5	11.0	10.0	8.5	10.0	9.5	0 1
38	10.5	12.0	10.0	11.5	9.5	11.0	11.0	10.5	9.5	12.0	9 .5	8.3
39	10.0	11.5	11.5	11.5	9.0	11.0	10.0	14.5	9.5	10.5		12.
40	11.0	11.5	11.0	12.0	11.0	12.0	9.5	1.1.0	0.0	6.91	8 .5	10

С

																						1	1:
40	38	ಬ ೦೦	37	A		3 6	35	34	33	32	31	30	29	28	27	26	25	24	23	CEN	No.		
-	1	1	10.0	AREA G		I	11.0	1	L	1	1	1	11.0	١	10.5	١	1	l	I	CEMETERY	Mesiodistal crown diam. Right M1	M 81	•
ŧ	1	1	12.	M2		l	10.0	ı	I	I.	1	l	10.0	l	10.0	I	l	١	I	Y R 37	Labiolingual crown diam. Right M 1	M 81 (1)	≤
l	i	1	11.5			i	10.0	1	I	1	ı	1	11.0	- 1	10.5	I	L	ı	١		Mesiodistal crown diam. Left M 1	3	ໝ
Í	ı	4000	12.0			1) 10.5	I	ı	1	1	ı	10.0	I	10.0	1	1	ı	ı		Labiolingual crown diam. Left M 1	M 81 (1)	COLLE
ı	1	1	0.11.0			1	5 10.0	ı	1	1	1	1	0 10.0	1	1	1	1	I	I		Mesiodistal crown diam. Right M2	M 81	OLLECTIVE
ı	-	I	0 12.0			1	0 10.5	1	ı	ı	1	ı	9.0	ľ	10.0	1	ŧ	ı	ı		Labiolingual crown diam. Right M2	M 81 (1)	TABLE-
ı	1	ı	0 11.0			ı	9.0	l	ı	1	i	L	10.5	1	1	I	1	1	1		Mesiodistal crown diam. Left M2	M 81	- B Contin
1	1	1	12.0			I	10.5	ı	1	ı	I	1	9.0	ı	ſ	ı	I	I	1		Labiolingual crown diam. Left M2	M 81 (1)	nued
ı	1	I,	10.5			I	10.0	1	1	ı	ļ	1	. 9.0	1	ı	1	1	1	I		Mesiodistal crown diam. Right M 3	M 81	Φ
i	1	ı	11.5			1	10.0	1	ı	ı	1	1	8.5	1	1	١	1	ı	1		Labiolingual crown diam. Right M3	M 81 (1)	
1	1	1	5 10.0			ı	9.0	1	ľ	1	ı	1	8.0	ı	10.0	1	1	ı	1		Mesiodistal crown diam. Left M3	M 81	
ı	j	ı	11.0			ı	10.0	1	1	1	1	1	7.5	ı	9.0	1	1	ı	ı		Labiolingual crown diam. Left M3	M 81 (1)	

	Crov	vn ind	lex of M 81 (1		ary te	eth	Crown	inde	x of M 81 (1	mandibu)/M 81	lar	teeth
No.	Right M 1	Left M 1	Right M2	Left M2	Right M3	Left M3	Right M 1	Left M 1	Right M2	Left M2	Right M3	Left M3
'EWET'	ERY R 3'											
23	90.00	100.00	111.11	100.00	100.00	100.00	_		_	_		_
24	_	_	_	_	_	_	_		_	. —	_	_
25	120.00	120.00	110.00	104.55	111.11	100.00			·	_		_
26	_	_	_	_	_	_	_			_		_
27	_	_	_			_	95.24	95.24	_	_	_	90.00
28	_	_	_							_	_	_
29	105.00	110.00	100.00	10 5.00	106 .25	100.00	90.91	90.91	90.00	85.71	94.44	9 3.75
30	_		_						_	_		
31		_			-				_	_	_	
32	104.76	104.76	115.00	95.65					_	_		_
33							manuta.	_	_	_	_	
34				_					_	_	_	
35	115.00	110.00	110.53	115.79		×	90.91	105.00	105.00	116.67	100.00	111.11
36	110 .53	95.24	105.26	105.26			_		_	-		
AREA		10455	105.00		115.70		120.00	104.35	109/09	109.09	109.5	2 110.0
37	110.00		125.00		115.79		120.00	104.00		Alleman		
38	114.29	115.00	115.79	90.91	117.65	89.47						

40 104.55 109.09 109.09 121.05 110.53 117.65 —

COLLECTIVE TABLE — B Continued

Linear measurements on craniograms

	W 42	W 43	W 44	W 48	W 49	W 50	W 51	W 52	W 53
	Frontal chord	Parietal chord	Occipital chord	Calvarial height	Lambda calvarial height	Bregma position line	Frontal perpendicular	Parietal perpendicular	Occipital perpendicular
No.		<u> </u>	. 0			— ш	<u> </u>	———	
EMETER	Y R 37							,	
23 24	 113.5	102.0	— 95.5	104.0	70.0		- 21.0	10.0	-
25	112.0	117.0	95.5 87.5	104.0	70.0	99.0	31.0	18.0	28.0
26	115.0	132.5	92.5	114.5	74.0 84.0	96.5 103.5	24.0 29.0	28.5 30.5	23.5 23.5
27					64.0	105.5	29.0	- -	23.
28	112.0	114.0	102.5	112.0	71.0	101.5	26.5	23.0	36.
29	116.5	118.5		112.0	71.5	101.5	25.0	26.0	
30	110.0	120.5	92.0	100.0	73.0	94.0	27.0	25. 0	28.
31	107.5	102.0	91.0	97.0	68.0	92.0	27.0	19.5	28.
32	105.0	_			**************************************	_	24.0?	-	_
33	104.0	101.5	97.0	104.5	64.0	93.5	26.0	21.0	`38.
34	_		materials.	_		_	_		
35	109.5	106.5	96.0	95.0	66.5	90.5	21.5	19.5	32
36					-			-	
		a Ugani sanca							
REA G	ν.								
37	117.0	125.0	92.0	110.0	80.0	104.0	29.0	24.0	31
38	108.0	110.5	98.0	104.0	70.0	96.0	28.0	22.5	29
39	106.5	116.5	92.0	106.0?	72.0	95.5	27.0	28.0	27
40						~~		-	

Angular measurements on craniograms W 3 W 4 W 9 W 12 W 13 W 14 W 15 .W 7 W 5 W 11 Superior facial length angle Occipital inclination angle angle Frontal inclination angle Parietal curvature angle Frontal curvature angle Alveolar profile angle angle profile angle Occipital curvature flexional Calvarial base Occipital Facial 1 No. CEMETERY R 37 23 84° 24 60.5° 80.5° 85° 13° 122° 140° 117.5° 115° 57.5° 3**9°** 25 60° 86° 80° 13° 133° 127° 123.5° 123.5° 64° 95° 15.5° 125.5° 130.5° 125° 125° 26 27 __ 92.5° 128° 42.5° 79° 85° 14° 110° 28 64.5° 135° 109° 73.5° 29 61° 82.5° 83° 10° 132° 132° 78.5° 48° 118° 3**0** 58.5° 91° 88° 12.5° 127.5° 134.5° 116.5° 76° 82° 13° 116° 57° 79° 125.5° 138.5° 115° 31 86.5° 36° 87° 130°? 32 --103.5° 64.5° 73° 7.5° 126° 134.5° 102° 33 ___ 34 63° 38.5° 112.5° 55.5° 82° 77° 16.5° 136.5° 13**9**.5° 112° 35 36 AREA G 111° 111° 63° 87.5° 14.5° 127° 13**8°** 37 39° 70.5° 117.5° 117° **80**.5° **8**3.5° 9° 123° 13**5**.5° 63° 38 117° 3**9°** 73° 117° **8**2.5° 84° 127.5° 13.5° 64° 126° 3**9** 40

				M 1	M 8	M 2 a	M 17	M 9	M 21	M 25	M 24 b
No.	Skl No.	Sex	Age	Maximum Cranial length	Maximum Cranial breadth	Nasion-inion length	Basion-bregma height	Minimum frontal breadth	Vertical porion height	Median sagittal arc	Vertieal transversal arc
AREA	G										
41	III S 22	M	A	182.0	144.0	168.0	128.0	101.0	109.0	354.0	300.0
42	III S 23	M	A	191.0	139.0	175.0	138.0	92.0	117.0	397. 0	311.0
43	III S 47	M	A	176.0	138.0	166.0	132.0	99.0	115.0	370.0	308.0
44	I S 13	F	A	_	_	_	_	86.0	109.0?	354.0	-4-
45	II S 5	F	A	175.0	135.0	160.0	126.0	92.0	110.0	372.0	292.0
46	III S 21	F	A	174.0	125.0	157.0	123.0	90.0	105.0		277.0
47	I S 1		Ch	166.0	126.0	157.0	110.0		101.0	343.0	301.0
48	I S 15		Ch	174.0		158.0	119.0	92.0	102.0	352.0	297.0?
49	III S 1		Ch	155.0?	_			84.0		_	-
50	III S 4		Ch	160.0	129.0		_	83.0?	_	350.0	323.07
51	III S 8		Ch	166.0	111.0		109.0	83.0	99.0	342.0	277.0

MOUND AB

52	H 11635	М	A	182.0	133.0	175.0	135.0	92.0	114.0	375. 0	309.0
53	5440 D	F	A	19 3.0?	135.0		-	99.0	-	********	319.0
54	Mound AB (Rec.)	F	Α	179.0	125.0	169.0	136.0	93.0	109.0	35 9.0	298.0
55	H.P. XXX 421		Ch	166.0	127.0	155.0	123.0	86.0	104.5	340.0	307.0

	M 23	M 40	M 48	M 47	M 45	M 55	M 54	M 50	M 51	M 51	M 52	M 52
No.	Horizontal circumferense	Prosthion-basion line	Nasion-prosthion line	Nasion-gnathion line	Bizygomatic breadth	Nasal height	Nasal breadth	Ant. interorbital breadth	Orbital breadth (right)	Orbital breadth (left)	Orbital height (right)	Orbital height (left)
AREA	G											
41	515.0	94.0	66.5	_	133.0	53. Q	29.0	22.5	44.0	41.0	33.0	34.0
42	523.0	96.0	70.0	_	123.0	52.5	26.0	17.0	39.0	3 9 .5	32.0	30.5
43	505.0	92.0	61.5	_	127.5	49.0	25.0	21.5	40.0	39.0	32.0	34.5
44	_	_	_	_	110.0?	42.0?	23.5	17.0?	35.0	35.5?	28.0	29.0
45	490.0	93.0	67.0	_	118.5	47.0	2 3.0	15,0	42.0	40.5	32.0	32.0
46	486.0	90.5	66.0	_	121.0	48.5	24.0	20.0	3 9 .5	39 .5	34.0	34.0
47	473.0	73.0	42.0	_	_	31.5	_	14.0	34.0		35. 0 ?	
48	_		54.0		106.0	42.0	21.0	16.0	36.0	34.0	28.0	28.5
49		74.0	46.5	_		33.5	20.0	18.0	33.0	32.0	28.5	31.5
50	449.0		_		_	_	_		-		_	₹,
51	448.0	_	44.0	_	-	31.0	18.0	17.5	35.0	3 2 .5	29 .5	28.5

MOUN	D AB											
52	506.0	92.5	69.0	116.0	134.0	52.5	27.0	21.0	42.0	42.5	35.0	35. 0
53	530.0?	*******		Manager		_	equality		_	and the same of th	_	*****
54	495.0	100.0	66.0	113.0	Monte	48.0	22.0	22.0	39.0	Specific Description of the Control	32.0	33.0
5 5	471.0	and a	*****	-	***		****	17.0	*******	WHISTON.	_	32.0

.744									11.74		14.00	14.70
	M 60	M 61	M 62	M 63	M 66	M 65	М 70	M 71 (1)	M 71	M 69	M 68	M 79
No	Maxillo-alveolar length	Maxillo-alveolar breadth	Palatal length	Palatal breadth	Bigonial breadth	Bicondylar breadth	Ht, of mandibular ramus	Max. breadth of mandibular ramus	Min. breadth of mandibular ramus	Ht. at mandibular symphysis	Mandibular length	Mandibular angle
AREA	G					v						*
41	52.5	_	45.0	48.0	_		_	_	_	-	. —	_
42	57.5	70.5	51.0	47.0	93.0	115.0?	56.5	43.5	32.0	35.0	83.0	120°
43	51.5	57.0	40.0	38.0		_	_	_		_	_	
44		57.0		32.0					·-	_	_	_
45	58.0	63.5	49.0	40.0		_	·		· <u> </u>	_		- .
46	55. 0	58.5	48.0	38.0		· —		_				
47	35.0	52.0	32.0	29.0	-	, , , , , , , , , , , , , , , , , , ,	_	_#	_	· _	¥	
48	43.0	57.0	34.0	33.0		_	·	_	,—	<u> </u>	_	
49	38.0	53.0	35.0	28.0	-	_	_			-		-
50		_	_	_		_		_		-	_	_
51	39.0	52.5	39.0	30.0		y y		_			<u> 9</u>	1 2

į	MOUND	AB		*		*							
	52	53.5	56.5	39:0	\$4.0	104.0	125.0?	58.0	42.0	31.5	27.5	78.0	113°
, 10	53	-	- 1 · 1	0.9	_		_		.500				-
4	54	55:0	63:5	48.0	41.0	84.0	110.0	60.0	42.0	35.0	27.0	78.0	127°
	55					_	******			-			named a

		M (1+8+17)									
		3	M 8/M 1	M 17/M 1	M 17/M 8	M 21/M 1	M 21/M 8	M 9/M 8	M 47/M 45	M 48/M 45	M 66/M 45
No.	Calculated cranial capacity in cc.	Cranial module	Length-breadth index	Length-height index	Breadth-height index	Length-auricular height index	Breadth-auricular height index	Transverse fronto-parietal index	Total facial index	Superior facial index	Jugo-mandibular index
AREA	G										
41	1377.94	151.33	79.12	70.33	88.89	59.89	75. 69	70.14		50.00	
42	1465.78	156.00	72.77	72.25	99.28	61.26	84.17	66.19		56.91	75.61
43	1355.57	148.67	78.41	75.00	95.65	65.34	8 3.33	71.74	_	48.24	
44	_	_	_	_	_	_				_	_
45	1270.89	145.33	77.14	72.00	9 3.33	62.86	81.48	68.15	_	56.54	_
46	1139.8 5	140.67	71.84	70.69	98.40	60.34	84.00	72.00	-	54.55	
47		134.00	75.90	66.27	87.30	60.84	80.16	(market	Name of Street		
48		_		68.39	_	58.62			-	50.94	
49			_	_					-	and the same of th	
50	_		80.62	MARKET THE	***************************************	-		64.34?			
51		128.67	66.87	65.66	98.20	59.64	89.19	74.77			

MO	UND	AR
TATA	OTAB	CATIO

52	1343.73	150.00	73. 08	74.18	101.50	62.64	8 5.71	69.71	86 .57	51.49	77.61
53		5,000,00	69.95?	ear-m			and an inter-	73.33	***************************************	-	professor.
54	1204.38	146.67	69.8 3	75. 98	108.80	60.89	87.20	74.40		-heroma	
55	orecond.	138.67	76.51	74.10	96.85	62.95	82.28	67.72			

D

			M 54/M 55 M		62/M 62 M	68/M 65 N	1 66/M 65	M 45/M 8	M 48/M 17	M 40/M 1	M 9/M 45
	M 52/M 51	M 52/M 51	M 54/M 55 M	61/M 60 M	63/W 02						2 2 2
Nø.	Orbital index (right)	Orbital index (left)	Nasal index	Maxillo-alveolar index	Palatal index	Mandibular index	Breadth index of mandible	Trans, craniofacial index	Vert. craniofacial index	Longitudinal craniofacial index	Jugo-frontal index
AREA	G										
41.	75.00	82.93	54.72	_	106.67	_		92.36	51.95	51.65	75.94
42.	82.05	77.22	49.52	122.61	92.16	72.17?	80.87?	88.49	50.72	50.26	74.80
43.	80.00	88.46	51.02	110.68	95.00	_	_	92.39	46.59	52.27	77.65
44.	80.00	81.69?	55. 9 5?	_		_		_		_	78.18
45.	76.19	79.01	48.94	109.48	81.63	_		87.78	53.17	53.14	77.64
46.	86.08		49.48	106.36	79.17	· —		96.80	53.66	52.01	74.38
47.	102.94	? —		148.57	90.62				38.18	43.98	_
48.	77.78	83.82	50.00	132.56	97.06				45.38		86.79
49.	86.36			139.47	80.00					47.74	? —
50.							-		_		_
51.	82.86	87.69	58.06	134.62	76.92	_	-	-	40.37		_

MOT	TIME	AR	

59	83.33	82,35	51.48	106.60	87.18	62.40?	83.20?	100.75	51,11	50.82	68.66
bj	3	-	_			**************************************		No.			
5	82.05	-	45.83	115.45	85.42	70.91	76.36	-	48 .53	55.87	_
	ß				sur-rest*		-		-	-	*****

Maxillary teeth

	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)
No.	Mesiodistal crown diam. Right M1	Labiolingual crown diam. Right M1	Mesiodistal crown diam. Left M1	Labiolingual crown diam. Left M1	Mesiodistal crown diam. Right M 2	Labiolingual crown diam. Right M2	Mesiodistal crown diam. Left M2	Labiolingual crown diam. Left M2	Mesiodistal crown diam. Right M3	Labiolingual crown diam. Right M3	Mesiodistal crown diam. Left M3	Labiolingual crown diam. Left M3
AREA	A G				9	The second secon						
41	11.5	11.5	_	_	10.0	11.5	_	_	7.5	10.5	8.5	7.5
42	_	_	10.0	11.0	10.0	12.0	10.5	11.5	8 .5	9.5	9.0	10.0
43	10.0	12.0	-	_	9.0	10.0	10.0	11.0	_	_	8.5	10.5
44	9.0	11.0	9.5	11.0	8.5	11.5	9.5	11.5	8.0	11.5	8.5	12.5
45	11.0	12.0	11.0	11.5	9.5	12.0	10.0	11.0	10.0	11.0		
46	11.0	11.0	9.0	11.0	9.0	11.0	11.0	11.0	_		_	
47	6.5	7.0	6.5	7.0	8.0	7.5	8.0	7.0	and the same of th	_	_	_
48	_		-				9.0	10.0	10.0	9.0	10.0	10.0
49	7.0	8.0	6.0	7.0			9.0	8.5		_		-
50	_							_		_	_	-
51	7.0	8.0	7.0	7.0	9.5	8.0	9.0	8.5	10.0	9.0	9 .5	8.5

	 100 000000
MOL	ΔR

52	9.0	11.0	9.0	10.5	*******				- '		attenting	-
53	10.0	9.0			8.5	10.0	_			-	erround,	_
54	11.0	11.0	11.0	11.0	9.0	10.0	11.0	9.0			******	****
55			7.0	8.0		*****	9.0	8.0	pathorm	- Times		and the same of

AREA	No.		Ĭ
Ω	Mesiodistal crown diam. Right M 1	M 81	ž
	Labiolingual crown diam. Right M 1	M 81 (1)	3
	Mesiodistal crown diam. Left M 1	M 81	a n
	Labiolingual crown diam. Left M 1	M 81 (1)	<u>م</u>
	Mesiodistal crown diam. Right M2	M 81	ь Б
	Labiolingual crown diam. Right M2	M 81 (1)	–
	Mesiodistal crown diam. Left M 2	M 81	7
	Labiolingual crown diam. Left M2	M 81 (1)	e e
	Mesiodistal crown diam. Right M3	M 81	0
	Labiolingual crown diam. Right M3	M 81 (1)	Þ.
	Mesiodistal crown diam. Left M3	M 81	
	Labiolingual crown diam.	× 81	

Left M3

| | | | | | | | | | | | | | |

10.0

1 | 1 | 1 | 1 | 10.0

| | | | | | 9.51

| | | | | | | | | | | | | | | | |

| | | | 9.0

| | | | | 9 |

ČII UT	57.4	5 3	52	MOUNI
1	11.5	1	1	AB
1	10.0			
	11.0			
1	10.0	1	e'	
T	9.0	1	Ì	
and the second	9.5	1	l,	
	10.0		1	
	9.0			
1	9.0	l	1 .	
1	9.0	1	1	
1	10.0	1	1	

9.0

	Cro	wn in	dex of M81(m a x i l l 1)/M 81	ary to	e e t h	Crow	n index		mandib (1)/M 81	ular	teeth	
No.	Right M 1	Left M1	Right M2	Left M2	Right M3	Left M3	Right M 1	Left M 1	Right M2	Left M2	Right M3	Left M3	
AREA G													
41	100.00	_	115.00	_	140.00	88.24	_		_	_	. —	_	
42		110.00	120.00	109.52	111.76	111.11	100.00	100.00	94.74	105.56	89.47	89.47	
43	120.00	_	111.11	110.00	_	123.53	_	_	_	_		_	
44	122.22	115.79	135.29	121.05	143.75	147.06	_	_	_		_		
45	109.09	104.55	126.32	110.00	110.00		_		_	_		_	
46	100.00	122.22	122.22	100.00	_					_			
47	107.69	107.69	93.75	87.50		_				_			
48		_	_	111.11	90.00	100.00	_			_			
49	114.29	116.67		94.44		-	_	_			_		
50		_	_	_	_		_		_	-	_	-	
51	114.29	100.00	84.21	94.44	90.00	89.47	_	-	_	_	_		

M	OI	TAT	Th	Λ	D
IVE		1131	11.7	-	

52	122.22	116.67	_			 -		patition (-	******	-
53	90.00	addressed	117.65	- Manager		 	named .	-			-
54	100.00	100.00	111.11	81.82		 86.96	90.91	105.56	90.00	100.00	90.00
55	Annania A	114.29		88.89	*******	 		-	-	-	Person

COLLECTIVE TABLE—C Continued

Linear measurements on craniograms

			-						
	W 42	W 43	W 44	W 48	W 49	W 50	W 51	W 52	W 53
No.	Frontal chord	Parietal chord	Occipital chord	Calvarial height	Lambda calvarial height	Bregma position line	Frontal perpendicular	Parietal perpendicular	Occipital perpendicular
AREA G									
41		_		_			_		_
42		_		_			_		_
43	108.0	117.0	99.5	106.0	73.0	97.0	26.0	25.5	26.5
44		_					_	_	_
45	108.5	118.0	98.0	112.0	76.0	99.0	27.0	29.0	28.5
46	106.5	112.0	87.0	104.5	70.0	93.5	26.5	26.0	20.5
47	_	_		_			_	_	
48				-		_			_
49		-					_	··	_
50		-	******				-	_	_
51	-	-		, <u></u>		`\	· _		9 3

TVIO	TOTAL	T TEN-	AR
TABAR	P C 1 1%	1107	AL BZ

52	117.5	111.0	96.5	110.5	72.5	101.0	24.5	27.5	30.5
53					_	-	_	_	
54-	110.0	109.5	_	107.0	68.0	97.0	23.5	22.5	
.55	Promo					*******			

COLLECTIVE TABLE — C Continued

Angular measurements on craniograms

	W 3	W 4	W 5	W 9	W 11	W 12 .	W 13	W 14	W 15	W 7
No.	Frontal inclination angle	Occipital inclination angle	Facial profile angle	Calvarial base angle	Frontal curvature angle	Parietal curvature angle	Occipital curvature angle	Occipital flexional angle	Superior facial length angle	Alveolar profile angle
AREA	G							ALCONOMICATOR STANCE AS Enclosed		carrotinis destructivos destructivos
41		_	_			_	_	_		_
42	_	_	_	_			_	_		
43	63°	85°	83°	12.5°	126°	133.5°	123°	123°	3 8°	7 0 .5°
44	_		_		8 <u> </u>		_	_		
45	66°	87.5°	79.5°	16.5°	126.5°	127°	118.5°	120°	43°	78.5°
46	61.5°	90°	77°	16°	126.5°	129°	127°	127°	43°	64°
47		_		_	-	_			-	_
48	_	_		_	-			_	_	
49		_				_	_	_		
5 0	_	, , , , , , , , , , , , , , , , , , ,	_				_	_		_
51			_	_	_	_		_		_

MOUN	ID AB									
52	5 9 .5°	82°	92°	13°	134°	127°	114°	114°	40°	116°
53				-		-		-	Water	******
54	62°	80.5°	7 9 .5°	16°	133.5°	134.5°			3 8 °	95 .5°
55	-				-			-	******	***

	S.4*			M 1	M 8	M 2a	M 17	M 9	M 21	M 25	M 24b
No.	Skl No.	Sex	Age	Maximum Cranial length	Maximum Cranial breadth	Nasion-inion length	Basion-bregma height	Minimum frontal breadth	Vertical porion height	Median sagittal arc	Vertical transversal arc
CEME	ETERY H (SURIAL)								
56	H184(K)	M	A	_	_	_			_		_
57	H307 (a)	M	A	189.0	142.0	176.0	133.0	99.5	_	391.0	<u></u> -
58	H484(a)	M	A				133.0		_		_
59	H 487 (a)	M	A	191.0		184.0	135.0	92.5	_	383.0	_
60	H 502 (G)	M	· A	189.0			_		_	_	
61	H 695	M	A	193.0	153.0?	176.0?	137.0	92.5	_	_	
62	H 698	M	A	183.5	139.0	170.0	136.0	98.5	119.0	372.0	318.0
63	H 88	F	A	194.0	_	_	_		_	_	
64	Н 306 (а)	F	A	188.0?						-,	_
65	H 488	F	A	183.5	132.0?	167.0	134.0	94.0	110.0	376.0	297.0
66	H 501 (a)	F	A								` _
67	Н 699	F	A	167.0	134.0	166.0	135.0	90.0	109.0		
68	H 710	F	A	177.0	_	163.0	_	95.0			_
CEM	ETERY H (JAR BU	JRIAL)								
69	H 206 (B)	M	A	180.0?	138.0	Manag	17.				010.0
70	H 255 (a)	M	A .	198:0 ^h	135.0	180.0	134.0	00 -			312.0
71	H 344	M	A	184.0?	141.0	165.0	136.0	98.5	118.0	384.0	307.0
72	H 61	F	A		131.0	Ministra	136.0	95.0 104.0	119.0	379.0	316.0
73	H 153 (a)	F	A	181.0		173.0	117.0?		117.0	0550	304.0
								89.0		355.0	*******

.

	M 23	M 40	M 48	M 47	M 45	M 55	M 54	M 50	M 51	M 51	M 52	M 52
No.	Horizontal circumference	Prosthion-basion line	Nasion-prosthion line	Nasion-gnathion line	Bizygomatic breadth	Nasal height	Nasal breadth	Ant. interorbital breadth	Orbital breadth (right)	Orbital breadth (left)	Orbital height (right)	Orbital height (left)
CEMET	ERY H	(OPEN I	BURIAL)						,	V		
56	-		72.0		_	53.5	26.5	22.0	41.5	_	-	_
57	537. 0			_	-	53.0		20.5	45.0		34.0	
58	_	100.0		_		_	27.5?	_		44.5	-77	34.0
59	72.00	94.0	70.5			55.0	25.0	18.5	39.0	_	31.0	_
60	* <u>-</u>	100.0	66.0	116.0	_	47.5	21.0?	19.0?	_	37.5	_	36.5
61	540.0	95.0	75.0			54.5	28.0?	18.5	43.0	40.5	35.0	34.5
62	514.0	94.0	68.0	116.5	136.0	52.5	25.0	20.0	41.0	40.0	33.0	33.0
6 3	· _	_	69.0	121.5		49.5	26.0	21.0	-	37.5		35.5
64		_	_		_	46.0		18.0	38.0	42.5	32.5	31.0
65	505.0?	·-			_	39.0	26.0	18.0	42.0	40.0	31.0	30.0
66	·	100.0	71.0	121.0	120.0?	51.0	21.0	19.0	40.5	41.5	36.0	
67	_	100.0	58.0	106.0	115.0	43.5	23.0	15.0	41.5	41.5		38.0
68	492.0			-					- Te (e)			
		¥						•				
CEMET	ERY H	(JAR BU	JRIAL)									
69	540.0	_			_	-		- Section 1			et man	-
70	545.0	107.0	71.0	123.0	131.0	51.0	27.0	22.0	42.5	43.5	34.0	33.5
71	522.0	98.0	60.0		136.0	47.0	27.0	22.0	39.0	40.0	30.5	30.0
72	483.0	<u> </u>		NAME OF THE PARTY	-		manuf.		42.0	-	34.0	the design
73				-	126.0	40.0?	22.0	15.0		- Thomas	Philips	No.
10	500.0											

E

	M 60	M 61	M 62	M 63	M 66	M 65	M 70	M 71 (1)	M 71	M 69	M 68	M 79
No.	Maxillo-alveolar length	Maxillo-alveolar breadth	Palatal length	Palatal breadth	Bigonial breadth	Bicondylar breadth	Ht. of mandibular ramus	Max. breadth of mandibular ramus	Min. breadth of mandibular ramus	Ht. at mandibular symphysis	Mandibular length	Mandibular angle
CEMET	ERY H	(OPEN B	URIAL)									
56	52.0	62.0	48.0?	32.5	_	_	_	_	_	_	_	_
57		_	_		_	_	_	_	_		_	_
58	57.0	67.0?	49.0	41.0			_		_	_	_	_
59	57. 0	63.0	43.5	39.5		_	_	_	_		_	_
60	58.5	65.5	47.5	37.0	82.5	-	60.0	39.5	34.0	32.0	91.0	110°
61	56.0	75.0		48.0			_	_		_	_	_
62	50.0	62.0	46.0	36.5	75.5	112.5?	64.0	40.0	·36.0	28.5	83.0	111°
6 3	58.0	62.5	48.5	33.0	_			37.0	30.0	34.0	75.0	_
64		59.0		36.0	_		58.0	42.5	29.0	31.0	81.0	123.5
65			44.0?	34.5	84.0		54.5	45.0	35. 0	29.5	74.0	121°
66	57.5	62.5	48.0	37.5	83.5	112.0	56.0	42.0	35.5	30.0	81.5	125°
67	_		Marine		79.0		_	_	30.0	29.0	70.0	_
68	_					_	_		_		_	_
CEME'	TERY H	(JAR BU	RIAL)									
69	50.0	62.0	42.0	39.0		-		-		32.5	-	
70	56.0	63.0	-4	41.0	100.0	111.0	62.0	46.0	34.5	24.0	88.0	117°
71	56.0		47.0		74.5				35.0	31.5	80.5	125°
72	. Marine	name.	44.0	parameter		name of the last o	******		en en	-	*******	Named
73	-	-				_	Principal				-	

	,	M (1+8+1)	7) М 8/М 1	M 17/M 1	M 17/M 8	M 21/M 1	M 21/M 8	в м9/м	B M 47/M 45	M 48/M 45	M 66/M 4
No.	Calculated cranial capacity in cc.	Cranial module	Length-breadth index	Length-height index	Breadth-height index	Length-auricular height index	Breadth-auricular height index	Transverse fronto-parietal index	Total facial index	Superior facial index	Jugo-mandibular index
EME	TERY H	(OPEN BU	RIAL)								-
56	_	_		-				_			
57	1474.07	154.67	75.13	70 .37 .	9 3.66		_	70.07			
58	-	_						_		_	
59		_	_	70.68	_	_			_	***************************************	
60	. —	_							_	encoder.	element .
61	1600.70?	161.00?	79.27?	70.98	89.54?		Name of Street, Street	60 .46?	_	matte	etoeno.
62	1440.81	152.83		74.11	97.84	64.85	8 5.61	70.86	85.66	50.00	55.51
63	_	_	75.75			- Control of Control o		_			-
64							_	-	_		
6 5	1298.21?	149.83?	不	73.02 1	01.52?	59.95	8 3.33?	71.21?			-
66	-	_	71.93?			-	-	-	100.83?	59.17? 6	9.587
67	1204.52	145.33	80.24	80.84	100.75	65.27	81.34	67.16	92.17	50.43 6	8.70
68		1.1	_				_		_	_	_

CEMETERY H (JAR BURIAL)

69			76.67?	-		-		-		-	
70	1482.65	155. 6 7	68.18	67.68	99.26	59.60	87.41	72.96	98.89	54.20	76.34
71	145 9 .13	153.67	76.63?	73.91?	96 .45	64.67?	84.40	67.38		44.12	54.78
72			-	_	103.82		89 .31	79.39	-	Pro-	-
73		,,,,,,	-	64.64?	-	-		-	*****	************	_

*	M 52/M 51	M 52/M 51	M 54/M 55	M 61/M 60	M 63/M 62	M 68/M 65	M 66/M 65 N	1 45/M 8	M 48/M 17	M 40/M 1	M 9/M 45
No.	Orbital index (right)	Orbital index (left)	Nasal index	Maxillo-alveolar index	Palatal index	Mandibular index	Breadth index of mandible	Trans. craniofacial index	Vert. craniofacial index	Longitudinal craniofacial index	Jugo-frontal index
CEMET	ERY H (OPEN BU	JRIAL)								
56		_	49.53	119.23	67.71?	_		_			_
57	75.56				_		_		-	_	
58		76.40		117.54?	83.67		-	_		_	_
59	79.49	-	45.45	110.53	90.80		_	_	52.22	49.21	
60	Menderal	9 7.33	44.21	111.97	77.89	_	_		-	52.91	_
61	81.40	85.19	51.38?	133. 9 3	parameter.	_	_		54.74	49.22	
6 2	80.49	82.50	47.62	124.00	79.35	73.78?	67.11?	97.84	50.00	51.23	72.43
6 3	-	94.67	52.53	107.76	68.04		_			_	·
64	8 5.53	72.94		_	horizon	Message				-	_
65	73.81	75.00	66.67	_	78.41?	_	_		_		
66	88.89		41.18	108.70	78.12	72.77	74.55				_
67		91.57	52.87	t-maker.		_		85.82	42.96	59.88	78.26
68			_			-	-		_	_	_
СЕМЕ	TERY H	(JAR BI	JRIAL)								
69		. Comments	,	124.00	92.86			-			-
70	80.00	77.01	52.94	112.5 0	Shain	79.28	90.09	97.04	52.99	54.04	75.19
71	78.21	75.00	57.45				-	96.45			
72	80.95		01740	gaspeer	-	****	Titellow	30.10	-11.01		Monaged
73		-	55.00?	******	guarra		Mining				70.6

			M a	x i l	l a	r y	t	e e	t h			
	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)
No.	Mesiodistal crown diam. Right M 1	Labiolingual crown diam. Right M 1	Mesiodistal crown diam. Left M 1	Labiolingual crown diam. Left MI	Mesiodistal crown diam. Right M2	Labiolingual crown diam. Right M2	Mesiodistal crown diam. Left M 2	Labiolingual crown diam. Left M 2	Mesiodistal crown diam. Right M3	Labiolingual crown diam. Right M3	Labiolingual crown diam. Left M3	Mesiodistal crown diam. Left M3
CEM	ETERY	H (OPE)	N BURIA	L)								
56	9.5	11.0	9.0	10.5	9.0	10.5	9.5	10.5	8.5	9.5	_	,
57	_	_	_	_	_			_	_	_	. —	
58	10.0	11.0	10.0	11.0	9.0	10.5	10.0	10.5	9.5	9.0	9.0	10.5
59	10.5	12.0	9.5	12.0	8.5	11.0	9.0	11.5	7.5	10.5	9.0	11.0
60	10.0	12.0	10.0	11.5	9.5	10.5	9.0	10.0	9.0	10.0	8.0	10.5
61	11.0	10.5	11.5	11.0	11.0	10.0	9.5	10.0	9.5	10 .5	9.5	8.5
62	10.5	11.0	10.0	11.0	8.5	11.0	8.5	10.5	8.5	10.5	8.5	10.0
-63	10.0	10.5	10.0	10.0	9.0	11.5	8.0	11.0	9.0	10.0	_	_
64	9.0	11.5			10.5	10.0			10.0	10.0	-	-
65	_	_			_	-	10.0	9.0		_	8.0	9.5
66	10.5	11.5	11.0	11.5	9.0	11.0	10.0	10.0	8.0	9.0	8.0	9.0
-67	-	_	-	-	eares	Managa	en en	-	_			
68	_						_				-	
			-									
CEN	IETERY	H (JAR	BURIAL)								
69	11.0	12.5	10.5	12.0	9.5	12.5		wants .	8.5	11.5	10.5	11.5
70	11.0	12.5		Wester	11.0	11.5	10.0	12.0	-		9.0	11.0

71

72

73

10.5 11.5

72 73

		М	a n	d i	b u	1	a r	t e	e e	t h		
	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)	M 81	M 81 (1)
No.	Mesiodistal crown diam. Right M 1	Labiolingual crown diam. Right M 1	Mesiodistal crown diam. Left M 1	Labiolingual crown dlam. Left M1	Mesiodistal crown diam. Right M 2	Labiolingual crown diam. Right M 2	Mesiodistal crown diam. Left M2	Labiolingual crown diam. Left M2	Mesiodistal crown diam. Right M3	Labiolingual crown diam. Right M 3	Mesiodistal crown diam. Left M3	Labiolingual crown diam. Left M3
CEM	ETERY	H (0P)	EN BURI	AL)								
56		-	_	_	_	_			10.0	8.5	_	_
57		-				_	_	_	_	_	_	_
58	10.5	9.5	11.0	10.5	10.5	10.0	11.0	9.5	_	_	10.5	9.5
59		_							_	-	_	_
60	10.5	9.5	10.5	10.0	9.5	9.0	10.0	9.0	10.0	8.5	10.0	9.0
61	_	_	_	-	_		_	_	-		_	
62	11.0	10.5	11.0	10.5	9.5	9.0	9.0	9.0	10.0	9.5	11.0	8.5
6 3	10.5	10.5	10.0	10.0	9.5	10.5	10.0	9.5	10.0	10.0	10.0	9.5
64		_			_						-	_
6 5	-	_		_	10.0	10.0						
66	10.0	10.0	10.5	10.5	10.0	9.0	9.0	10.0	_		9.0 `	8.5
67		-				_						_
68	-			-		annua .				_	Name of Street, Street	_
			-					PRODUCTION CONTRACTOR				
CEM	ETERY	H (JAF	R BURIA	L)								
69	11.5	11.0	12.5	11.5	10.5	11.0	11.0	11.5	10.5	11.0	9.5	10.5
70	-					agencies	*****	-	-	-		
71	10.5	11.5	mage	*****	41.5	10.5			_			

- (1.5 10.5 -

11.0

12.0 - 9.0 11.0 - -

	Cr	own in	ndex of M81	maxi (1)/M 81	llary	teeth	Cro	wn inde		mandib (1)/M 81	ular	teeth
No.	Right M 1	Left M 1	R. at M2	Left M2	Right M3	Left M3	Right M1	Left M 1	Right M2	Left M2	Right M3	Left M3
CEMET	rery h	(OPEN	BURIAL)						`			
56	115.79	116.67	116.67	110.53	111.76	_		_	_		85.00	_
57	_	_	_	_	<u> </u>	_	_		_			_
58	110.00	110.00	116.67	105.00	94.74	116.67	90.48	95.45	95.24	86.36	_	90.48
59	114.29	126.32	129.41	127.78	140.00	122.22	_	_	_		-	
60	120.00	115.00	110.53	111.11	111.11	131.25	90.48	95.24	94.74	90.00	85.00	90.00
61	95.45	95.65	90.91	105.26	110.53	89.47	_		_			
62	104.76	110.00	129.41	123.53	123.53	117.65	95.45	95.45	94.74	100.00	95.00	77.27
63	105.00	100.00	127.78	137.50	111.11		100.00	100.00	110.53	95.00	100.00	95.00
64	127.78	_	95.24	_	100.00	-			*******	_		
65	_	_	-	90.00		118.75	_ ,		100.00	_		-
66	109.52	104.55	122.22	100.00	112.50	112.50	100.00	100.00	90.00	111.11		94.44
67		_						_			-	-
68	_			_					_			
			М									
CEMET	rery h	(JAR B	URIAL)									
69	113.64	114.29	131.58		135.29	109.52	95.65	92.00	104.76	104.55	104.76	110.53
70	113.64		104.55	120.00	_	122.22	Marketon				_	_
71						-	109.52	-	91.30			_
72		***			enperint	Amoun	-		www	Nones		_
73		******				torus.	Statements		109.09		122.22	2

Linear measurements on craniograms

	W 42	W 43	Linear n	.W 48	on cran W 49	w 50	W 51	W 52	W 53
				1					
No.	Frontal chord	Parietal chord	Occipital chord	Calvarial height	Lambda calvarial height	Bregma position line	Frontal perpendicular	Parietal perpendicular	Occipital perpendicular
CEMETER	Y H (OPEN	N BURIAL)			y /		8	
56			_	_			_		_
57	118.5	110.5	110.0	109.5	72.5	103.5	28.0	19.5	34.5
58	_	11 —	_	_			_		
59	_	_	_	·		· <u></u>	· _	_ ·	_
60			_	* 			- <u>-</u>		_
61	_					_	_	· <u> </u>	-
62	113.5	105.5	110.0	108.5	66.5	99.0	22.5	20.5	31.5
63	_		_	_	_	_	_	_	_
64	**************************************		_		_	_	·	_	
65	110.0	108.0	101.5	107.5	69.5	97.5	24.0	24.5	38.0
66	-		_	-		1	_	_ `	
67	_	_				_			-
68		_	_	_					
							-		
CEMETER	Y H (JAR)	BURIAL)							
69			promotes				_	- Marine III	
70	107,5?	116.5?	103.0	114.0	64.5	93.5?	23.5	29.0	33.0
71	114.5	Acres 100 and	200	110.0		100.0	28.0		-
72							No.	Principal Princi	
73	-	energy.		manage .					Proceeding

Angular measurements on craniograms W 3 W 4 W 5 W 9 W 11 W 12 W 13 W 14 W 15 W 7 Superior facial length angle Occipital inclination angle Occipital curvature angle Frontal inclination angle Parietal curvature angle Occipital flexional angle Frontal curvature angle Alveolar profile angle angle profile angle Calvarial base Facial No. CEMETERY H (OPEN BURIAL) 56 61° 81.5° 9° 128° 139° 114.5° 114.5° 57 58 59 60 61 89° 10.5° 136.5° 137° 120.5° 121° 40° 100° 81.5° 62 60.5° 63 6475.5° 82° 130.5° 130° 104° 110° 32° 79° 62.5° 65 66 67 68 CEMETERY H (JAR BURIAL) 69 113° 114° 11° 132° 125° 3**8**° 85° 78° 80.5° 61° 70 8° 123.5° 33.5° **8**3° 68° 60.5° 71 72 73

				M 1	M 8	M 2a	M 17	М 9	M 21	M 25	M 24b
No.	Skl No.	Sex	Age	Maximum Cranial length.	Maximum Cranial breadth	Nasion-inion length	Basion-bregma height	Minimum frontal breadth	Vertical porion height	Median sagittal arc	Vertical transversal arc
CEME	ETERY H (JA	R BUI	RIAL)						<		
74	H 154 (a)	F	A	176.0	128.0	166.5	139.0	85.0		362.0	_
75	H 206 (d)	F	A	173.0	133.0	165.0	125.0	93.0	_	344.0	_
76	H 245 (e)	F	AII	175.0	147.0	_	_		_	_	_
77	H 246 (c)	F	Α	182.0	135.0	167.0	135.0	96.0	111.0	371.0	298.0
78	H 247 (a)	F	Α	176.0		_	130.0		108.5	_	282.0
79	H 247 (b)	F	A	178.0	133. 0	·	125.5	87.5	108.0	353.0?	292.0
80	H 7435 (b)	F	A	178.0	129.0?	165.0	117.0	91.0	100.0	350.0	283.0
81	Pot no. 11	F	A	173. 0	131.0	166.0	122.0	92.0	103.0	352.0	284.5
82	Pot no. 12 (a)) F	A	173.0	131.0	162.0	119.0	90.0	107.5	360.0	289.0
83	Pot no. 12 (b) F	A		119.0		124.0	-	_	-	285.0
84	H 206 (a)	F	Juv.			_		_	_		` _
85	H 245 (a)	F	Juv.	167.0	128.0	160.0	124.0	83.0	105.0	335. 0	284.0
86	H 231 (c)		Infant	_	-				_		_

	M 23	м 40	M 48	M 47	M 45	M 55	M 54	M 50	M 51	M 51	M 52	M 52
No.	Horizontal circumference	Prosthion-basion line	Nasion-prosthion line	Nasion-gnathion line	Bizygomatic breadth	Nasal height	Nasal breadth	Ant. interorbital breadth	Orbital breadth (right)	Orbital breadth (left)	Orbital height (right)	Orbital height (left)
CEM	ETERY	H (JAR	BURIA	L)								
74	493.0	_	_	_	-	54.0	_	16.5	35.0	37.0	-	_
75	491.0	91.0	54.0	96.0?	_	43.0	28.0	18.5	_	41.5	35.0	31.5
76			-	_			25.5			_		
77	573.0		-	_	_			20.0	39.0	39.0	33.0	
78	488.0	100.0	63.5	_	119.0	48.0	26.0	21.0	41.5	_	33.0	_
79	504.0	96.5 ,	60.0			46.0	26.0	17.5	39.0	40.0	32.5	32.0
80	499.0	100.5?	57.0		_	44.0?	23.5	18.5	39.0?	41.5	_	30.0
81	494.0	89.5	65.0		125.0	49 .5	24.0	22.0	40.0	39.5	30.0	34.0
82	488.0	85.0	56.5	_	116.0	45.0	24.0	18.5	39.0	34.0	31.5	30.0
83		_			_		_	_				
84	_	_	uned)		_					-	operations.	
85	478.0	90.0	60 .5		117.0	48.5	25.0	16.0	39.0	39.0	34.0	_
86			36.5		specimo	28.0	19.0	15.0	32.0	31.0	28 .5	26.0
		110										

	M 60	M 61	M 62	M 63	M 66	M 65	M 70	M 71 (1)	M 71	M 69	M 68	М 79
No.	Maxillo-alveolar length	Maxillo-alveolay kreadth	Palatal length	Palatal breadth	Bigonial breadth	Bicondylar breadth	Ht. of mandibular ramus	Max. breadth of mandibular ramus	Min. breadth of mandibular ramus	Ht. at mandibular symphysis	Mandibular length	Mandibular angle
CEMETI	ERY H (AR BUR	RIAL)									
74			_	_	_		57.0	41.5	34.0		88.0	123°
75		_	_	_	_	_	50.5	39.0	30.0	21.0	81.0	123°
76			_		_	_		_	_	25.0		_
77	_					_	_		_	ė 		_
78	60.0?	63.5?	42.0	_	_	_	_	_		_	_	_
79	-	-	39.0		_	_		_			_	_
80	57. 0	64.0	_		_	_	_		34.0	_		_
81	56.0	64.5	_	_	_	_		_	_	_	_	_
82	49.0	64.0	41.0?	35.0?	_		_	_	_		_	
83	-		-		_		_	_	-			_
84		_		-	75.5	103.0	41.0	39.0	31.0	_	72.0	123°
85	46.5	62.5	37.0	35.0	84.5	_	-	35.0	31.0	26.0	74.0	121°
86				_	53.5	80.5	24.5	24.0	21.0	19.5	46.5	133°

		M (1+8+17)									· · · · · ·
		3	M 8/M 1	M 17/M 1	M 17/M 8	M 21/M 1	M 21/M 8	M 9/M 8	M 47/M 45	5 M 48/M 45	M 66/M 45
No.	Calculated cranial capacity in c.c.	Cranial module	Length-breadth index	Length-height index	Breadth-height index	Length-auricular height index	Breadth-auricular height index	Transverse fronto-parietal index	Total facial index	Superior facial index	Jugo-mandibular index
CEME	TERY H	(JAR BU	RIAL)								
74	1300.50	147.67	72.73	78.98	108.59	_	÷	66.41			8
75	1260.68	143.67	76.88	72.25	93.98	_	_	69.92	_	_	
76	_	_	84.00	_	_	_				_	specialis
77	1324.35	150.67	74.18	74.18	100.00	60.99	82.22	71.11		_	
78	-	_	_	73.86		61.65	_		_	53.36	-comm
79	1253.44	145.50	74.72	70.51	94.36	60.67	81.20	65.79			
80	1145.03	141.33?	72.47?	6 5.73	90.70?	56.18	77.52?	70.54?			-
81	1160.87	142.00	75.72	70.52	93.13	59.54	78.63	70.23	_	52.00	_
82	1203.30	141.00	75.72	68.79	90.84	62.14	82.06	68.70	_	48.71	_
83	_		-	_	104.20	-	name of the same o			_	
84	_									_	
85	_	139.67	76.65	74.25	96.88	62.87	82.03	64.84	_	51.71	72. 22
86	_		_		_			_			

	M 52/M 51	M 52/M 51	M 54/M 55	M 61/M 60	М 63/М 62	M 68/M 65	M 66/M 65	M 45/M 8	M 48/M 17	M 40/M 1	M 9/M 45
No.	Orbital index (right)	Orbital index (left)	Nasal index	Maxillo-alveolar index	Palatal index	Mandibular index	Breadth index of mandible	Trans. craniofacial index	Vert. craniofacial index	Longitudinal craniofacial index	Jugo-frontal index
CEMET	ERY H	(JAR BU	RIAL)								
74	_		_	_	-	_	_	_	_	_	_
75	_	75.90	65.12	_	-	_	_		43.20	52.60	_
76	_	_	_	_	_	_		_	_		_
77	84.62	_	-	_			_		_		_
78	79.52	_	54.17	105.83?	_	_	_		48.85	56.82	_
79	8 3.33	80.00	56.52	_	_	_	_		47.81	54.21	_
80		72.2 9	53.41?	112.28			_	_	48.72?	56.46?	
81	75.00	86.08	48.48	115.18	_	-		95.42	53.28	51.73	73.60
82	80.77	88.24	53.33	130.61	8 5.37?	_		88.55	47.48	49.13	77.59
83		-	_						_		
84	_	_				69.90	73.30		_	· <u>-</u>	_
85	87.18	_	51.55	134.41			-	91.41	48.79	53. 89	70.94
86	89.06	83.87	67.86		*****	57.7 6	66.46	_	_	_	

	85	84	83	82	81	80	79	78	77	76	75	74	No.	
	10.0	1	1	l	11.0	I	1	١	l	1	1	11.0	Mesiodistal crown diam. ≤ x	
	11.5	l	I	١	11.0	ı	1	I	l	ł	I	12.0	Labiolingual crown diam.	
	10.5	١	l	1	10.5	I	1	ı	I	ı	I	1	Left M 1	M a
	11.0	1	1	1	11.5	1	ı	ı	I	1	ı	I		×
	10.5	I	1	1	1	ı	f	ı	ı	ı	I	9.5	Mesiodistal crown diam.	_ _ a
	12.0	Ĺ	l	ı	ı	1	1	1	1	ı	1	11.0	Labiolingual crown diam.	r
	10.0	1	ſ	1	10.0	1	ı	ı	1	1	1	1	Mesiodistal crown diam. ≤ Left M 2	·
	11.5	I	ı	1	11.0	I	ı	I	1	I	1	1	Labiolingual crown diam.	e e
	1	1	-	١	j	I	ı	4	l	1	1	I	Mesiodistal crown diam. ≥ Right M3	· -
l	1	1	1	ſ	1	Å	1	1	ı	ı	ı	1	Labiolingual crown diam.	
1	1	I	Ī	Ţ	9.0	A	I	ı	1	I	ı	I	Mesiodistal crown diam.	
1	Í	I	I		10.0			1	1	I	1	1	Labiolingual crown diam.	

86

88	80	84	83	82	81	80	79	78	77	76	75	74	CEME	No.		xlviii
ł	1 8	1	1	ı	1	I	ı	1	I	l	Ī	10.5	CEMETERY H	Mesiodistal crown diam. Right M1	M 81	
1	1	I	1	I	1	1	1	1	I	I	1	10.0	(JAR BURIAL	Labiolingual crown diam. Right M1	M 81 (1)	3
1	ł	ı	1	ı		1	Į.	1	I	l	10.0	11.0	URIAL)	Mesiodistal crown diam. Left M1	M 81	ם
l	1	I	1	1	1	1	t	L	I	1	11.0	9.5		Labiolingual crown diam. Left M1	M 81 (1)	COLLEC
1	1	1	1	I	1	1	I	1	ı	1	I	9.0		Mesiodistal crown diam. Right M2	M 81	CTIVE T
1	1	ı	1	-	1	1	I	1	1	I	I	Ø 7.7.		Labiolingual crown diam. Right M2	M 81 (1)	ABLE —
I	I,	l	I	l	1	1	1	1	I	1	9.0	10.0		Mesiodistal crown diam. Left M2	M 81	E Conti r
1	i	l	ŀ	-	I	1	I	I	I	ı	9.5	8.51		Labiolingual crown diam. Left M2	M 81 (1)	in u e d
1	l	l	I	ł	1	l	1	I	1	1	ı	9.5		Mesiodistal crown diam. Right M3	M 81	O C
1	I	1	1	. 1	1	1	1	1	ı	1	I	9.0		Labiolingual crown diam. Right M3	M 81 (1)	5
I	1	1	ı	ı	ı	ı	1	i	1	1	9.5	1		Mesiodistal crown diam. Left M3	M 81	
ı	I	1	1	1	1	1	1	1	l	l	9.5	I		Labiolingual crown diam. Left M3	M 81 (1)	

	Cro	own in	index of maxillary M 81 (1)/M 81			teeth	Cro	wn inde		mandil (1)/M 81	ular	teeth
						1						
No.	Right M 1	Left M1	Right M2	Left M2	Right M3	Left M3	Right M1	Left M1	Right M2	Left M2	Right M3	Left M3
EMET	rery h	(JAR BU	URIAL)									
74	109.09		115.79	_			95.24	86.36	94.44	85.00	94.74	
75	_			_	_	_	_	110.00		105.56		100.0
76		_ ,	- ,		, , -	_	_			_		
77	_	-			_		_	_		_		_
78	,	_	_							_		-
79	_		_	_			_		_			
80	_		_					_				_
81	100.00	109.52	_	110.00	_	111.11	_					
82		_	_		_			_			-	_
83				_							-	_
84	_					_				_	-	
85	115.00	104.76	114.29	115.00		_	-	_			-	-
86			_							_		Manage

Linear	measurements	on	craniograms
Linear	measurements	on	Crainograins

		W 42	W 43	W 44	W 48	W 49	W 50	W 51	W 52	W 53
No.		Frontal chord	Parietal chord	Occipital chord	Calvarial height	Lambda calvarial height	Bregma position line	Frontal perpendicular	Parietal perpendicular	Occipital perpendicular
CEMETE	RY	H (JAR	BURIAL)							
74		_		_	_	_		_	_	_
75		_	-		_	_	_	- '	_	_
76		_	- 1	_	_	· —	_	_	_	_
77		110.0	106.0	103.0	105.0	65.0	96.0	26.0	19.0	32.0
78		_	_		_	_			_	
79		107.5	111.5	_	_	70.0	_	27.5	25.0	_
80			_	-	_		_	_	_	
81		110.0	106.0	94.5	100.5	67.5	93.0	28.0	23.0	31.5
82						_	_	_		
83		_		_			-	_	_	-
84			* *		_		_	_		
85			5ee	7.0	-		_	_	_	
86		_	91 H		-		. <u> </u>	_	_	_

COLLECTIVE TABLE — E Continued

Angular measurements on craniograms

	W 3 W 4		W 5	W 9	W 11	W 12	W 13	W 14	W 15	W 7		
No.	Frontal inclination angle	Occipital inclination angle	Facial profile angle	Calvarial base angle	Frontal curvature angle	Parietal curvature angle	Occipital curvature angle	Occipital flexional angle	Superior facial length angle	Alveolar profile angle		
CEMETERY H (JAR BURIAL)												
74		_	_		_		_		_	_		
75		-	_	-		_	_	_	_			
76		-	_		_			_				
77	61°	82°		12.5°	128.5°	140°	115.5°	114.5°		_		
78	_		_		_	_	-		_			
79	_	_	84.5°	_	125.5°	131°	_	_	36.5°	88 .5°		
80	_								_	name.		
81	58°	83°	82.5°	14°	125°	133°	112.5°	115°	41.5°	67°		
82	_	_			_		same.		_	_		
83		_		-		_						
84	_	_			_		_	_				
85	_	_		_	_	_			_	_		
86				Name of Street	_							

-					M 66	M 65	M 70	M 71 (1)	M 71	M 69	M 68	M 79	M 68/M 65	M 66/M 65
No.		andible No.	Sex Ag		Bigonial breadth	Bicondylar breadth	Ht. of mandibular ramus	Max. breadth of mandibula r ramus	Min. breadth of mandibular ramus	Ht. at mandibular symphysis	Mandibular length	Mandibular angle	Mandibular index	Breadth index of mandible
CEM	ET	ERY R 37												A CONTRACTOR OF THE SAME OF TH
1	Н	784/A	M	A					31.0	28.0	_	_	_	_
2	Н	793/A	Μ	A	60.0	_			39.0		_			_
3	Н	802	М	A		_	_	_	_	32.0?		_		_
4	Н	815	M	A	98.0	_	84.0	51.0	37.0	38.0	83.0	111°		_
5	Н	801/B (III)	Μ	Juv.	_	-	-		30.0	29.0	74.0?	_	_	-
6	Н	798/A3	F	A	83.0	_			29.0	30.0	76.0	_	- ,	_
ARI		G	M	A	95.0	116.0	60.0	40.5	29.0	29.0	85. 0	123°	73.28	81.90
						~		was an annual section of the section						
CEN	/IE'	TERY H (O	PEN	BURIA	L)									
8		I 700		A		111.0	62.5	41.0	31.0	31.5	67.5	121°	60.81	79.28
9	Н	485		Juv.		-				25.0	74.0			
.Con							K =	4						
10		TERY H (J.		URIĂL)									
10	ŀ	^H 206 (c)	F	A	86.0	******	47.0?	38.0	30.0	24.0	76 .5	133°	Names	

10	CEMI	<i>©</i> &	CEMI	7	AREA	6	σι	4	ယ	12	р-1	CEMI	No.		
ı	CEMETERY	11.0	CEMETERY	10.0	G	1	11.0	1	[11.0		CEMETERY	Mesiodistal crown diam. Right M 1	M 81	
l	H (JAR BURIAL)	9.0	H (OPEN	9.0		I	10.0	1	1	10.0	I	R 37	Labiolingual crown diam. Right M1	M 81 (1)	×
1	JURIAL)	11.0	BURIAL	10.5			11.0	I	1	11.0	I		Mesiodistal crown diam. Left M 1	M 81	a
1		© :: : : : : : : : : : : : : : : : : :	C				10.0	I	1	10.5	1		Labiolingual crown diam. Left M 1	M 81 (1)	<u>-</u> .
1		10.0		10.0		1	10.0	1	10.5	10.0	1		Mesiodistal crown diam. Right M2	₩ 81	<u>Б</u>
1		7.0		1			9.0	1	9.0	10.0	1		Labiolingual crown diam. Right M2	M 81 (1)	_ a
1		10.0		10.0			10.0	1	1	10.0	1		Mesiodistal crown diam. Left M2	M 81	٦
ı		88		.53		1	8.0	1	1	10.0	1		Labiolingual crown diam. Left M2	M 81 (1)	e e
I		1 1		51		1	1	1	10.0	10-5	I		Mesiodistal crown diam. Right M3	M 81	e c
1		1	I	9.0		I	-	1	9.0	9.5	1		Labiolingual crown diam. Right M3	M 81 (1)	5
4		1	1	S			1	1	Оп	9.5			Mesiodistal crown diam. Left M3	M 81	
İ		1		9.0		1	1	1	1	9.0	1		Labiolingual crown diam. Left M3	M 81 (1)	

(Crow	n inde		mandib 1)/M 81	ular	teeth
ř.	M 1	-	M 2	61	13	က
No.	Right N	Left M	Right 1	Left M2	Right M	Left M3
CEMET	ERY R	37				
1					_	
2	90.91	95.45	100.00	100.00	90.48	94.74
3		_	85.71	-	90.00	
4		-	_	_	_	_
5	90.91	90.91	90.00	80.00	_	_
6			_	_	-	_
AREA 7	G 90.00		_	95.00	94.74	94.74
CEME	TERY H	(OPEN	BURIAL	·)		
8				matina		
9	81.82	77.27	70.00	80.00	-	-
	-		*			
CEMIE	TERY H	(JAR	BURIAL)	******		_

					Cra	nial conto	ur in nor	ma vertic	alis			al aspec	t of the aticus
No.	Skl No.	Sex	Age	Ellipsoides	Pentagonoides	Rhomboides	Ovoides	Sphenoides	Sphaeroides	Byrsoides	Phaenozygy	Orthozygy	Cryptozygy
CEM	ETERY R	37											
1	H 779 (e)	M	A	+							+		
2	Н 793	M	A					+			+		
3	H 793 (A)	M	A				+				+		
4	H 793 (B)	M	A				+						
5	H 794	\mathbf{M}	A					+			+		
6	H 796 (B)	M	A					+			+		
7	H 798 (A)	${\bf M}$	A					+			+		
8	H 798 (B)	M	A				+						
9	H 798 (C)	M	A					+					
10	H 806	M	A	+							+		
11	Н 811	M	A	+									
12	Н 818	M	A	+							+		
13	H 820 (iii)	M	A					+			+ +		
14	Skl. 1	M	A							+	T	+	
15	Skl. 10	M	A					+					
16	H 801 (B)	M	Juv.								+		
17	H 779 (a)	F	A	+							+		
18	H 779 (c)	F	A	+									
19	H 779 (d)	F	A									+	
20	Н 780	F	A		+							•	
21	Н 788	F	A								+		
22	H 791/A	F	A				+				· ·		

	S	upraorbit	al ridge	es	For	rehead s	slope	1	Nasal	profile	4	Shape	of nasa	l bones
*			·	ent		Slightly receding	80	,	a a		Concavo-convex	Narrow constricted	Broad constricted	shaped
No.	Absent	Slight	Medium	Prominent	Vertical	Slightl	Receding	Straight	Concave	Convex	Concav	Narrow	Broad	Wing shaped
	ETERY	***************************************						L						
1				+			+		+					+
2				+			+							
3			+			+			+					+
4	+				+				+				+	
5				+	*,		+	+						+
6				+	-		+							+
7				+			+		+					+
8				+			+							
9				+	*/*	+								
10 11		-T.		+			+		+			+		
12		+				+								
13		·		+			+		+			+		+
14				+			+		+			т		+
15				, -+-			+		+					+
16					ķ:				+			+		
17		+				+								
18	+				+									
19	+				+				+			+		
20			+			+			+					+
21	+													

-			Nasal ro	ot depres	sion	1	Margo piri	formis int	erior		Subnasa	prognath	nism
	No.	Absent	Shallow	Medium	Depressed	Amblykraspedotic	Bothrokraspedotic	Oxykraspedotic	Orygmokraspedotic	Absent	Slight	Medium	Marked
_	CEMET	ERY	R 37										
	1			+		+						+	
	2				+	ı							+
	3			+				+				+	
	4	+				+					+		
	5				+			+					+
	6			+				+					+
	7				+	+							+
	8				+								
	9												
	10			+				+					+
	11												
	12	+						+					+
	13				+			+				+	
	14				+	+							+
	15				+			+					
	16							+		+			+
	17		+					+		_			·
	18		+					+		+			
	19	+										+	
	20			+				+				+	
	21		+					+				F	
	22	+											
	Н												

21

22

		Occipital p	rotuberance		Occipital	contour		Dental a	arch		Shape o	f foramer	magnui	m
No.	Round	Moderately protruding	Markedly protruding	Flat	Wedge shaped	House shaped	Upsiloid	Ellipsoid	Paraboloid	Elliptic	Oval (base posterior)	Oval (base anterior)	Round	Rhomboid
CEME?	TERY	R 37												
1			+		+		+							
2		+				+			+		+			
3	+					+	+			+				
4	+					+			+		+			
5	+					+			+			+		
6		+				+			+		+			
7		÷				+	+			+				
8														
9		+				+								
10			+			+			+					+
11			+			+			+				•	
12														
13			+			+			+			+		
14 15		+				+	+						+	
16		+				+	+				+			
17						r			+		+			
18		+				+			+	+				
19		+				+			+				+	
20		2 8												
~ 0		+			+		+				+			

+

					_	Cr	ranial con	tour in no	rma vert	icalis			rtical aspe rcus zygor	
No.	Skl No.	Se	X	Age	Ellipsoides	Pentagonoides	Rhomboides	Ovoides	Sphenoides	Sphaeroides	Byrsoides	Phaenozygy	Orthozygy	Cryptozygy
CEM	ETERY	R 37	,											
2 3	Н 795/А		F	A				+				+		
24	Н 798/А	I	F	A							+	+		
25	Н 798 (a)	F	A				+					+	
26	H 801/A		F	A	+									+
27	H 804		F	A										
28	H 805		F	A	+								+	
29	H 806/A		F	A		+								+
30	H 812		F	A	+									+
31	Н 816		F	A	+							+		
32	H 817		F	A					+				+	
33	Н 820 ([)	F	A				+						+
34	Н 820 (F	A					+					1
				A					+					+
36	H 798/A	. 2	F	Juv.				+						+

ARE	A G							
37	I S 11	M	A		+		+	
38	II S 18	M	A	+				+
3 9	II S 42	M	A			+		+
40	III S 2	M	Α	+			+	

Slight Nedium Prominent Prominent Straight Concave Concave Concave Marrow constricted Broad constricted Wing shaped		
nn al ly receding ht ht constricted constricted shaped		
nent nent ng ve-convex vo-convex constricted shaped		
al ng hat hat hat hat hat hat hat hat hat hat		
ng nt vo-convex vo-constricted constricted shaped		1
ve vo-convex vo-convex constricted shaped	receding	
vo-convex vo-convex constricted shaped	aight	<u> </u>
vo-convex v constricted constricted shaped		
vo-convex v constricted constricted shaped		
v constricted constricted shaped		
constricted	constricted	
shaped	constricted	
	shaped	

CEMETERY R 37

23		+				+						
24		+		+				+				+
25			+		+			+			+	
26	+			+				+		+		
27	+			+								
28	+			+				+				+
29			+		+		+					+
30			+		+			+				
31			+		+				+			+
32	+				+		+			+		
33		+		+		*						
34		+			+			+			+	
35		+			+			+				+
36		.+			+		+				+	

AREA G



	1	Nasal roo	t depressi	on	I.	Margo pirif	ormis infe	rior		Subnasal	prognathis	m
No.	Absent	Shallow	Medium	Depressed	Amblykraspedotic	Bothrokraspedotic	Oxykraspedotic	Orygmokraspedotic	Absent	Slight	Medium	Marked
	rery i				<u> </u>				1 ~			
23		+					+					+
24		+					+		+			
25			+				+				+	
26	+						+		+			
27	+						+		+			
28	+						+		+			
29			+					+				+
30			+									
31			+				+				+	
32	+						+			+		
33	+						+		+			
34			+		+							+
35			+				+					+
36	+						+		+			

AREA G

37		+	+		
38		+		+	+
39		+		+	
40	_L_			+	

tely protruding ly protruding shaped d d	Occipital protuberance
ly protruding shaped shaped d	protuberance
shaped shaped d	ance
shaped	e
	Occipita
a bid	al contour
Įq	Den
	tal arch
	1
Elliptic	5
Oval (base posterior)	Shape of
Oval (base anterior)	foramen
Round	magnum
Rhomboid	1

CEMETERY R 37

23	+				+		+				
24	+				+	+					
25	+				+	+					+
26		+		+					+		
27											
28		+			+			+			
29		+		+			+				
30		+			+	+			+		
31	+				+ "	+					
32					+ -		+				
33		+			+	+				, + <u>;</u>	
34	+						+				
35	+				+		+				
36		+			+		+	+			

AREA G

37		+	+	+	+
38	+		- - 4	+	+
39		+	+	+	+
:40	+		Д.	+	

)					Cr	anial conto	ur in nor	ma vertica	lis			al aspect is zygoma	
No.	Skl No.	Sex	Age	Ellipsoides	Pentagonoides	Rhomboides	Ovoides	Sphenoides	Sphaeroides	Byrsoides	Phaenozygy	Orthozygy	Cryptozygy
ARE	A G												
41	III S 22	M	A							+	+		
42	III S 23	M	A		+							+	
43	III S 47	M	A					+			+		
44	I S 13	F	A	+									
45	II S 5	F	A					+			+		
46	III S 21	F	A	+							+		
MOU	ND AB												
52	Н 11635	M	A					+			+		
54	Mound AB (Rec.)	F	A	+									

56	H 184 (K)	M	A	
57	H 307 (a)	M	A	+
5 8	H 484 (a)	M	A	
5 9	H 487 (a)	M	Α	
60	H 502 (G)	M	A	

	Supraork	oital ridges		Fo	rehead slope		Nasa	l profile		Shape	of nasa	bones
No.	Absent Slight	Medium	Prominent	Vertical	Slightly receding Receding	Straight	Concave	Convex	Concavo-convex	Narrow constricted	Broad constricted	Wing shaped
AREA	A G											
41			+		+		+			+		
42		+			+				+			+
43	+			+			+			+		
44	+			+								
45	+			+ .		+	,			+		
46	+				+		+			+		

MOUND AB

CEMETERY H (OPEN BURIAL)

		Nasal ro	ot depressi	ion	Ma	argo pirii	ormis info	erior		Subnasal	prognath	ism
No.	Absent	Shallow	Medium	Depressed	Amblykraspedotic	Bothrokraspedotic	Oxykraspedotic	Orygmokraspedotic	Absent	Slight	Medium	Marked
AREA	G				-	×						
41		+					+			+		
42		+			+					+		
43	+						+				+	
44							+					
	+						+			+		
45												

MOUND AB

CEMETERY H (OPEN BURIAL)

 56
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +
 +

I

	O cci	pital pr	otubera	nce	Occipital	contour	D	ental a	arch	Access on the contract of the	Shape o	f forame	en magn	um
No.	Round	Moderately protruding	Markedly protruding	Flat	Wedge shaped	House shaped	Upsiloid	Ellipsoid	Paraboloid	Elliptic	Oval (base posterior)	Oval (base anterior)	Round	Rhomboid
REA	A G													
41		+				+			+					
42	+				+				+		+			
43		+				+			+		+			
44	+									+				
45	+				+				+				+	
46	+					+			+		+			
MOU	IND AE	3												
52				+		+	+				+			
54		+				+			+	+				

					Cran	ial contou	ur in nor	ma vertic	alis			cal aspect us zygoma	
No.	Skl No.	Sex	Age	Ellipsoides	Pentagonoides	Rhomboides	Ovoides	Sphenoides	Sphaeroides	Byrsoides	Phaenozygy	Orthozygy	Cryptozygy
CEMI	ETERY H	(OPE	N BU	RIAL)									
61	H 69 5	M	A		+						+		
62	Н 698	M	A				+				+		
63	H 88	F	A	+									
64	H 306 (a)	F	A										
65	H 488	F	A					+			+		
66	H 501 (a)	F	A										
67	Н 699	F	A				+						
68	H 710	F	A	+									
				_									
CEM	ETERY H	(JA	R BUF	RIAL)									
69	H 206 (B)	M	A								+		
70	H 255 (a)	М	A	+							+		
71	Н 344	Μ	A				+				+		
· 72	H 61	F	A										
73	H 153 (a)	F	A					+					
74	Н 154 (а)	F	A	+									
75	H 206 (d)	F	A				+					+	
76	H 245 (e)	F	A				+						

77 H 246 (c) F A 78 H 247 (a) F A


Si	upraorbital ri	dges	Fore	nead slope	Nasa	l profile	Shape o	f nasal	bones
No. No.	Slight Medium	Prominent	Vertical	Slightly receding	Straight	Convex Concavo-convex	Narrow constricted	Broad constricted	Wing shaped

61			+			+					
62				+			+				
63	+				+			+			+
64		+			+				+		+
65		·¥j	+			+		+			+
66	+				+						
67								+			+
68	+				+						

69					•	
70		+	+	+	+	
71		+	+	+	+	
72						
73	+	+		+		
74	+	, +				
75	+		+	+		+
76						
777	+	*		+		+
78			+			

	Nasal root depression					М	ar go p irif	ormis infe	rior		Subnasa	prognath	ism
													ř
No.	Absent	Shallow		Medium	Depressed	Amblykraspedotic	Bothrokraspedotic	Oxykraspedotic	Orygmokraspedotic	Absent	Slight	Medium	Marked

61							+
62			+		+		+
63	+				+	+	
64		+					
65			+				+
66	+					+	
67	+				+		+
68		+					



	Oce	cipital pr	otuberance	e	Occipital	contour	De	ental arc	h	S	hape of	foramen	magnun	1
No.	Round	Moderately protruding	Markedly protruding	Flat	Wedge shaped	House shaped	Upsiloid	Ellipsoid	Paraboloid	Elliptic	Oval (base posterior)	Oval (base anterior)	Round	7 m o 4 D

61	+		+		+					
62	+			+			+		+	
63		+		+			+			
64					+					
65		+		+						
66						+				
67	+			. +			+			+
68		+						+		

69	+				+	+		
70		+			+	+		+
71	+				+		+	
72	+				+			
73		+	4					+
74	•	+			+			
75	+	. ' .		+	,	4,		+
76	+,				+			
77		+			+			
18								

			1			Cranial	contour i	n norma	verticalis	* 5 (*** 5) ***		rtical asp rcus zygo	ect of the omaticus
No.	Skl No.	Sex	Age	Ellipsoides	Pentagonoides	Rhomboides	Ovoides	Sphenoides	Sphaeroides	Byrsoides	Phaenozygy	Orthozygy	Cryptozygy
CEM	ETERY H	[(JA]		<u> </u>									
79	H 247 (b)	F	A	+							+		
80	Н 7435 (b) F	A	+									
81	Pot no. 1	1 F	A					+			+		
82	Pot no. 12	(a) F	A					+				+	
83	Pot no. 12	(b) F	A	+									
85	H 245 (a)	F	Juv.				+					+	

-				_										
	Sup	raorbit	al rid	ges	, F	orehead	slope	1	Nasa	l profile		Shape	e of nas	al bones
No.	Absent	Slight	Medium	Prominent	Vertical	Slightly receding	Receding	Straight		Сопуех	Concavo-convex	Narrow constricted	Broad constricted	Wing shaped
CEME	ETERY I	H (JA	AR BU	JRIAL)										
79		+			+			+						+
80		+				+								
81		+			+				+					+
82	+				+				+					
83														
85	+						+							

		Nacal roo	ot depress	ion	M	argo pirif	ormis infe	erior		Subnasal	prognathis	im
No.	Absent	Shallow	Medium	Depressed	Amblykraspedotic	Bothrokraspedotic	Oxykraspedotic	Orygmokraspedotic	Absent	Slight	Medium	Marked
	·	H (JAR	BURIAL)		,							
CEMET	LEKY 1								+			
79		+										
80											+	
81	+						+		+			
82		+			+				+			
83												
85		+					+		+			

Occipital	protuberance	Occipital contour	Dental arch	Shape of foramen magnum
Round Moderately protruding	Markedly protruding Flat	Wedge shaped House shaped	Upsiloid Ellipsoid Paraboloid	Elliptic Oval (base posterior) Oval (base anterior) Round Rhomboid

CEMETERY H (JAR BURIAL)

				+	+		79
	+			+	+		80
+		+		+	+		81
	+		+	+		+	. 82
+				+	+		83
+		+		+		+	85
		+					00

In 'eranial contour in norma verticalis' (Collective Tables G to ${\rm K})$ words byrsoides and sphenoides should be transposed.

-				1	Н	u m	e r u	5	1	R	a d	i u	s
				M 1	М 3	M 4	M 7a	М 7	M 7/M 1	M 1	M 2	M 3	M 3/M 2
N	o. Skl No.		Age	Maximum length	Breadth of proximal epiphysis	Breadth of distal epiphysis	Circumference of the shaft at the middle	Minimum circumference of diaphysis	Robusticity index	Maximum length	Physiological length	Minimum circumference of diaphysis	Robusticity index
CE	METERY R 37												
1	Skl 1 [Rt.]	M	A	361.0	56 .5	72.0	77.0	65.0	18.01	_	_	_	_
2	Skl 10 [Rt.]	M	Α	389.0	55.5	67.0	6 7. 0	55.0	14.14	_	_	_	_
3		M	Α	_	_	****		_		_		37.0	
4		M	A		_	_		. —	_	_	_		_
5	H 793 [Lt.]	M	Α	_				_			_	_	
6	H 794 [Lt.]	M	Α	_	_	_	_	_			_	_	
7	H 796 (B) [Rt.]	M	Α	344.0?	51.0?	66.0?	73.0	6 3. 0	18.31?		_		
8	H 798 (ii) [Rt.]	M	Α	321.0	_		60.0	56.0	17.45		_	_	
9	H 798 (ii) [Lt.]	M	Α	_	_	-				_	_	_	_
10	H 811 [Rt.]	M	Α	_							_	_	summittee.
11	H 815 [Lt.]	M	A										
12	H 818 [Lt.]	M	Α	301.0	45.0?	57. 0	57.0	55. 0	18.27				_
13	Н 821 [Lt.]	M	Α			_					_		
14	Skl 2[Lt.]	F	Α		- Allerton			_		_			and territors
15	H 780 [Rt.]	F	Α.			_	-				_		
16	H 792 [Rt.]	F	A							_		_	
17	H 804 [Rt.]	F	A					-					e-Printerio
18	H 805 (a) [Lt.]	F	Α	314.0	43.0		55.0	50.0	15.92		www.		
19	H 806/A [Rt.]	F	A	*******		52.5	-	52.5				accepta	MATERIAL STATES
20	H 806/A [Lt.]	F	A	293.0	42.0	53. 0	52.0	50.0	17.06		. SOLUTION		
21	H 820 [Rt.]	F	A	**********	decision).		-	program.	-				Titlering
22	H 820 (c) [Lt.]	F	A			and the same of th	- cyclastic		NAME OF STREET			******	-

	ι	JIn	a				F	e m	u	r		
	М 1	M 2	5/13 M	3/M 2	M 1	M 2	M 10	M 9	М 6	M 7	M 8	M 18
No.	Maximum length	Physiological length	Minimum circumference of diaphysis	Robusticity index	Maximum length	Length in natural position	Sagittal diameter at upper flattening	Transversal diameter at upper flattening	Sagittal diameter at middle of diaphysis	Transversal diameter at middle of diaphysis	Circumference of the shaft at the middle	Vertical diameter of caput
CEMET	ERY R	37										
1		_		_	518.0	514.0	30.0	33.0	34.5	32.5	100.0	54.0
2			_		544.0	542.0	29.0	35. 0	29.0	31.0	92.0	53.0
3	-	277.5	40.0	14.41	545.0	543.0	29.5	36.0	27.0	31.0	93.0	53.5
4	_	-	_		467.0	462.0	23.5	29.0	22.0	28.0	79.0	44.4
5	-		-		_		_	_	_		_	_
6	-	-	_		479.0	476.0	29.0	34.5	33.5	28.5	95.0	49.0
7	_	-	× .	-	_	_		_	_	_	_	_
8	_		_		_	-	_	_	_			
9	_				_		_		_		_	-
10		-			475.0?		27.5	35.5	29.0	27.5	93.0	
11	_			-		_	24.0	31.0	26.0	27.0	83.0	_
12		_			_			_	_	_	_	_
13	_	_	_			_	-	_		_	_	
14		**************************************					_					-
15		-		_	437.0	431.0	25.0	27.5	26.0	25.0	80.0	40.0?
16		253.0	29.0	11.46			Politica					_
17	,					,	Notherina		equip. (III)	Francis		_
18	3	******	-					-		-		
18			-	200	· <u>~</u>			*****	*****	, · -		Phone:
20				access.					-	******		enement .
2		~	*******	and the same of th			on and a					
2	2	Arrane	26.0	desired	_			-				-

					F e	m	u ı	r		-	
	M 19	M 20	M 11	M 12	M 21	M 8/M 2	M6+M7 M2	M 6/M 7	M 10/M 9	M 11/M 12	M18+M19 M 2
No.	Transversal diameter of caput	Circumference of caput	Distal sagittal diameter	Distal transversal diameter	Epicondylar breadth	Length thickness index	Robusticity index	Pilastric index	Platymeric index	Popliteal index	Robusticity index of caput
EMETI	ERY R	37									
1	53.5	172.0	32.0	42.5	92.0	19.46	13.04	106.15	90.91	75.29	20.91
2	52.0	169.0	34.0	44.0	90.0	16.97	11.07	9 3.55	82.86	77.27	19.37
3	52.5	168.0	33.5	42.0	88.0	17.13	10.68	87.10	81.94	79.76	19.52
-1	45.0	140.0	29.0	37. 0	76.0	17.10	10.82	78.57	81.03	78.38	19.26
5	_	_		_		_	_	_		_	
6	49.5	156.0	31.0	42.5	87.0	19.96	13.03	117.54	84.06	72.94	20.69
7	-		-		_		_	_	_	-	
8		_		_	-		one con				**************************************
9							-	-	_	-	_
10	-		30.5	45.0	_	- Articles		105.45	77.46	67.78	
11	_		27.0	42.5		-		96.30	77.42	6 3.53	and the same of th
12		-	-	-		-		-		(percent and	_
13	_				_						
1-4	_			_			Persona	_			
15	41.0	130.0	26.0?	26.5	73. 0	18.56	11.83	104.00	90.91	98.11?	18.79?
6								Property.		waterspan	
.7	pulse and the second		The state of the s		-	No.	_			property (Ministration.
8				_		eneman.		*********			
9			*****					_		Named	Principles .
0	_				-		-	Named			Name of the last o
1				**Propos	*******	-		-		-	-
2		-		******	-			- Names		-	

		•	T i	b i	a	*		Fi	b u	ı I a
	M 1a	M 2	M 8	M 9	M 10b	M 9/M 8	M 10b/M 1a	M 1	NJ 4a	M 4a/M 1
No.	Maximum length (Spino-malleolar)	Physiological length	Sagittal diameter at the middle	Transversal diameter at the middle	Minimum circumference of diaphysis	Index of cross-section in the middle	Robusticity index	Maximum length	Minimum circumference	Robusticity index
EMET	TERY R 3	37								
1	_			_	-		_	_	_	~
2	457.0?	433.0	36.0	24.0	83.0	66.67	18.32?	_	_	<u> </u>
3	460.0	433.0	36.0	21.5	83.5	58.90	18.35	435.0	39.0	8.97
4	_	348.0	28.0	20.0	71.0	71.43		-		_
5	374. 0	348.0?	31.5	19 .5	68.0	61.90	18.50		_	
6		370.0	35. 0	23.0	88.0	65.71			_	_
7		_	_	_		_	_	-	_	
8						_			_	_
9	_		33.0	20.0	72.0	60.61			_	
10	_							_	_	_
11	_					_	_	-	_	_
12	378.0	35 7.0	27.0	20.5	64.0	75. 9 3	17.16	368.0	34.0	9.24
13	379.0	352.0	34.0	19.5	76.0	57.35	20.21	-		-
14	372.0	352.5	30.0	19.5	71.0	65.00	-		_	
15				-	emonatus.			-	_	
16				_	and the second	_		~		_
17		-	24.0	19.0	6 2. 0	79.17	, and the same of	_		
18	Street	, , , , , , , , , , , , , , , , , , ,	-		******					
19	pupositi	SP-COMP	• •		,	_	-	-	appendix.	*******
20			aurtuppe		orannos	Maria		_		
21	386.0	384.0	23.5	20.0	6 3.5	8 5.11	16.71			_
22	No.	- Augus	Aminosis	(mineral)			_	reases		

					н	u m	e r u	ı s		F	a d	i u	S
				M 1	M 3	M 4	M 7a	M 7	M 7/M 1	M 1	M 2	M 3	M 3/M 2
No.	Skl No.	Sex	Age	Maximum length	Breadth of proximal epiphysis	Breadth of distal epiphysis	Circumference of the shaft at the middle	Minimum circumference of diaphysis	Robusticity index	Maximum length	Physiological length	Minimum circumference of diaphysis	Robusticity index
CEN	METERY H (OF	EN	BURIA	L)									
23	H 184 (K) [Lt.]		A	321.0	5 0 .5	62.0	67.0	61.0	19.00	256.0	240.5	42.0	17.46
24	H 484 (a) [Rt.]		A	_				_	_	_			
25	H 484 (a) [Lt.]	М	A	_	_		_		_	_	_		_
26	H 502 (G) [Rt.]	M	A	324.0	50.0	64.5	6 5.5	58.0	17.90	251.0	232.0	41.0	17.67
27	H 502 (G) [Lt.]	M	A	327.0	52. 0	64.0?	62.0	55.0	16.82	_		_	_
28	H 636 [Rt.]	M	A	_		_				_	_		
29	H 694 [Lt.]	Μ	A	326.0	51.5	64.0	6 5.5	60 .5	18.56	-		_	_
30	H 695 [Rt.]	M	A	329.0	52. 0	63.0	69.0	61.0	18.54		-		
31	H 696 [Rt.]	M	A	320.0	research	5 9 .5	66.0	58.0	18.12				- Constitute
32	H 696 [Lt.]	M	A				_		-	_			
33	H 697 [Rt.]	M	A				_	65.5	_			_	
34	H 697 [Lt.]	Μ	Α							_	_		-
35	H 698 [Rt.]	M	A				69.0	58.0		_	257.0	41.0	15. 9 5
36	H 698 [Lt.]	M	A	3 41.0	51.0	_	6 3. 0	57. 0	16.72			_	TO COMPANY.
37	H 88 [Rt.]	F	A	Palaborin				_			Approximate to the second	Amagana	mendena
38	H 88 [Lt.]	F	A			Name of the last o			Antonores	-	Phaetin		Name of the last o
39	H 306 (a) [Rt.]	F	Α	298.0	44.0	57.5	62 .5	55. 0	18.46		-	-	
40	H 306 (a) [Lt.]	F	A	292.0?	43.0	No.	62.0	55.0	18.84?	2 33. 0	218.0	35.0	16.06
41	H 485 [Lt.]	F	A			or a constant	55. 0	49.0	tiouri militari.	-	Brookele	_	- Thomas
42	H 488 [Rt.]	F	A		er-schools	55. 0		55.0		221.0	208.0	38.0	18.27
43	H 488 [Lt.]	F	A	267.0	40.0	54.5	63.0	5 4.0	20.22	221.0	206 .5	37.0	17.92

		U I	n a	×			F e	m	u	r ,		
	M 1	M 2	M 3	M 3/M 2	M 1	M 2	M 10	М 9	M 6	M 7	М 8	М 18
No.	. Maximum length	Physiological length	Minimum circumference of diaphysis	Robusticity index	Maximum length	Length in natural position	Sagittal diameter at upper flattening	Sagittal diameter at upper flattening	Sagittal diameter at middle of diaphysis	Transversal diameter at middle of diaphysis	Circumference of the shaft at the middle	Vertical diameter of caput
CEMETI	ERY H (OPEN B	BURIAL)			,						
23	_	_	_	_	_	_	_	-	_	_	<u> </u>	_
24	_	_		_	_	_	_	_		_	_	_
25	-		_	_	_			_		_	_	_
26	271.0	244.0	32.0	13.11	_		22.5	31.5	25.0	26.0	85.0	44.5
27	268.0	241.0	31.0	12.86		_	22.0	30.5	23.0	27.5	83.0	45.0
28					_	_	-	_		_	_	_
29	en en en		-		_		_	_		_	_	-
30			_				-	_		_	-	
31					_		_	_		~		—
32		witness				_	-	_	_	_	-	
33	270.0?	249.0			and the same of th					-	_	
34	Philippine.	248.0?	37. 0	14.92?		Town	******	terripa.		_	n-mannage .	
35			manufaction of the second				25.0	33.5	29.5	27.0	88.0	_
36	289.0	262.0	34.0	12.98			27.0	30.0	31.0	28.5	95.0	48.5
37		238.0	26.0	10.92				_	_		_	_
38	2 5 3. 0	232.0	, 27.0	11.64		Anaman		***************************************	. —		_	-
3 9	~~							-				_
40		228.0	32,0	14.04	garagement			-	_		_	
41	et derin.			- N	-		-	***********	Promotes .	, —	-	
42	243.0	218.0	32.0	14.68	379.0	37 8 .5	21.0	27.0	26.0	25.0	80.0	38.0
43	242.0	217.0	31.0	14.29	378.0	378.0	20.0	28.0	26.0	26.0	80.0	38.0

	1				F	e m	u	r			
	M 19	M 20	M 11	M 12	M 2	1 M 8/M	12 M6	-M7 M 6/M	7 M 10/M 9	M 11/M 1	M18+M19 M 2
No.	Transversal diameter of caput	Circumference of caput	Distal sagittal diameter	Distal transversal diameter	Epicondylar breadth	Length thickness index	Robusticity index	Pilastric index	Platymeric index	Popliteal index	Robusticity index of caput
CEMET	ERY H	(OPEN I	BURIAL)								
23			_	_		-		_	_		-
24	_	_	_	_		_	-	-	_	_	_
25	_	_	_	_	_	_	_			_	_
26	45.0	147.0	27.0	38.0		_	_	96.15	71.43	71.05	
27	44.0	141.0	_	_	_	_	-	83.64	72.13	_	_
28	_	_		-	_	` —	-		_	-	_
29	_	_		_	-		_	_	_		
30	_		_		_	_		_	_		
31	-	_				_		_	_		_
32	_	_	_		_		_		_	_	
33	_	-				_	_		_	_	_
34	_	_				-	_		_	_	_
35		_	29.0	42.0	8 3.5		_	109.26	74.63	69.05	
36	47.5	152.0	29.0	44.0	82.0	_		108.77	90.00	65.91	
37					_	-	-	-		-	-
38	_	—			_	-			_		_
3 9	_			- Compa						******	
40	-				_		Prilitaria				Protecting.
41		101.0	25.5						******	district	
42	37.5	121.0	25.5	31.0	66.0	21.14	13.47	104.00	77.78	32.26 1	9.95
43 K	3 6 .5	117.0	26.0	36.0	67.0	21.16	13.76	100.00	71.43	72.22 1	9.7L

		т	ı	b i	a		1	FI	b u	1 a
	M 1a	M 2	M 8	M 9	M 10b	M 9/M 8	M 10b/M 1a	M 1	NI 4a	M 4a/M 1
No.	Maximum length (Spino-malleolar)	Physiological length	Sagittal diameter at the middle	Transversal diameter at the middle	Minimum circumference of diaphysis	Index of cross-section in the middle	Robusticity index	Maximum length	Minimum circumference	Robusticity index
CEMET	ERY H (OPEN BUR	IAL)							
23	_	344.0	37.0	22.0	80.0	59.46		_	_	_
24	410.0	381.5	3 9 .5	23.0	85.0	58.23	21.12	_	_	_
25	_		_	_	_	_	_	_	40.0	_
26		_			_	_	_	_	_	_
27		372.0	33.0	23.0	80.0	69.70		_	_ ,	_
28			31.0	20.0	70.0?	64.52	-	_	_	_
29					-		-	_		-
30			Account	-		_	_	_	_	_
31			-	_	_	-	_	_		_
32		_			-	-	_	369.0	40.0	10.84
33					-	100,000	_	_	_	_
34			and the same of th	_	-		_	-	_	
35	_		34.0	24.0		70.59	-		_	
36			35.0	25.0	85.0	71.43	_		_	
37		-				, -				
38	****	_		_	-		_		_	
89	352,0?	832.0	80.9	20.5	68.0	68.33	19.54	_		
40	_	mage		enth.				*****		annum .
41			-		-	·, —		- 1		
42	328.0	308.0	26.5	17.0	67.0	64.15	21.07	- Annual	_	
48	3		and the same of th			-	_		recent.	-

			Н	u m	e r u	s		1	R a d	i u	s
		M 1	М 3	M 4	M 7a	M 7	M 7/M 1	M 1	M 2	M 3	M 3/M 2
o. Skl No. Se	x Age	Maximum length	Breadth of proximal epiphysis	Breadth of distal epiphysis	Circumference of the shaft at the middle	Minimum circumference of diaphysis	Robusticity index	Maximum length	Physiological length	Minimum circumference of diaphysis	Robusticity index
EMETERY H (OPE	N BURI	AL)								*	
4 H 501 (a) [Rt.]	F A	298.0	46.5	_	6 5. 0	56.0	18.79	_		_	
5 H 5 0 1 (a) [Lt.]	F A	289.0	_	_	64.0	55. 0	19.0 3				
6 H 699 [Lt.]	F A	_	_	54.5	58.0	53. 0	_	_	_		
	F A	277.0	44.5	5 8.0	60.0	5 4.0	19.49				
8 H 710 [Lt.]	F A	279.0	44.5	58.0	60.0	53.5	19.18			-	
EMETERY H (^{JAR}	BURIA	L)								•	
	M A	-		_		_	_				
	F A				_						-
51 H 154 (a) [Lt.]	F A					_					
52 H 206 (c) [Rt.]		319.0	4 5. 0	54.0	5 2.0	51. 0	15. 99		_	-	
53 H 206 (c) [Lt.]		318.0	43.5	55.5		55. 0	17.30	_			
54 H 206 (d) [Rt.]		279.0	4 3. 0	53.5	5 9.0	51. 0	18.28	-	_	-	_
11 200 (0) 1									242.0	37.0	15.29

	•	J I	n a				F	e n		r		
-	M 1	M 2	М 3	M 3/M 2	M 1	M 2	M 10	М 9	M 6	M 7	M 8	M 1
No.	Maximum length	Physiological length	Minimum circumference of diaphysis	Robusticity index	Maximum length	Length in natural position	Sagittal diameter at upper flattening	Transversal diameter at upper flattening	Sagittal diameter at middle of diaphysis	Transversal diameter at middle of diaphysis	Circumference of the shaft at the middle	Vertical diameter of caput
EMET	ERY H	OPEN B	URIAL)									
44	_	_	_	_	_	_		_	_	_	_	_
45			_	_	-	_	24.0	28.0	27.0	23.0	80.0	_
46	_		_	_		_	-	_	_	_	_	_
47	245.0?	227.0	32.5	14.32	_	_	22.5	28.0	26.0	26.0	81.0	_
48		224.0	32.0	14.29	410.0	408.5	23.0	30.0	24.0	27.5	81.0	41.0
EME	TERY H	(JAR B	- URIAL)		410.0	408.5	23.0	30.0	24.0	27.5	81.0	41.0
EME:	 ГЕRY Н 240.0?		-	14.29	410.0	408.5	23.0	30.0	24.0	27.5	81.0	41.0
CEME 2 49 50		(JAR B	URIAL) 31.0 —		410.0	408.5 	_ _ _			_	_	41.0
CEME 3 49 50 51		(JAR B	- URIAL)		410.0	408.5 — —	_ _ _ 24.0		_ _ _ 26.0	 25.0	— — 80.0	. <u>-</u>
EEME 3 49 50 51 52		(JAR B	URIAL) 31.0 —		410.0 — — —	408.5 — — —	_ _ _			_	_	41.0
CEME 3 49 50 51		(JAR B	URIAL) 31.0 —		410.0 ———————————————————————————————————	408.5 	_ _ _ 24.0		_ _ _ 26.0	 25.0	— — 80.0	. <u>-</u>

					F e	m	u	r		-	
	M 19	M 20	M 11	M 12	M 21	M 8/M 2	M6+M7	M 6/M 7	M 10/M 9	M 11/M 12	M 18+M 19
No.	Transversal diameter of caput	Circumference of caput	Distal sagittal didameter	Distal transversal diameter	Epicondylar breadth	Length thickness index	Robusticity index	Pilastric index	Platymeric index	Popliteal index	Robusticity index of caput
CEMET	ERY H	(OPEN B	URIAL)								
44	_	_	_	_	_	-		-	-		_
45	_	_	29.0	42.5	71.5?	_	_	117.39	8 5.71	68.24	_
46	_	_			_	-	-	_		-	_
47		_	27.0	39.0	79.0?	_		100.00	80 .36	69 .23	_
48	_	_	25.0	43. 0	73. 0 ?	19.83	12.61	87.27	76.67	58.14	_
CEMETI	ERY H (JAR BUF	RIAL)								
49	_			_	-		_				Territoria
5 0	_	_	_			-	-		_	-	
51	_		27.0	35. 0		-	_	104.00	85.71	77.14	_
52	41.0	127.0	.		-	-		91.07	73.33		-
53	-				-		_		-	- '	-
54	-	_	-		-	-			-		-
55		_		Maria	_	-	_		-	-	_

	1		T i	b	i a			FI	b u	l a
	M 1a	M 2	M 8	M 9	M 10b	M 9/M 8	M 10b/M 1a	M 1	M 4a	M 4a/M 1
No.	Maximum length (Spino-malleolar)	Physiological length	Sagittal diameter at the middle	Transversal diameter at the middle	Minimum circumference of diaphysis	Index of cross-section in the middle	Robusticity index	Maximum length	Minimum circumference	Robusticity index
CEME?	TERY H (OPEN BUR	IAL)							
44	_	325.0	29.5	20.0	70.0	67.80	_	_	_	_
45	_		_		_			_		_
46		_			_		_	_	_	_
	0000	3 08.0	28.0	18.0	66.0	64.29	20.37	323. 0	35. 0	10.84
47	332. 0	300.0	20.0	10.0	00.0					

49							_	-		
5 0	3 6 5. 0	345.0	31.5	22. 0	74.0	69.84	20.61	_		
51	_									_
52					and the	_	_	_		_
53							-			
54	energia.	name and a second							-	
55	energia.						***************************************			********

				,	ACCORDING	TO MANOUVE	RIER'S TABL	ES	
No.	Skl. No.	Sex	Humerus (a)	Radius (b)	Ulna (c)	Femur (d)	Tibia (e)	Fibula (f)	Average
CEN	IETERY R 3	7							
1	H 7 9 3	M		_	_	1699.38	1682.40	_	1690.89
2	Н 794	M	_	_		1728.00	_		1728.00
3	Н 796 (В)	M	1723.00		_	_			1723.00
4	H 798 (ii)	M	1651.50		-	_			1 6 51.5 0
5	Н 811	M				1720.00	_		1720.00
6	Н 818	M	1575.75	_		_	1690.40	1680.60	1648.92
7	Н 821	M			_	-	1692.60	_	1692.60
8	Skl. 1	\mathbf{M}	1805.25			18 35. 60			1820.42
9	Skl. 10	M	192 7. 6 3	_	_	1929.14	1 989 .36	1909.69	1938.96
10	н 780	F	_			15 9 7.43	_		1597.43
11	H 805 (a)	F	1605.20				-	_	1605.20
12	H 806/A	F	1550.80	_	_	_	_	_	1550.80
13	н 820	F		_		_	1715.00	-	1715.00

CEMETERY	H	(OPEN	BURIAL)
----------	---	-------	---------

14	H 184 (K)	M	1 6 51.5 0	-				_	1651.50
15	H 484 (a)	M	-	***************************************		_	1795.80		1795.80
16	H 502 (G)	M	1664.50	1703.33	1706.50	-	_	-	1691.44
17	Н 694	M	1666.00				-	_	1666.00
18	H 695	M	1674.25						1674.25
	н 696	M	1649.00	-			*****	1682,40	1665.70
	Н 697	M	Marine A.		1709.66	wysiania		NO.00.	1709.66
21	Н 698	M	1711.25	-	1818.00	ententa .	-	-	176.1.62
22	H 88	F	_		165 5. 00		-	enema.	1655.00

			PEARSON'S SION FORMU	LAE'	DUPERTUIS & HADDEN'S 'GENERAL FORMULAE'						
No.	Femur (a)	Humerus (b)	Tibia (c)	Radius (d)	Average	Femur (a)	Tibia (b)	Humerus (c)	Radius (d)	Average	
CEMETE	ERY R 37	•									
1	1691.0	-	_	_	1691.0	1736.0	_		-	1736.0	
2	1713.6	_	_	_	1713.6	1762.9	_	-		1762.9	
3	_	1701.9?	_	_	1701.9	-	_	1757.4?		1757.4?	
4	_	1635.4		_	1635.4	_		1689.1		1689.1	
5	1706.1?	_	_		1706.1?	1753. 9		_	_	1753.9	
6	_	1577.5	1672.9	_	1625.2		1709.1	1629.7	_	1669.4	
7	_		1680.0		1680.0	_	1716.3		-	1716.3	
8	1786.9	1751.1	_	_	1769.0	1850.2	—	1807.9		1829.0	
9	1835.8	1832.2	186 3. 0 ?		184 3.7?	1908.4	1900.5?	1891.0	_	1900.0	
10	1578.4	_	_		1578.4	1626.6	_	_	_	1626.6	
11		1579.5	_	_	1579.5	_	_	1637.0		1637.0	
12		_		Palace	_		-	_	_	-	
13		_	164 1.5		1641.5		1688.3	_		1688.3	

CEMETE	RY H	OPEN BURI	(AL)							
14		16 35.4		1696.6	1666.0		_	1689.1	1738.4	1713.8
15			1743.0		1743.0		1779.7		_	1779.7
16	-	1644.1		1680 .3	1662.2	-	_	1698.0	1720.2	1709.1
17	-	1649.9		_	1649.9	Antonia	_	1703.9	- ,	1703.9
hß		1658.5			1658.5	-	_	1712.8		1712.8
,19	_	1632.5			1632 .5	-	-	1686.1	-	1686.1
20		·	4 -	٣	o	H	_		_	
21	-				ph -104	-	-			
92	- Services				armaters.	Prince	The contract of the contract o	genelité	_	_



			ACCORDING TO MANOUVRIER'S TABLES								
No.	Ski No.	Sex	Humerus (a)	Radius (b)	Ulna (c)	Femur (d)	Tibia (e)	Fibula (f)	Average		
CEM	ETERY H (PEN BU	URIAL)								
23	H 306 (a)	F	155600	1635.00		gar-many.	1600.66	-	1597.22		
24	H 488	F	1285.00	1585.25	1601.38	1462.50	1531.00	_	1523.03		
2 5	H 501 (a)	F	1552.10			- Application	Destroyle	No. ordered	1552.10		
26	H 710	F	1491.00	_	1612.00	15 50 .43	1556.00	1543.00	1550.49		
EME	ETERY H (JA	AR BURI	AL)								
27	H 601 (1)	M			1601.25	-	-	_	1601.25		
28	H 61	F			-	-	1640.00	-	1640.00		
29	H 206 (c)	F	1619.50			-	-	_	1619.50		
30	H 206 (d)	F	1494.00	_	-	_			1494.00		

			PEARSON'S SSION FOR	MULAE'	DUPERTUIS & HADDEN'S 'GENERAL FORMULAE'					
No.	Femur (a)	Humerus (b)	Tibia (c)	Radius (d)	Average	Femur	Tibia (b)	Humerus (c)	Radius (d)	A verage
CEMET	ERY H	OPEN BU	JRIAL)		*					
23		1535.4	1566.2		1550.8	-	1607.2	1586.7		1597.0
24	1465.6		1495.7	1551. 0	1504.1	1492.3	1:31.2	_	1591.6	1538.4
25	-	1535.4			1535.4		-	1586.7	_	1586.7
26		1477.6	1509.8		1493.7		1546.4	1520.7		1533.6
Семеті	ERY H (JAR BUR	(AL)							
27		_	_		_				-	_
28	-	_	1592.1	_	1592.1		1635.1	_	_	1635.1
29	-	15 9 3.3			15 9 3.3		_	1652.7	_	1652.7
30		1483.1		nicement.	1483.1	-	-	1526.9	-	1526.9

22.4.23